

# DEUTERIUM IN THE UNIVERSE

All the heavy hydrogen in space may have been made in the first 15 minutes after the "big bang." Observations leading to estimates of its abundance thus provide evidence on conditions at that time

by Jay M. Pasachoff and William A. Fowler

The calculations of modern cosmology lead to the conclusion that the universe originated between 10 and 20 billion years ago. The particular cosmology that seems to have the greatest weight of observational evidence behind it is the "big bang" theory, which posits that all the matter in the universe was compressed into a superdense kernel that exploded and has been expanding ever since. Within the past few years evolving techniques of observation have made it possible to look backward in space and time to what appear to be the first few minutes after the big bang, and to learn about the physical state and early evolution of the universe.

One item of observational evidence in favor of the big-bang theory was the discovery that the universe is permeated uniformly by radiation corresponding to what would be radiated by a theoretically perfect "black body" with a temperature of three degrees Kelvin (degrees Celsius above absolute zero). This radiation is believed to be a remnant of the original big bang. The newest evidence has come to the fore within the past two years. It is the detection in interstellar space of atoms that may have been formed within the first 15 minutes after the big bang. These atoms are the atoms of deuterium, commonly known as heavy hydrogen.

The big-bang theory stemmed from the discovery by Edwin P. Hubble 50 years ago that all other galaxies appear to be receding from ours, and that the most distant galaxies seem to be receding the fastest. The big-bang theory accounts for Hubble's observations by the expansion of the universe, as does the steady-state cosmology advanced by Hermann Bondi and Thomas Gold and independently by Fred Hoyle. The

that the universe has always looked the same as it does today, and it requires that hydrogen be continuously created to make up for the decreasing density of the universe caused by its expansion. Most astronomers believe this explanation has been ruled out by the discovery of the three-degree background radiation. A variant big-bang cosmology posits that the universe will stop expanding and then contract. The recent measurements of interstellar deuterium have provided a method for determining whether that will happen or whether the universe will simply go on expanding forever.

The "standard" big-bang theory states that in the first 100 seconds after the initial explosion of the universe only protons, neutrons, electrons, positrons and various types of neutrinos existed and that each particle was independent. As the exploding universe cooled, the protons and neutrons began to combine. Since the nucleus of an atom of "ordinary" hydrogen is simply a proton, the combinations of protons and neutrons were the nuclei of the heavier isotopes of hydrogen and of the other atoms. The simplest compound nucleus is the deuterium, composed of one neutron and one proton, which with a single electron forms an atom of deuterium. The nucleus of the third isotope of hydrogen, tritium, consists of one proton and two neutrons. The combination of two protons and one neutron forms the nucleus of an atom having a quota of two electrons; it is the light isotope of helium. The nucleus of the common isotope of helium has two protons and two neutrons. The nuclei of heavier elements are built up in the same way.

In 1948 Ralph A. Alpher, Hans Bethe and George Gamow first calculated the

elements could have been synthesized in the big bang. In 1957 one of us (Fowler), together with Geoffrey Burbidge, E. Margaret Burbidge and Hoyle, laid out a blueprint showing how both the lighter and the heavier elements could have been built up in the interior of stars. Seven years later Hoyle and Robert J. Tayler calculated that the large amount of helium observed throughout the universe (between 20 and 30 percent of the total mass) could not have been manufactured in ordinary stars. The nuclei require a temperature of 10 billion degrees K. for their synthesis, a temperature that could have been attained only in the big bang, in the explosion of supermassive stars ("little bangs") or in supernovas, the explosions of ordinary stars.

In 1966, after the discovery of the three-degree background radiation, P. J. E. Peebles tried to calculate what the relative abundance of deuterium and helium would have been if those two elements had been formed in the big bang. At the same time Robert V. Wagoner joined Fowler and Hoyle to check whether or not the new data sufficiently changed the outcome of previous calculations enough to allow heavy elements to have been synthesized in the big bang. They found that it was still not possible. Could these elements have been formed in events other than the big bang? Let us deal first with their synthesis in the big bang and then discuss the possible exceptions.

The calculations of Wagoner, Fowler and Hoyle show that there is only one isotope left over that is unique to the big bang. It is deuterium. Although helium would also have been made in the big bang, additional helium has been synthesized since then in stars and cannot now be distinguished from the original


helium. Deuterium, however, is only depleted by the processes that go on inside ordinary stars; it is "cooked" into heavier elements by thermonuclear reactions. Therefore whatever deuterium we find now may well have persisted since the origin of the universe. The deuterium would have been formed in the first 1,000 seconds after the big bang, when the rapidly expanding universe had cooled off just enough to allow protons and neutrons to combine. In some sense,

when, deuterium may actually allow us to look backward 10 to 20 billion years in time.

In order to use deuterium as a key to gaining a fuller understanding of the big bang, it is necessary to determine how abundant deuterium is with respect to ordinary hydrogen. Ideally we should like the cosmic abundance ratio, if it exists as a meaningful constant, to be uncomplicated by processes inside stars or

by the chemical interactions of elements and compounds on planets or on grains of ice or dust in space. The ratio is best measured in the gas between the stars.

The density of the interstellar gas is very low: about one atom per cubic centimeter. The space between the stars is so vast, however, that the total amount of gas is very large. It is well known that almost all this gas is hydrogen. The problem is to find out if any of the hydrogen is deuterium and if so how much.



WIDE-FIELD PHOTOGRAPH OF THE MILKY WAY, made by Wulfhard Schlosser of the Ruhr-Universität at Bochum in Germany, shows the region of the center of the galaxy in the direction of which the authors and their colleagues conducted their search at

radio wavelengths for absorption from interstellar deuterium. Galactic center is in the constellation Sagittarius, just above knot of stars to right of center. Round dark object is secondary mirror of astronomical camera. Three linear forms are the mirror supports.

ISOTOPES OF HYDROGEN AND HELIUM differ in the number of their nuclear particles. The lightest isotope is ordinary hydro-

gen (left), which has a single proton in its nucleus (dark color) and a single electron (dark gray). Deuterium has a proton and

neutron (l  
one neutro

How can deuterium be observed? In an atom of ordinary hydrogen the single proton and single electron act as though they are spinning like tiny tops. The spin of the electron can be either in the same direction as the spin of the proton or in the opposite direction. If the spins are in the same direction, the atom is in a higher energy state, and the electron spin can flip over so that it is spinning in the opposite direction with respect to the proton. This spin-flip emits a small amount of energy at the radio wavelength of 21 centimeters. Gaseous hydrogen can also absorb energy at this wavelength as well as emit it. The atoms that absorb the energy are thus put into the higher energy state. Although the spin-flip for any one atom is rare, there are enough atoms in the galaxy for the radiation at 21 centimeters to be quite strong.

Deuterium also has a single electron, and the radiation from its spin-flip is at a wavelength of 92 centimeters. Deuterium, however, is much less abundant than ordinary hydrogen. In the water of the earth's oceans there are 6,600 atoms of ordinary hydrogen for every atom of deuterium. The abundance of deuterium in the solid body of the earth could be quite different. Thus the value for the oceans does not necessarily reflect the abundance ratio of deuterium to ordinary hydrogen on the earth, much less the ratio in interstellar space.

Soon after the spectral line of hydrogen at 21 centimeters was discovered, several astronomers began to search for the line of deuterium in interstellar space. The surveys culminated 10 years ago when Sander Weinreb of the National Radio Astronomy Observatory (NRAO) at Green Bank, W.Va., turned the observatory's 85-foot radio telescope to the constellation Cassiopeia to look for the 92-centimeter line in the spec-

trum of the strong radio source Cassiopeia A. Although Weinreb observed for weeks, he was totally unable to detect deuterium. He stated on the basis of his negative observations that the ratio of deuterium to ordinary hydrogen in that direction appeared to be less than 1 : 13,000, since any larger amount would have been detected. That upper limit is half the ratio of deuterium to ordinary hydrogen on the earth.

Radio astronomy has continued its rapid development since the time of Weinreb's survey, and within the past few years many complex molecules have been detected in the clouds of gas and dust in interstellar space. Most of the molecules have been found at relatively short wavelengths, from about 21 centimeters down to two millimeters. Few spectral lines have been discovered at wavelengths longer than a meter, although the failure has not been for lack of trying. In 1969 Carl A. Gottlieb, Dale F. Dickinson and one of us (Pasachoff) used the 150-foot radio telescope of the Air Force Cambridge Research Laboratories to search at meter wavelengths for various molecules, some of which included deuterium as a constituent. Two such molecules were the hydroxyl radical (OH), with deuterium in place of ordinary hydrogen, and water, with deuterium in place of one of the two ordinary hydrogens. The search was later continued at the NRAO in collaboration with David Buhl, Patrick Palmer, Lewis E. Snyder and Ben Zuckerman. No molecules were detected.

The chance of finding the faint spectral line of deuterium, or for that matter any other spectral line at such wavelengths, seemed very small. Nevertheless, in conversations in 1970 the authors of this article felt that another attempt

should be made. The project would call for many weeks of observing, but it seemed that the 130-foot radio telescope of the Owens Valley Radio Observatory of the California Institute of Technology might be available. Diego A. Cesarsky and Alan T. Moffet of the staff at Owens Valley joined one of us (Pasachoff) in making the observations. In the following discussion of these observations "we" refers to Cesarsky, Moffet and Pasachoff and students from Williams College and Cal Tech who worked with us.

We decided to observe for two weeks at a time because we wanted to analyze the results as we went along. Many problems could have interfered with the investigation. For example, the 92-centimeter line lies in a region of military air-to-ground communications. Although the aircraft transmissions were sometimes bothersome, we were able to remove those time periods from our data.

The first two weeks of observing were in March, 1972. The main direction in which we chose to observe was toward the center of our galaxy. The galactic center, however, is in the southern sky, and at the latitude of the Owens Valley it is above the horizon for only about six hours a day. Since radio telescopes can operate in broad daylight, we were able to observe the galactic center whenever it was above the horizon, to observe in the direction of the Great Nebula in Orion much of the rest of the time and to observe in the direction of Cassiopeia A during any intervening hours. The data from this first observing run looked good, showing that the telescope and the receiving system were working satisfactorily. Furthermore, there was actually a suspicion of a faint absorption line at 92 centimeters.

That summer we had three more observing runs of two weeks each, extend-

ing the re  
concentrated  
the galac  
gen in fr  
other par  
working  
ratio of  
gen is co  
so that  
amount  
same di

SPIN ( of ordi charac droger same c

HELIUM 3



HELIUM 4



(black), tritium has a proton and two neutrons. Helium 3 has two protons and one neutron in its nucleus and two electrons. Helium 4 has two protons and two neutrons.


results of the first run. We continued our efforts in the direction of the galactic center. There is more hydrogen in front of that source than in any other part of the sky. We assumed as a hypothesis that the abundance of deuterium to ordinary hydrogen is constant throughout the universe, presumably the greatest total abundance of deuterium would lie in the direction as the greatest total

amount of ordinary hydrogen. The galactic center is a strong source of radio waves, and the gas in the 40,000 light-years between it and the earth absorbs some of the radiation at the wavelengths of 21 and 92 centimeters.

Our analysis of the data by computer during that summer and fall strengthened our belief that we were actually observing the absorption line of deuterium. In the summer of 1973 five more weeks

of observing by Cesarsky and Pasachoff yielded results that were compatible with the findings in 1972. We attempted to observe over a wider range of wavelengths, but it seemed as though we were now encountering too much outside interference in the wider band. In the narrower range of wavelengths observed in common during both years, the largest absorption fluctuation again corresponded to the spectral line of deuterium at 92 centimeters.

The line is still only barely visible in the data, which show expected random fluctuations [see top illustration on page 113]. Adding the results of both years' observations together has slightly improved the ratio of the signal to the noise and the line seems more than three times as strong as the fluctuations. Even if the "line" turns out to be only a particularly large random fluctuation in the data, the magnitude of the surrounding noise places an important upper limit on how strong the absorption could be. From our data that upper limit is one part in 3,000. If the feature we observed is the deuterium line, then the abundance ratio of deuterium to ordinary




OF THE ELECTRON FLIPS on rare occasions in an atom of ordinary hydrogen or deuterium, with the emission of radiation characteristic of the atom. In the schematic atom of ordinary hydrogen (top left) the nucleus and the electron are spinning in the same direction. In the second drawing at top right the electron has

flipped so that it is spinning in the opposite direction, with the emission of radiation at a wavelength of 21 centimeters. The same kind of spin-flip in the deuterium atom (third and fourth drawings) gives rise to 92-centimeter radiation. When the atoms absorb radiation of same wavelength, the spin of their electron flips back.




in addition to the facts of physics. Chemical combinations of the elements are complicated principles and processes many of which are unknown. That is particularly true for elements coming on the surface of dust grains in interstellar clouds, which is a leading theory for the way in which such molecules form. The ratio of deuterium to ordinary hydrogen in interstellar space could very well be quite different from the ratio of DCN to HCN. Calculations show that the difference in the abundance from such chemical fractionation could be on the order of a factor of 600. In the observation of Jefferts, Penzias and Wilson was adjusted to take the fractionation factor into account, the ratio of deuterium to ordinary hydrogen in the Kleinmann-Low nebula was found to be 1:100,000, just below the range inferred from the observations of Cerny, Moffet and Pasachoff. The fact that the ratio of deuterium to hydrogen is indeed substantially lower than the ratio of DCN to HCN has since been confirmed by two independent limits on the abundance of deuterium by sky observing at radio wavelengths by another group observing a transit of deuterium at visible wavelengths. The abundance ratio of deuterium to ordinary hydrogen has also been determined from space, from the third Orbiting Astronomical Observatory satellite named *Copernicus*. One experiment conducted by a group from Princeton University including Lyman Spitzer, Jr., Jerry F. Drake, Edward B. Easley, Donald C. Morton, John B. Peterson, Jr., and Donald G. York, uses the 36-foot radio telescope to observe spectra at ultraviolet wavelengths between 950 and 1,450 angstroms and between 1,650 and 2,000 angstroms. The second range includes spectral lines from molecular hydrogen ( $H_2$ ), detected only once before in an experiment placed aboard a sounding rocket. Molecular hydrogen has not been detected from the earth's atmosphere because it has no lines in the visible region of the spectrum, and its absorption in the ultraviolet region is absorbed by the atmosphere.

Results from the Princeton experiment aboard *Copernicus* showed that whenever the telescope was pointed toward reddened stars (that is, in directions where the interstellar material in front of the stars affects the overall distribution of the energy radiated by the stars favoring the longer—redder—wavelengths), the spectra revealed that at least 10 percent of the intervening mat-



ABSORPTION LINE OF DEUTERIUM shows up as the central dip in observations of the galactic center at a wavelength of 92 centimeters. The observations were made with the 130-foot radio telescope at the Owens Valley Radio Observatory in California. The curve is the sum of results from observations made in 1972 and 1973. The dip is shifted in frequency by a small amount, indicating that the gas incorporating the deuterium is moving at a velocity of 3.7 kilometers per second toward the solar system. Spectrum is calibrated in terms of temperature of antenna, equivalent to energy of radiation received by antenna.



EMISSION LINE OF DEUTERATED HYDROGEN CYANIDE (DCN), which is actually composed of three separate lines (arrows), was observed by Keith B. Jefferts, Arno A. Penzias and Robert W. Wilson of the Bell Laboratories at millimeter wavelengths with the 36-foot radio telescope of the National Radio Astronomy Observatory on Kitt Peak in Arizona. The molecule was detected in Kleinmann-Low nebula in constellation Orion.

ter was in the form of molecular hydrogen. The same result was obtained with 11 different stars. For a similar number of unreddened stars, however, molecular hydrogen was not detected at all to a limit of one part in 10 million.

Furthermore, the Princeton group measured two lines of HD (deuterated  $H_2$ ) at wavelengths of 1.054 and 1.066 angstroms. The first results, for nine stars, indicate that the abundance ratio of HD to  $H_2$  is 1 : 1,000,000. Again the result had to be adjusted by calculations of chemical combinations to yield a corrected ratio of deuterium to ordinary hydrogen. John H. Black and Alexander Dalgarno of the Center for Astrophysics of the Harvard College Observatory and the Smithsonian Astrophysical Observatory have calculated that the ratio of deuterium to ordinary hydrogen is between 1 : 5,000 and 1 : 500,000 for the nebula Zeta Ophiuchus.

After these promising measurements had been made Rogerson and York attempted to use the telescope aboard *Copernicus* to observe the Lyman series of transitions of the deuterium atom. These transitions absorb radiation in the ultraviolet region of the spectrum when the electron in the deuterium atom is raised to a higher energy level from the "ground" state. The wavelengths of the transitions are slightly shorter than the wavelengths of the corresponding

transitions for the atom of ordinary hydrogen (and were also slightly shorter than the wavelengths for which the experiment was actually designed). The calculation of the ratio of deuterium to ordinary hydrogen from these measurements is straightforward and not subject to the correction factors that must be applied to the molecular observations.


Rogerson and York searched for the transitions in the interstellar gas in front of hot stars of spectral Type B. Such stars have few lines in their own spectra and therefore any absorption lines detected would have been formed in interstellar space between the star and the earth. In the direction of the star Beta Centauri, Rogerson and York were able to measure four lines in the Lyman series for deuterium. The ratio of deuterium to ordinary hydrogen in that direction is 1 : 70,000, with an error of 15 percent. The work is being continued for other stars.

The big-bang theory assumes that the universe is isotropic (that it looks more or less the same in all directions), so that a knowledge of how uniformly deuterium is distributed through space bears on whether or not the big bang is indeed a good model for the origin of the universe. If the abundance of deuterium is found to be nonuniform, for example by comparing the observations from *Copernicus* of nearby stars with our ob-

servations toward the galactic center, then either the big bang is not a good model or the deuterium was formed in some way other than the big bang.

In order for the measurements of deuterium to be significant in cosmological calculations, the measurements must reflect a general cosmic abundance and not simply local variations. For this reason it is difficult to interpret the abundance of deuterium in the solar system in terms of a big-bang origin. The abundance should nonetheless be discussed briefly if only because deuterium had not been detected elsewhere in the solar system until last year.

Some deuterium has been found in the water content of carbonaceous meteorites. Until the Apollo 11 astronauts landed on the moon, meteorites were the only samples of matter from space. One of the Apollo experiments was designed to capture ionized atoms from the solar wind: the flux of ions expelled from the sun. The astronauts caught the ions in a "window shade" of aluminum foil that unrolled on the moon. The foil was then brought back to earth for analysis. Johannes Geiss of the University of Bern and Hubert Reeves of the Nuclear Research Center at Saclay and the Institut of Astrophysics in Paris made deductions about the abundance of deuterium from the light isotope helium 3 that has



SPECTRUM OF THE STAR BETA CENTAURI in the ultraviolet region shows Lyman-gamma absorption lines for ordinary hydrogen (HI) and deuterium (DI). The lines are from the interstellar gas between the star and the solar system and they are superposed

on a broad Lyman-gamma line from the star (*gentle dip*). They were detected with the ultraviolet spectrometer aboard the third Orbiting Astronomical Observatory, named *Copernicus*. A line for oxygen (OI) also happens to be in this region of the spectrum.

found in the foil. It is known that the sun has transformed into helium 3 out of the deuterium that was present in the primordial solar nebula, the cloud of dust and gas out of which the sun formed. Therefore the amount of helium in the foil directly reflects an upper limit for the amount of deuterium that would have been present in the solar nebula. Geiss and Reeves have calculated that the abundance ratio of deuterium to ordinary hydrogen in the solar nebula was 1 : 40,000.

It is also possible that the deuterium in the material from which the planets were made was not transformed into helium 3 as the protosun materialized out of the solar nebula. In March, 1972, Reinhard Beer of the Jet Propulsion Laboratory of the California Institute of Technology and his collaborators reported that they had detected  $\text{CH}_3\text{D}$ , a deuterated form of methane ( $\text{CH}_4$ ), in the spectrum of Jupiter at infrared wavelengths. Beer and Frederic W. Taylor calculated the molecular-correction factors and found that the abundance ratio of deuterium to ordinary hydrogen in Jupiter was between 1 : 13,000 and 1 : 35,000, depending on certain assumptions about the structure of the planet's atmosphere.

Although the interpretation of results from such a relatively complicated molecule as  $\text{CH}_3\text{D}$  is very uncertain, the calculations seem to show a clear discrepancy with the value of 1 : 6,600 found on the earth. Within the past year John Trauger and Frederick L. Roesler of the University of Wisconsin and Nathaniel P. Carleton and Wesley A. Traub of the Center for Astrophysics have detected three lines of deuterated molecular hydrogen ( $\text{HD}$ ) on Jupiter in the far-infrared region of the spectrum. From their observations they calculate a ratio of deuterium to ordinary hydrogen of 1 : 48,000, with a possible error of 20 percent. The discrepancy between the values on the earth and those on Jupiter might be due to the fact that the ratio of heavy water ( $\text{HDO}$ ) to ordinary water ( $\text{H}_2\text{O}$ ) on the earth could be enhanced over the ratio of deuterium to ordinary hydrogen by some chemical fractionation process.

A most unusual observation of deuterium was made two years ago. A group of workers from the University of New Hampshire were observing the sun during the large solar flares of August, 1972, with a gamma ray instrument aboard orbiting Solar Observatory 7. During the particularly large flare of August 7 they detected a peak in the gamma ray




KLEINMANN-LOW NEBULA is embedded in Great Nebula in Orion. Location, which is indicated by contours, is rich in many molecules, including deuterated hydrogen cyanide.

spectrum corresponding to the energy that would have been released by the formation of deuterons. That peak meant deuterium was being synthesized on the surface of the sun. This event, however, is probably unrelated to the total cosmic abundance of deuterium. The amount of deuterium formed in this way is small, and it may well be quickly consumed by further nuclear processes on the sun before it can be expelled into space.

The results of the observations of deuterium to date are summarized in the table on page 112. What do these indications of the amount of deuterium in interstellar space tell us about the early universe? One can link the ratio of deuterium to ordinary hydrogen to the big bang by theoretical calculations. For this purpose we shall adopt the standard model of the big bang developed by Wagoner, Fowler and Hoyle, which accepts three basic assumptions about the universe today: first, the universe is

homogeneous and isotropic; second, the principle of equivalence must hold, that is, a gravitational field cannot be distinguished from an acceleration (a principle fundamental to the theory of relativity); third, the three-degree background radiation was indeed generated by the big bang. The model also makes several other assumptions. The baryon number must be positive, that is, the amount of antimatter is not equal to the amount of matter, ruling out theories to the contrary. The lepton number is small, that is, the flux of neutrinos does not overwhelm the amount of radiation. The general theory of relativity provides the correct theory of gravitation. Lastly, only the kinds of subatomic particles that we now know about are present; there are no new ones.

In the standard model it happens that the amount of deuterium formed just after the big bang is sensitive to the deu-



- DEUTERIUM
- HELIUM 3
- HELIUM 4
- LITHIUM 6
- LITHIUM 7
- · - · - BORON 11
- · · · · ALL OTHERS - MASS 12

"STANDARD" MODEL of the big-bang theory of the origin of the universe predicts an abundance of deuterium that is very sensitive to the postulated density of the universe at the time it originated, as is shown in this diagram devised by Wagoner. The cosmic abundances of a number of other light elements are given as well. Various kinds of observations indicate that the density of the universe is approximately  $10^{-31}$  gram per cubic centimeter.

ity of the  
density we  
few second  
terium jus  
quickly co  
result wou  
venely, if  
the end re  
deuterium  
stoppage].  
of the deut  
is possible  
verse at th  
first synth.  
of the univ  
pansion ne  
what the  
present. T  
our observ  
gram per c  
is not enc  
which will  
definitely.  
verse's pr  
draw conc  
and event  
In addi  
the big ba  
new hypo  
bootstrap  
Carditz of  
Steven C.  
Werner  
Bonn. It  
baryons"  
The supe  
the same  
their mas  
greater. A  
becomes  
forces tha  
clear parti  
The st  
sumes the  
particles t  
because t  
them can  
tory. The  
laws gov  
particles,  
in the lab  
400 times  
rocs. hold  
stars of 1  
postulate  
known pa  
standard  
there are  
prior reas  
other.


If the st  
initial  
compar  
density

of the universe at that time. If the density were relatively high in the first few seconds after the big bang, the deuterium just formed would have been quickly cooked into helium and the end result would be less deuterium. Conversely, if the universe was less dense, the end result would have been more deuterium [see illustration on the opposite page]. Thus from the measured value of the deuterium in the universe today it is possible to find the density of the universe at the time that the elements were first synthesized. Knowing the density of the universe then and the rate of expansion now, it is possible to calculate what the density of the universe is at present. The density we deduce from our observations is of the order of  $10^{-31}$  gram per cubic centimeter. That density is not enough to "close" the universe, which will thus continue to expand indefinitely. Therefore knowing the universe's present density enables us to draw conclusions about its present state and eventual future.

In addition to the standard model of the big bang there is a very speculative hypothesis called the statistical-bootstrap model, advanced by Robert D. Carlitz of the University of Chicago, Steven C. Frautschi of Cal Tech and Werner Nahm of the University of Bonn. It assumes that massive "superbaryons" formed just after the big bang. The superbaryons were approximately the same size as ordinary baryons, but their mass was as much as  $10^{38}$  times greater. At that point the force of gravity comes competitive with the normal forces that govern the interaction of nuclear particles.

The statistical-bootstrap model assumes that there are many elementary particles that have not yet been observed because the energies required to make them cannot be attained in the laboratory. The model also assumes that the laws governing the existence of such particles, which have been determined in the laboratory for particles with up to 10 times the mass of an ordinary baryon, hold true for a further factor in powers of  $10^{37}$ . In one sense this new hypothesis, in which there are many unknown particles, is the opposite of the standard theory, which assumes that there are none. There is, in fact, no a priori reason to favor one theory over the other.

If the statistical-bootstrap model turns out to be accurate, there will be many consequences for cosmology. First, no helium would have been formed in the



"STATISTICAL BOOTSTRAP" MODEL of the big-bang theory, advanced by Robert D. Carlitz of the University of Chicago, Steven C. Frautschi of the California Institute of Technology and Werner Nahm of the University of Bonn, predicts a relation between the abundance of deuterium (white line) and the density of the universe quite different from what is predicted by the standard model. At the density of  $10^{-31}$  gram per cubic centimeter the abundance of helium 3 (thin black line) and of ordinary helium (thick black line) is considerably less than is predicted by the standard model or observed. The density of  $10^{-31}$  gram per cubic centimeter is insufficient to "close" the universe, that is, to stop its expansion.

early universe because no nuclear particles would have been available for a full year after the big bang, and by that time the density of the universe would have been too low for helium to be synthesized. Wagoner has calculated the effect of the statistical-bootstrap model on the formation of deuterium. Contrary to the predictions of the standard model, the statistical-bootstrap model predicts that a less dense universe would result in a lower abundance of deuterium [see illustration above].

There are two other ways to measure the density of the universe. The first entails simply adding up the masses of everything that can be observed (stars, galaxies and so forth) and dividing the total mass in a given unit volume by that volume. The most recent work along this line has been done by Stuart L. Shapiro of Princeton. His results depend somewhat on the value accepted for the Hubble constant, that is, on the rate at which the universe is expanding. The

value of the Hubble constant is currently a topic of debate in some circles. The traditional value is 75 kilometers per second per megaparsec, which corresponds to a density of  $10^{-31}$  gram per cubic centimeter. (One megaparsec is 3.3 million light-years.) A newer value measured by Allan R. Sandage of the Hale Observatories and Gustav Tammann of the University of Basel is 55 kilometers per second per megaparsec, which yields three-fourths the density of the former value.

The second method of measuring the density of the universe considers the dynamics of interactions of clusters of galaxies; it predicts a density that is substantially higher than the value obtained by the first method. Both values for the density could be correct if there is a substantial amount of invisible material in the universe. This "missing mass" could be in the form of molecular hydrogen in the intergalactic material, although the measurements from the Copernicus sat-

