

The Limiting Eigenvalue Density for the Ensemble of m -Circulant Matrices

Gene Kopp and Murat Kolođlul; Advisor Steven J. Miller
Williams College

<http://www.williams.edu/go/math/sjmilller/>

SMALL Presentation, August 4, 2010

Classical Random Matrix Theory

Origins of Random Matrix Theory

Classical Mechanics: 3 Body Problem Intractable.

Heavy nuclei (Uranium: 200+ protons / neutrons) worse!

Get some info by shooting high-energy neutrons into nucleus, see what comes out.

Fundamental Equation:

$$H\psi_n = E_n\psi_n$$

H : matrix, entries depend on system

E_n : energy levels

ψ_n : energy eigenfunctions

Origins of Random Matrix Theory

- Statistical Mechanics: for each configuration, calculate quantity (say pressure).
- Average over all configurations – most configurations close to system average.
- Nuclear physics: choose matrix at random, calculate eigenvalues, average over matrices (real Symmetric $A = A^T$, complex Hermitian $\overline{A}^T = A$).

Random Matrix Ensembles

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1N} \\ a_{12} & a_{22} & a_{23} & \cdots & a_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{1N} & a_{2N} & a_{3N} & \cdots & a_{NN} \end{pmatrix} = A^T, \quad a_{ij} = a_{ji}$$

Fix p , define

$$\text{Prob}(A) = \prod_{1 \leq i < j \leq N} p(a_{ij}).$$

This means

$$\text{Prob}(A : a_{ij} \in [\alpha_{ij}, \beta_{ij}]) = \prod_{1 \leq i < j \leq N} \int_{x_{ij}=\alpha_{ij}}^{\beta_{ij}} p(x_{ij}) dx_{ij}.$$

Want to understand eigenvalues of A .

Eigenvalue Distribution

$\delta(x - x_0)$ is a unit point mass at x_0 :

$$\int_{\mathbb{R}} f(x) \delta(x - x_0) dx = f(x_0).$$

To *each* matrix A , attach a probability measure:

$$\mu_{A,N}(x) := \frac{1}{N} \sum_{i=1}^N \delta\left(x - \frac{\lambda_i(A)}{\sqrt{N}}\right)$$

$$\int_{\mathbb{R}} f(x) \mu_{A,N}(x) dx = \sum_{i=1}^N f\left(\frac{\lambda_i(A)}{\sqrt{N}}\right)$$

$$M_n(A, N) := n^{\text{th}} \text{ moment} = \frac{1}{N^{\frac{n}{2}+1}} \sum_{i=1}^N \lambda_i(A)^n = \frac{\text{Trace}(A^n)}{N^{\frac{n}{2}+1}}.$$

Eigenvalue Trace Formula

We want to understand the eigenvalues of A , but it is the matrix elements that are chosen randomly and independently.

Eigenvalue Trace Lemma

Let A be an $N \times N$ matrix with eigenvalues $\lambda_i(A)$. Then

$$\text{Trace}(A^k) = \sum_{n=1}^N \lambda_i(A)^k,$$

where

$$\text{Trace}(A^k) = \sum_{i_1=1}^N \cdots \sum_{i_k=1}^N a_{i_1 i_2} a_{i_2 i_3} \cdots a_{i_k i_1}.$$

Our ensemble:
m-Circulant Matrices

m-Circulant Ensemble

We look at matrices A of the following form, which we call *m*-circulant or *m*-doped palindromic Toeplitz.

$$\begin{pmatrix} b_{1,0} & b_{1,1} & b_{1,2} & \cdots & & b_{1,\frac{N}{2}-1} & b_{1,\frac{N}{2}} & b_{1,\frac{N}{2}-1} & \cdots & b_{1,1} \\ b_{1,1} & b_{2,0} & b_{2,1} & \cdots & & & b_{2,\frac{N}{2}-1} & b_{2,\frac{N}{2}} & \cdots & b_{2,2} \\ b_{1,2} & b_{2,1} & b_{3,0} & \cdots & & & & \ddots & \ddots & b_{3,3} \\ & & & \ddots & & & & & & \vdots \\ & & & & b_{m,0} & b_{m,1} & b_{m,2} & \cdots & & b_{m,m+1} \\ \vdots & \vdots & \vdots & & b_{m,1} & b_{1,0} & b_{1,1} & \cdots & & b_{1,m+2} \\ & & & & b_{m,2} & b_{1,1} & b_{2,0} & \cdots & & b_{2,m+3} \\ & & & & \vdots & \vdots & \vdots & \ddots & & \vdots \\ b_{1,1} & b_{2,2} & b_{3,3} & \cdots & b_{m,m+1} & b_{1,m+2} & b_{2,m+3} & \cdots & & b_{m,0} \end{pmatrix}$$

Matrices are real symmetric, becomes a probability space when we choose the **red** entries independently from a fixed distribution p of mean 0 and variance 1, and fill in the rest of the matrix as per the structure defined.

Averaging

Look at the *expected value* for the moments:

$$\begin{aligned} M_n(N) &:= \mathbb{E}(M_n(A, N)) \\ &= \frac{1}{N^{\frac{n}{2}+1}} \mathbb{E}(\text{Trace}(A^n)) \\ &= \frac{1}{N^{\frac{n}{2}+1}} \sum_{1 \leq i_1, \dots, i_n \leq N} \mathbb{E}(a_{i_1 i_2} a_{i_2 i_3} \cdots a_{i_n i_1}). \end{aligned}$$

As $N \rightarrow \infty$, these moments converge to the moments of the limiting spectral distribution. A bounding arguments involving Chebyshev's inequality and the Borel-Cantelli lemma shows that a "typical" m -circulant matrix of large dimension has an eigenvalue distribution "close" to this limiting density.

Matchings

We rewrite our formula for the moments as

$$M_n(N) = \frac{1}{N^{\frac{n}{2}+1}} \sum_{\sim} \eta(\sim) m_{d_1(\sim)} \cdots m_{d_l(\sim)}.$$

where the sum is over equivalence relations on $\{(1, 2), (2, 3), \dots, (n, 1)\}$. The $d_j(\sim)$ denote the sizes of the equivalence classes, and the m_d the moments of p .


Finally, the coefficient $\eta(\sim)$ is the number of solutions to the system of Diophantine equations:

Whenever $(s, s+1) \sim (t, t+1)$,

- $i_{s+1} - i_s \equiv i_{t+1} - i_t \pmod{N}$ and $i_s \equiv i_t \pmod{m}$, or
- $i_{s+1} - i_s \equiv -(i_{t+1} - i_t) \pmod{N}$ and $i_s \equiv i_{t+1} \pmod{m}$.

Contributing Terms

As $N \rightarrow \infty$, the only terms that contribute to this sum are those in which the entries are matched in pairs and with opposite orientation.



Therefore, the odd moments go to zero as $N^{-1/2}$.

Algebraic Topology

If we think of these pairings as topological identifications, the contributing ones are precisely the ones that give rise to orientable surfaces.




Figure: A three holed torus.

It turns out that the contribution from such a pairing is m^{-2g} , where g is the genus (number of holes) of the surface. The proof is a combinatorial argument involving Euler characteristic.

Computing the Even Moments

Our formula for the even moments becomes

$$M_{2k} = \sum_{g=0}^{\lfloor k/2 \rfloor} \varepsilon_g(k) m^{-2g} + O_k \left(\frac{1}{N} \right),$$

with $\varepsilon_g(k)$ the number of pairings of the edges of a $(2k)$ -gon giving rise to a genus g surface. J. Harer and D. Zagier (1986) gave generating functions for the $\varepsilon_g(k)$. Their results and a bit of analysis yield explicit formulas for the limiting spectral density.

Results

Theorem

The limiting spectral density function $f_m(x)$ of the real symmetric m -circulant ensemble is given by the formula

$$f_m(x) = \frac{e^{-\frac{mx^2}{2}}}{\sqrt{2\pi m}} \sum_{l=1}^m \sum_{s=0}^{l-1} \binom{m}{l} \frac{(2s-1)!!}{(l-1)!} \binom{2(l-1)}{2s} \cdot (mx^2)^{l-1-s} (-1)^s.$$

Results (continued)

Theorem

As $m \rightarrow \infty$, the limiting spectral densities approach the semicircle distribution.

Results (continued)




Figure: Plots for $f_1, f_2, f_4, f_8, f_{16}$ and the semicircle.

Now that we have an explicit formula for the moments

$$M_{2k} = \sum_{g=0}^{k/2} \varepsilon_g(k) m^{-2g}$$

$$\varepsilon_g(k) = \frac{(2k)!}{(k+1)!(k-2g)!} \times \left(\text{Coefficient of } x^{2g} \text{ in } \left(\frac{x/2}{\tanh(x/2)} \right)^{k+1} \right)$$

Now that we have an explicit formula for the moments

$$M_{2k} = \sum_{g=0}^{k/2} \varepsilon_g(k) m^{-2g}$$

$$\varepsilon_g(k) = \frac{(2k)!}{(k+1)!(k-2g)!} \times \left(\text{Coefficient of } x^{2g} \text{ in } \left(\frac{x/2}{\tanh(x/2)} \right)^{k+1} \right)$$

construct char. fn of limiting spectral distribution:

$$\phi(t) = \sum_{k=0}^{\infty} \frac{1}{(2k)!} (it)^{2k} M_{2k}$$

Now that we have an explicit formula for the moments

$$M_{2k} = \sum_{g=0}^{k/2} \varepsilon_g(k) m^{-2g}$$

$$\varepsilon_g(k) = \frac{(2k)!}{(k+1)!(k-2g)!} \times \left(\text{Coefficient of } x^{2g} \text{ in } \left(\frac{x/2}{\tanh(x/2)} \right)^{k+1} \right)$$

construct char. fn of limiting spectral distribution:

$$\begin{aligned} \phi(t) &= \sum_{k=0}^{\infty} \frac{1}{(2k)!} (it)^{2k} M_{2k} \\ &= \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \varepsilon_g(k) m^{-2g} (-t^2)^k / (2k)!. \end{aligned}$$

Using $(2k - 1)!! = \frac{(2k)!}{2^k k!}$, rewrite the characteristic function:

$$\phi(t) = m^{-1} \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k - 1)!!} \frac{(-t^2/2m)^k}{k!}$$

Using $(2k - 1)!! = \frac{(2k)!}{2^k k!}$, rewrite the characteristic function:

$$\phi(t) = m^{-1} \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k-1)!!} \frac{(-t^2/2m)^k}{k!}$$

Now consider two functions

$$F(y) := \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k-1)!!} y^k$$

Using $(2k - 1)!! = \frac{(2k)!}{2^k k!}$, rewrite the characteristic function:

$$\phi(t) = m^{-1} \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k-1)!!} \frac{(-t^2/2m)^k}{k!}$$

Now consider two functions

$$F(y) := \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k-1)!!} y^k$$

which Harer proves in his paper to be

Using $(2k - 1)!! = \frac{(2k)!}{2^k k!}$, rewrite the characteristic function:

$$\phi(t) = m^{-1} \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k - 1)!!} \frac{(-t^2/2m)^k}{k!}$$

Now consider two functions

$$F(y) := \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k - 1)!!} y^k$$

which Harer proves in his paper to be

$$= \frac{1}{2y} \left(\left(\frac{1+y}{1-y} \right)^m - 1 \right)$$

Using $(2k - 1)!! = \frac{(2k)!}{2^k k!}$, rewrite the characteristic function:

$$\phi(t) = m^{-1} \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k - 1)!!} \frac{(-t^2/2m)^k}{k!}$$

Now consider two functions

$$F(y) := \sum_{k=0}^{\infty} \sum_{g=0}^{k/2} \frac{\varepsilon_g(k) m^{k+1-2g}}{(2k - 1)!!} y^k$$

which Harer proves in his paper to be

$$= \frac{1}{2y} \left(\left(\frac{1+y}{1-y} \right)^m - 1 \right)$$

$$G(y) := e^y = \sum_{k=0}^{\infty} y^k / k!.$$

Bibliography



A. Basak and A. Bose, *Limiting spectral distribution of some band matrices*, preprint 2009.



A. Basak and A. Bose, *Balanced random Toeplitz and Hankel matrices*, preprint.



A. Bose, S. Chatterjee and S. Gangopadhyay, *Limiting spectral distributions of large dimensional random matrices*, J. Indian Statist. Assoc. (2003), **41**, 221–259.



A. Bose and J. Mitra, *Limiting spectral distribution of a special circulant*, Statist. Probab. Lett. **60** (2002), no. 1, 111–120.



W. Bryc, A. Dembo, T. Jiang, *Spectral Measure of Large Random Hankel, Markov, and Toeplitz Matrices*, Annals of Probability **34** (2006), no. 1, 1–38.



Persi Diaconis, "*What is a random matrix?*", Notices of the Amer. Math. Soc. **52** (2005) 1348 – 1349.



Persi Diaconis, *Patterns of Eigenvalues: the 70th Josiah Willard Gibbs Lecture*, Bull. Amer. Math. Soc. **40** (2003) 155 – 178.



F. Dyson, *Statistical theory of the energy levels of complex systems: I, II, III*, J. Mathematical Phys. **3** (1962) 140–156, 157–165, 166–175.



F. Dyson, *The threefold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics*, J. Mathematical Phys., **3** (1962) 1199–1215.



L. Erdős, J. A. Ramirez, B. Schlein and H.-T. Yau, *Bulk Universality for Wigner Matrices*, preprint.
<http://arxiv.org/abs/0905.4176>



L. Erdős, B. Schlein and H.-T. Yau, *Wegner estimate and level repulsion for Wigner random matrices*, preprint. <http://arxiv.org/abs/0905.4176>



W. Feller, *Introduction to Probability Theory and its Applications, Volume 2*, first edition, Wiley, New York, 1966.



F. W. K. Firk and S. J. Miller, *Nuclei, Primes and the Random Matrix Connection*, *Symmetry* **1** (2009), 64–105; doi:10.3390/sym1010064.



P. J. Forrester, N. C. Snaith, and J. J. M. Verbaarschot, *Developments in Random Matrix Theory*. In *Random matrix theory*, *J. Phys. A* **36** (2003), no. 12, R1–R10.



G. Grimmett and D. Stirzaker, *Probability and Random Processes*, third edition, Oxford University Press, 2005.



C. Hammond and S. J. Miller, *Eigenvalue spacing distribution for the ensemble of real symmetric Toeplitz matrices*, *Journal of Theoretical Probability* **18** (2005), no. 3, 537–566.



B. Hayes, *The spectrum of Riemannium*, *American Scientist* **91** (2003), no. 4, 296–300.



S. Jackson, S. J. Miller and V. Pham, *Distribution of Eigenvalues of Highly Palindromic Toeplitz Matrices*, <http://arxiv.org/abs/1003.2010>.



D. Jakobson, S. D. Miller, I. Rivin, and Z. Rudnick, *Eigenvalue spacings for regular graphs*. Pages 317–327 in *Emerging Applications of Number Theory (Minneapolis, 1996)*, The IMA Volumes in Mathematics and its Applications, Vol. 109, Springer, New York, 1999.



V. Kargin, *Spectrum of random Toeplitz matrices with band structure*, *Elect. Comm. in Probab.* **14** (2009), 412–421.



N. Katz and P. Sarnak, *Random Matrices, Frobenius Eigenvalues and Monodromy*, AMS Colloquium Publications, Vol. 45, AMS, Providence, RI, 1999.



N. Katz and P. Sarnak, *Zeros of zeta functions and symmetries*, Bull. AMS **36** (1999), 1–26.



J. P. Keating and N. C. Snaith, *Random matrices and L-functions*. In *Random Matrix Theory*, J. Phys. A **36** (2003), no. 12, 2859–2881.



D.-Z. Liu and Z.-D. Wang, *Limit Distribution of Eigenvalues for Random Hankel and Toeplitz Band Matrices*, preprint, 2009.



A. Massey, S. J. Miller, J. Sinsheimer, *Distribution of eigenvalues of real symmetric palindromic Toeplitz matrices and circulant matrices*, Journal of Theoretical Probability **20** (2007), no. 3, 637-662.



B. McKay, *The expected eigenvalue distribution of a large regular graph*, Linear Algebra Appl. **40** (1981), 203–216.



M. L. Mehta, *Random Matrices*, 3rd edition, Elsevier, San Diego, CA (2004)



M. L. Mehta and M. Gaudin, *On the density of the eigenvalues of a random matrix*, Nuclear Physics **18** (1960), 420–427.



L. Takacs, *A Moment Convergence Theorem*, The American Mathematical Monthly **98** (Oct., 1991), no. 8, 742–746.



T. Tao and V. Vu, *From the Littlewood-Offord problem to the Circular Law: universality of the spectral distribution of random matrices*, Bull. Amer. Math. Soc. **46** (2009), 377–396.



T. Tao and V. Vu, *Random matrices: universality of local eigenvalue statistics up to the edge*, preprint.
http://arxiv.org/PS_cache/arxiv/pdf/0908/0908.1982v1.pdf



C. A. Tracy and H. Widom, *Level-spacing distributions and the Airy kernel*, *Commun. Math. Phys.* **159** (1994), 151–174.



C. Tracy and H. Widom, *On Orthogonal and Symplectic Matrix Ensembles*, *Communications in Mathematical Physics* **177** (1996), 727–754.



C. Tracy and H. Widom, *Distribution functions for largest eigenvalues and their applications*, *ICM Vol. I* (2002), 587–596.



E. Wigner, *On the statistical distribution of the widths and spacings of nuclear resonance levels*, Proc. Cambridge Philo. Soc. **47** (1951), 790–798.



E. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions*, Ann. of Math. **2** (1955), no. 62, 548–564.



E. Wigner, *Statistical Properties of real symmetric matrices*. Pages 174–184 in *Canadian Mathematical Congress Proceedings*, University of Toronto Press, Toronto, 1957.



E. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions. II*, Ann. of Math. Ser. 2 **65** (1957), 203–207.



E. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. Ser. 2 **67** (1958), 325–327.



E. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. Ser. 2 **67** (1958), 325–327.



J. Wishart, *The generalized product moment distribution in samples from a normal multivariate population*, *Biometrika* **20 A** (1928), 32–52.



N. C. Wormald, *Models of random regular graphs*. Pages 239–298 in *Surveys in combinatorics, 1999 (Canterbury)* London Mathematical Society Lecture Note Series, vol. 267, Cambridge University Press, Cambridge, 1999.