

Eigenvalue Statistics of Toeplitz and Block *m*-Circulant Ensembles

Gene Kopp[†], Murat Kolođlu[‡], Steven J Miller[‡]
†: University of Chicago; ‡: Williams College

email sjm1@williams.edu

<http://www.williams.edu/Mathematics/sjmiller/>

AMS Special Session on Random Processes
Holy Cross, April 9, 2011

Introduction

Goals

- See how the structure of the ensembles affects limiting behavior.
- Discuss the tools and techniques needed to prove the results.
- Due to time: only discussing eigenvalue density, not neighbor spacings.

Eigenvalue Distribution

To each A , attach a probability measure:

$$\begin{aligned} \mu_{A,N}(\mathbf{x}) &= \frac{1}{N} \sum_{i=1}^N \delta \left(\mathbf{x} - \frac{\lambda_i(A)}{2\sqrt{N}} \right) \\ \int_a^b \mu_{A,N}(\mathbf{x}) d\mathbf{x} &= \frac{\# \left\{ \lambda_i : \frac{\lambda_i(A)}{2\sqrt{N}} \in [a, b] \right\}}{N} \\ \text{k}^{\text{th}} \text{ moment} &= \frac{\sum_{i=1}^N \lambda_i(A)^k}{2^k N^{\frac{k}{2}+1}} = \frac{\text{Trace}(A^k)}{2^k N^{\frac{k}{2}+1}}. \end{aligned}$$

Wigner's Semi-Circle Law

Wigner's Semi-Circle Law

$N \times N$ real symmetric matrices, entries i.i.d.r.v. from a fixed $p(x)$ with mean 0, variance 1, and other moments finite. Then for almost all A , as $N \rightarrow \infty$

$$\mu_{A,N}(x) \longrightarrow \begin{cases} \frac{2}{\pi} \sqrt{1-x^2} & \text{if } |x| \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Fat-Thin Families

Need a family **FAT** enough to do averaging and **THIN** enough so that everything isn't averaged out.

Real Symmetric Matrices have $\frac{N(N+1)}{2}$ independent entries.


Examples of Fat-Thin sub-families:

- Band Matrices
- Random Graphs
- Special Matrices (Toeplitz, Circulant, ...)


McKay's Law (Kesten Measure) with $d = 3$

Density of Eigenvalues for d -regular graphs

$$f(x) = \begin{cases} \frac{d}{2\pi(d^2-x^2)} \sqrt{4(d-1)-x^2} & |x| \leq 2\sqrt{d-1} \\ 0 & \text{otherwise.} \end{cases}$$



McKay's Law (Kesten Measure) with $d = 6$



Fat-Thin: fat enough to average, thin enough to get something different than semi-circle (though as $d \rightarrow \infty$ recover semi-circle).

Toeplitz Ensembles


Toeplitz Ensembles

Toeplitz matrix is of the form

$$\begin{pmatrix} b_0 & b_1 & b_2 & \cdots & b_{N-1} \\ b_{-1} & b_0 & b_1 & \cdots & b_{N-2} \\ b_{-2} & b_{-1} & b_0 & \cdots & b_{N-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ b_{1-N} & b_{2-N} & b_{3-N} & \cdots & b_0 \end{pmatrix}$$

- Will consider Real Symmetric Toeplitz matrices.
- Main diagonal zero, $N - 1$ independent parameters.
- Normalize Eigenvalues by \sqrt{N} .

The Fourth Moment



$$M_4(N) = \frac{1}{N^3} \sum_{1 \leq i_1, i_2, i_3, i_4 \leq N} \mathbb{E}(b_{|i_1 - i_2|} b_{|i_2 - i_3|} b_{|i_3 - i_4|} b_{|i_4 - i_1|})$$

Let $x_j = |i_j - i_{j+1}|$.

The Fourth Moment

Case One: $x_1 = x_2, x_3 = x_4$:

$$i_1 - i_2 = -(i_2 - i_3) \quad \text{and} \quad i_3 - i_4 = -(i_4 - i_1).$$

Implies

$$i_1 = i_3, \quad i_2 \text{ and } i_4 \text{ arbitrary.}$$

Left with $\mathbb{E}[b_{x_1}^2 b_{x_3}^2]$:

$$N^3 - N \text{ times get } 1, \quad N \text{ times get } p_4 = \mathbb{E}[b_{x_1}^4].$$

Contributes 1 in the limit.

The Fourth Moment

$$M_4(N) = \frac{1}{N^3} \sum_{1 \leq i_1, i_2, i_3, i_4 \leq N} \mathbb{E}(b_{|i_1 - i_2|} b_{|i_2 - i_3|} b_{|i_3 - i_4|} b_{|i_4 - i_1|})$$

Case Two: Diophantine Obstruction: $x_1 = x_3$ and $x_2 = x_4$.

$$i_1 - i_2 = -(i_3 - i_4) \quad \text{and} \quad i_2 - i_3 = -(i_4 - i_1).$$

This yields

$$i_1 = i_2 + i_4 - i_3, \quad i_1, i_2, i_3, i_4 \in \{1, \dots, N\}.$$


If $i_2, i_4 \geq \frac{2N}{3}$ and $i_3 < \frac{N}{3}$, $i_1 > N$: at most $(1 - \frac{1}{27})N^3$ valid choices.

The Fourth Moment

Theorem: Fourth Moment: Let p_4 be the fourth moment of p . Then

$$M_4(N) = 2\frac{2}{3} + O_{p_4}\left(\frac{1}{N}\right).$$

500 Toeplitz Matrices, 400×400 .



Main Result

Theorem: Hammond and M– '05

For real symmetric Toeplitz matrices, the limiting spectral measure converges in probability to a unique measure of unbounded support which is not the Gaussian. If p is even have strong convergence).

Theorem: Massey, M– and Sinsheimer '07

For real symmetric palindromic matrices (first row a palindrome), converge in probability to the Gaussian (if p is even have strong convergence).

Results exist for highly palindromic (Jackson, M– and Pham '11).

Block *m*-Circulant Matrices

Study circulant matrices periodic with period m on diagonals.

6-by-6 real symmetric period 2-circulant matrix:

$$\begin{pmatrix} c_0 & c_1 & c_2 & c_3 & c_2 & d_1 \\ c_1 & d_0 & d_1 & d_2 & c_3 & d_2 \\ c_2 & d_1 & c_0 & c_1 & c_2 & c_3 \\ c_3 & d_2 & c_1 & d_0 & d_1 & d_2 \\ c_2 & c_3 & c_2 & d_1 & c_0 & c_1 \\ d_1 & d_2 & c_3 & d_2 & c_1 & d_0 \end{pmatrix}.$$

Look at the *expected value* for the moments:

$$\begin{aligned} M_n(N) &:= \mathbb{E}(M_n(A, N)) \\ &= \frac{1}{N^{\frac{n}{2}+1}} \sum_{1 \leq i_1, \dots, i_n \leq N} \mathbb{E}(a_{i_1 i_2} a_{i_2 i_3} \cdots a_{i_n i_1}). \end{aligned}$$

Matchings

Rewrite:

$$M_n(N) = \frac{1}{N^{\frac{n}{2}+1}} \sum_{\sim} \eta(\sim) m_{d_1(\sim)} \cdots m_{d_l(\sim)}.$$

where the sum is over equivalence relations on $\{(1, 2), (2, 3), \dots, (n, 1)\}$. The $d_j(\sim)$ denote the sizes of the equivalence classes, and the m_d the moments of p .

Finally, the coefficient $\eta(\sim)$ is the number of solutions to the system of Diophantine equations:

Whenever $(s, s+1) \sim (t, t+1)$,

- $i_{s+1} - i_s \equiv i_{t+1} - i_t \pmod{N}$ and $i_s \equiv i_t \pmod{m}$, or
- $i_{s+1} - i_s \equiv -(i_{t+1} - i_t) \pmod{N}$ and $i_s \equiv i_{t+1} \pmod{m}$.

- $i_{s+1} - i_s \equiv i_{t+1} - i_t \pmod{N}$ and $i_s \equiv i_t \pmod{m}$, or
- $i_{s+1} - i_s \equiv -(i_{t+1} - i_t) \pmod{N}$ and $i_s \equiv i_{t+1} \pmod{m}$.





Figure: Red edges same orientation and blue, green opposite.


Contributing Terms

As $N \rightarrow \infty$, the only terms that contribute to this sum are those in which the entries are matched in pairs and with opposite orientation.



Algebraic Topology

Think of pairings as topological identifications, the contributing ones give rise to orientable surfaces.



Contribution from such a pairing is m^{-2g} , where g is the genus (number of holes) of the surface. Proof: combinatorial argument involving Euler characteristic.

Computing the Even Moments

Theorem: Even Moment Formula

$$M_{2k} = \sum_{g=0}^{\lfloor k/2 \rfloor} \varepsilon_g(k) m^{-2g} + O_k \left(\frac{1}{N} \right),$$

with $\varepsilon_g(k)$ the number of pairings of the edges of a $(2k)$ -gon giving rise to a genus g surface.

J. Harer and D. Zagier (1986) gave generating functions for the $\varepsilon_g(k)$.

Harer and Zagier

$$\sum_{g=0}^{\lfloor k/2 \rfloor} \varepsilon_g(k) r^{k+1-2g} = (2k-1)!! c(k, r)$$

where

$$1 + 2 \sum_{k=0}^{\infty} c(k, r) x^{k+1} = \left(\frac{1+x}{1-x} \right)^r.$$

Thus, we write

$$M_{2k} = m^{-(k+1)} (2k-1)!! c(k, m).$$

A multiplicative convolution and Cauchy's residue formula yields the *characteristic function* of the distribution (inverse Fourier transform of the density).

$$\begin{aligned}
 \phi(t) &= \sum_{k=0}^{\infty} \frac{(it)^{2k} M_{2k}}{(2k)!} \\
 &= \frac{1}{2\pi im} \oint_{|z|=2} \frac{1}{2z^{-1}} \left(\left(\frac{1+z^{-1}}{1-z^{-1}} \right)^m - 1 \right) e^{-t^2 z/2m} \frac{dz}{z} \\
 &= \frac{1}{m} e^{-\frac{t^2}{2m}} \sum_{l=1}^m \binom{m}{l} \frac{1}{(l-1)!} \left(\frac{-t^2}{m} \right)^{l-1}
 \end{aligned}$$

Results

Fourier transform and algebra yields

Theorem: Kopp, Kolođlu and M-

The limiting spectral density function $f_m(x)$ of the real symmetric m -circulant ensemble is given by the formula

$$f_m(x) = \frac{e^{-\frac{mx^2}{2}}}{\sqrt{2\pi m}} \sum_{r=0}^m \frac{1}{(2r)!} \sum_{s=0}^{m-r} \binom{m}{r+s+1} \frac{(2r+2s)!}{(r+s)!s!} \left(-\frac{1}{2}\right)^s (mx^2)^r.$$

As $m \rightarrow \infty$, the limiting spectral densities approach the semicircle distribution.

Proof of $m \rightarrow \infty$ Convergence

The characteristic function for the spectral measures of the period m -circulant matrices can be written in terms of the Laguerre polynomial

$$\phi_m(t) = \frac{1}{m} e^{-\frac{t^2}{2m}} L_{m-1}^{(1)}(t^2/m),$$

or equivalently in terms of the confluent hypergeometric function

$$\phi_m(t) = \exp(-t^2/2m) M(m+1, 2, -t^2/m).$$

From 13.2.2 of [AS] we have

$\lim_{m \rightarrow \infty} \phi_m(t) = \phi(t) = J_1(2t)/t$; however, we need some control on the rate of convergence.

Proof of $m \rightarrow \infty$ Convergence (cont)

Let $r > 1/3$ and $\beta = \frac{2}{3}(1 - r)$. For all m and all t we have

$$|\phi_m(t) - \phi(t)| \ll_r \begin{cases} m^{-(1-r)} & \text{if } |t| \leq m^\beta \\ t^{-\frac{3}{2}} + m^{-\frac{5}{4}} \exp(-\frac{t^2}{2m}) & \text{otherwise,} \end{cases}$$

$$|f_m(x) - f_{\text{s.c.}}(x)| \leq \int_{-\infty}^{\infty} |\phi_m(t) - \phi(t)| dt \ll m^{-\frac{1-r}{3}}.$$

$\epsilon > 0$ and $r = \frac{1}{3} + 3\epsilon$ bound integral with $O(m^{-\frac{2}{9} + \epsilon})$.

Key idea in proof: 13.3.7 of [AS]

$$\phi_m(t) = e^{-\frac{t^2}{2m}} M(m+1, 2, -\frac{t^2}{m}) = \frac{J_1(2t)}{t} - \frac{1}{2m} \sum_{n=1}^{\infty} A_n \left(\frac{-t}{2m}\right)^{n-1} J_{n+1}(2t),$$

where $A_0 = 1$, $A_1 = 0$, $A_2 = 1$ and $A_{n+1} = A_{n-1} + \frac{2m}{n+1} A_{n-2}$ for $n \geq 2$.

For any $r > \frac{1}{3}$ have $A_n \ll_r m^m$.

Results (continued)




Figure: Plot for f_1 and histogram of eigenvalues of 100 circulant matrices of size 400×400 .

Results (continued)




Figure: Plot for f_2 and histogram of eigenvalues of 100 2-circulant matrices of size 400×400 .

Results (continued)




Figure: Plot for f_3 and histogram of eigenvalues of 100 3-circulant matrices of size 402×402 .

Results (continued)




Figure: Plot for f_4 and histogram of eigenvalues of 100 4-circulant matrices of size 400×400 .

Results (continued)




Figure: Plot for f_8 and histogram of eigenvalues of 100 8-circulant matrices of size 400×400 .

Results (continued)




Figure: Plot for f_{20} and histogram of eigenvalues of 100 20-circulant matrices of size 400×400 .

Results (continued)




Figure: Plot of convergence to the semi-circle.

Bibliography



A. Basak and A. Bose, *Limiting spectral distribution of some band matrices*, preprint 2009.



A. Basak and A. Bose, *Balanced random Toeplitz and Hankel matrices*, preprint.



A. Bose, S. Chatterjee and S. Gangopadhyay, *Limiting spectral distributions of large dimensional random matrices*, J. Indian Statist. Assoc. (2003), **41**, 221–259.



A. Bose and J. Mitra, *Limiting spectral distribution of a special circulant*, Statist. Probab. Lett. **60** (2002), no. 1, 111–120.



W. Bryc, A. Dembo, T. Jiang, *Spectral Measure of Large Random Hankel, Markov, and Toeplitz Matrices*, Annals of Probability **34** (2006), no. 1, 1–38.



Persi Diaconis, “*What is a random matrix?*”, Notices of the Amer. Math. Soc. **52** (2005) 1348 – 1349.



Persi Diaconis, *Patterns of Eigenvalues: the 70th Josiah Willard Gibbs Lecture*, Bull. Amer. Math. Soc. **40** (2003) 155 – 178.



F. Dyson, *Statistical theory of the energy levels of complex systems: I, II, III*, J. Mathematical Phys. **3** (1962) 140–156, 157–165, 166–175.



F. Dyson, *The threefold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics*, J. Mathematical Phys., **3** (1962) 1199–1215.



L. Erdős, J. A. Ramirez, B. Schlein and H.-T. Yau, *Bulk Universality for Wigner Matrices*, preprint.
<http://arxiv.org/abs/0905.4176>



L. Erdős, B. Schlein and H.-T. Yau, *Wegner estimate and level repulsion for Wigner random matrices*, preprint. <http://arxiv.org/abs/0905.4176>



W. Feller, *Introduction to Probability Theory and its Applications, Volume 2*, first edition, Wiley, New York, 1966.



F. W. K. Firk and S. J. Miller, *Nuclei, Primes and the Random Matrix Connection*, *Symmetry* **1** (2009), 64–105; doi:10.3390/sym1010064.



P. J. Forrester, N. C. Snaith, and J. J. M. Verbaarschot, *Developments in Random Matrix Theory*. In *Random matrix theory*, *J. Phys. A* **36** (2003), no. 12, R1–R10.



G. Grimmett and D. Stirzaker, *Probability and Random Processes*, third edition, Oxford University Press, 2005.



C. Hammond and S. J. Miller, *Eigenvalue spacing distribution for the ensemble of real symmetric Toeplitz matrices*, Journal of Theoretical Probability **18** (2005), no. 3, 537–566.



J. Harer and D. Zagier, *The Euler characteristic of the moduli space of curves*, Invent. Math. **85** (1986), 457–485.



B. Hayes, *The spectrum of Riemannium*, American Scientist **91** (2003), no. 4, 296–300.



S. Jackson, S. J. Miller and V. Pham, *Distribution of Eigenvalues of Highly Palindromic Toeplitz Matrices*, <http://arxiv.org/abs/1003.2010>. 2010.



D. Jakobson, S. D. Miller, I. Rivin, and Z. Rudnick, *Eigenvalue spacings for regular graphs*. Pages 317–327 in *Emerging Applications of Number Theory (Minneapolis, 1996)*, The IMA Volumes in Mathematics and its Applications, Vol. 109, Springer, New York, 1999.



V. Kargin, *Spectrum of random Toeplitz matrices with band structure*, Elect. Comm. in Probab. **14** (2009), 412–421.



N. Katz and P. Sarnak, *Random Matrices, Frobenius Eigenvalues and Monodromy*, AMS Colloquium Publications, Vol. 45, AMS, Providence, RI, 1999.



N. Katz and P. Sarnak, *Zeros of zeta functions and symmetries*, Bull. AMS **36** (1999), 1–26.



J. P. Keating and N. C. Snaith, *Random matrices and L-functions*. In *Random Matrix Theory*, J. Phys. A **36** (2003), no. 12, 2859–2881.



G. Kopp, M. Kolođlu and S. J. Miller, *The Limiting Eigenvalue Density for the Ensemble of m -Circulant Matrices*, preprint.



D.-Z. Liu and Z.-D. Wang, *Limit Distribution of Eigenvalues for Random Hankel and Toeplitz Band Matrices*, preprint, 2009.



A. Massey, S. J. Miller, J. Sinsheimer, *Distribution of eigenvalues of real symmetric palindromic Toeplitz matrices and circulant matrices*, Journal of Theoretical Probability **20** (2007), no. 3, 637-662.



B. McKay, *The expected eigenvalue distribution of a large regular graph*, Linear Algebra Appl. **40** (1981), 203–216.



M. L. Mehta, *Random Matrices*, 3rd edition, Elsevier, San Diego, CA (2004)



M. L. Mehta and M. Gaudin, *On the density of the eigenvalues of a random matrix*, Nuclear Physics **18** (1960), 420–427.



L. Takacs, *A Moment Convergence Theorem*, The American Mathematical Monthly **98** (Oct., 1991), no. 8, 742–746.



T. Tao and V. Vu, *From the Littlewood-Offord problem to the Circular Law: universality of the spectral distribution of random matrices*, Bull. Amer. Math. Soc. **46** (2009), 377–396.



T. Tao and V. Vu, *Random matrices: universality of local eigenvalue statistics up to the edge*, preprint.
http://arxiv.org/PS_cache/arxiv/pdf/0908/0908.1982v1.pdf



C. A. Tracy and H. Widom, *Level-spacing distributions and the Airy kernel*, *Commun. Math. Phys.* **159** (1994), 151–174.



C. Tracy and H. Widom, *On Orthogonal and Symplectic Matrix Ensembles*, *Communications in Mathematical Physics* **177** (1996), 727–754.



C. Tracy and H. Widom, *Distribution functions for largest eigenvalues and their applications*, *ICM Vol. I* (2002), 587–596.



E. Wigner, *On the statistical distribution of the widths and spacings of nuclear resonance levels*, Proc. Cambridge Philo. Soc. **47** (1951), 790–798.



E. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions*, Ann. of Math. **2** (1955), no. 62, 548–564.



E. Wigner, *Statistical Properties of real symmetric matrices*. Pages 174–184 in *Canadian Mathematical Congress Proceedings*, University of Toronto Press, Toronto, 1957.



E. Wigner, *Characteristic vectors of bordered matrices with infinite dimensions. II*, Ann. of Math. Ser. 2 **65** (1957), 203–207.



E. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. Ser. 2 **67** (1958), 325–327.



E. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. Ser. 2 **67** (1958), 325–327.



J. Wishart, *The generalized product moment distribution in samples from a normal multivariate population*, *Biometrika* **20 A** (1928), 32–52.



N. C. Wormald, *Models of random regular graphs*. Pages 239–298 in *Surveys in combinatorics, 1999 (Canterbury)* London Mathematical Society Lecture Note Series, vol. 267, Cambridge University Press, Cambridge, 1999.