

From the Manhattan Project to Number Theory: How Nuclear Physics Helps Us Understand Primes

Steven J Miller
Williams College

Steven.J.Miller@williams.edu
<http://www.williams.edu/go/math/sjmiller/>

Bronfman Science Lunch
Williams College, June 23, 2009

Introduction

Fundamental Problem: Spacing Between Events

General Formulation: Studying system, observe values at t_1, t_2, t_3, \dots

Question: What rules govern the spacings between the t_i ?

Examples:

- Spacings b/w Energy Levels of Nuclei.
- Spacings b/w Eigenvalues of Matrices.
- Spacings b/w Primes.
- Spacings b/w $n^k \alpha \bmod 1$.
- Spacings b/w Zeros of L -functions.

Sketch of proofs

In studying many statistics, often three key steps:

- 1 Determine correct scale for events.
- 2 Develop an explicit formula relating what we want to study to something we understand.
- 3 Use an averaging formula to analyze the quantities above.

It is not always trivial to figure out what is the correct statistic to study!

Background Material: Linear Algebra

Eigenvalue, Eigenvector

Say $\vec{v} \neq \vec{0}$ is an eigenvector of A with eigenvalue λ if
 $A\vec{v} = \lambda\vec{v}$.

Background Material: Linear Algebra

Eigenvalue, Eigenvector

Say $\vec{v} \neq \vec{0}$ is an eigenvector of A with eigenvalue λ if $A\vec{v} = \lambda\vec{v}$.

Example:

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 3 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = -1 \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

Background Material: Probability

Probability Density

A random variable X has a probability density $p(x)$ if

- $p(x) \geq 0$;
- $\int_{-\infty}^{\infty} p(x) dx = 1$;
- $\text{Prob}(X \in [a, b]) = \int_a^b p(x) dx$.

Background Material: Probability

Probability Density

A random variable X has a probability density $p(x)$ if

- $p(x) \geq 0$;
- $\int_{-\infty}^{\infty} p(x) dx = 1$;
- $\text{Prob}(X \in [a, b]) = \int_a^b p(x) dx$.

Examples:

- 1 Exponential: $p(x) = e^{-x/\lambda} / \lambda$ for $x \geq 0$;
- 2 Normal: $p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2}$;
- 3 Uniform: $p(x) = \frac{1}{b-a}$ for $a \leq x \leq b$ and 0 otherwise.

Background Material: Probability (cont)

Key Concepts

- Mean (average value): $\mu = \int_{-\infty}^{\infty} xp(x)dx.$

Background Material: Probability (cont)

Key Concepts

- Mean (average value): $\mu = \int_{-\infty}^{\infty} xp(x)dx.$
- Variance (how spread out): $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 p(x)dx.$

Background Material: Probability (cont)

Key Concepts

- Mean (average value): $\mu = \int_{-\infty}^{\infty} xp(x)dx.$
- Variance (how spread out): $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 p(x)dx.$
- k^{th} moment: $\mu_k = \int_{-\infty}^{\infty} x^k p(x)dx.$

Background Material: Probability (cont)

Key Concepts

- Mean (average value): $\mu = \int_{-\infty}^{\infty} xp(x)dx.$
- Variance (how spread out): $\sigma^2 = \int_{-\infty}^{\infty} (x - \mu)^2 p(x)dx.$
- k^{th} moment: $\mu_k = \int_{-\infty}^{\infty} x^k p(x)dx.$

Key observation

As a nice function is given by its Taylor series, a nice probability density is determined by its moments.

Classical Random Matrix Theory

Origins of Random Matrix Theory

Classical Mechanics: 3 Body Problem Intractable.

Origins of Random Matrix Theory

Classical Mechanics: 3 Body Problem Intractable.

Heavy nuclei (Uranium: 200+ protons / neutrons) worse!

Origins of Random Matrix Theory

Classical Mechanics: 3 Body Problem Intractable.

Heavy nuclei (Uranium: 200+ protons / neutrons) worse!

Get some info by shooting high-energy neutrons into nucleus, see what comes out.

Origins of Random Matrix Theory

Classical Mechanics: 3 Body Problem Intractable.

Heavy nuclei (Uranium: 200+ protons / neutrons) worse!

Get some info by shooting high-energy neutrons into nucleus, see what comes out.

Fundamental Equation:


$$H\psi_n = E_n\psi_n$$

H : matrix, entries depend on system

E_n : energy levels

ψ_n : energy eigenfunctions

Origins (continued)



- Statistical Mechanics: for each configuration, calculate quantity (say pressure).
- Average over all configurations – most configurations close to system average.
- Nuclear physics: choose matrix at random, calculate eigenvalues, average over matrices (real Symmetric $A = A^T$, complex Hermitian $\overline{A}^T = A$).

Random Matrix Ensembles

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1N} \\ a_{12} & a_{22} & a_{23} & \cdots & a_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{1N} & a_{2N} & a_{3N} & \cdots & a_{NN} \end{pmatrix} = A^T, \quad a_{ij} = a_{ji}$$

Random Matrix Ensembles

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1N} \\ a_{12} & a_{22} & a_{23} & \cdots & a_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{1N} & a_{2N} & a_{3N} & \cdots & a_{NN} \end{pmatrix} = A^T, \quad a_{ij} = a_{ji}$$

Fix p , define

$$\text{Prob}(A) = \prod_{1 \leq i \leq j \leq N} p(a_{ij}).$$

Random Matrix Ensembles

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1N} \\ a_{12} & a_{22} & a_{23} & \cdots & a_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{1N} & a_{2N} & a_{3N} & \cdots & a_{NN} \end{pmatrix} = A^T, \quad a_{ij} = a_{ji}$$

Fix p , define

$$\text{Prob}(A) = \prod_{1 \leq i < j \leq N} p(a_{ij}).$$

This means

$$\text{Prob}(A : a_{ij} \in [\alpha_{ij}, \beta_{ij}]) = \prod_{1 \leq i < j \leq N} \int_{x_{ij}=\alpha_{ij}}^{\beta_{ij}} p(x_{ij}) dx_{ij}.$$

Eigenvalue Trace Lemma

Want to understand the eigenvalues of A , but it is the matrix elements that are chosen randomly and independently.

Eigenvalue Trace Lemma

Want to understand the eigenvalues of A , but it is the matrix elements that are chosen randomly and independently.

Eigenvalue Trace Lemma

Let A be an $N \times N$ matrix with eigenvalues $\lambda_i(A)$. Then

$$\text{Trace}(A^k) = \sum_{n=1}^N \lambda_i(A)^k,$$

where

$$\text{Trace}(A^k) = \sum_{i_1=1}^N \cdots \sum_{i_k=1}^N a_{i_1 i_2} a_{i_2 i_3} \cdots a_{i_N i_1}.$$

Eigenvalue Distribution

$\delta(\mathbf{x} - \mathbf{x}_0)$ is a unit point mass at \mathbf{x}_0 :

$$\int_{-\infty}^{\infty} f(\mathbf{x})\delta(\mathbf{x} - \mathbf{x}_0)d\mathbf{x} = f(\mathbf{x}_0).$$

Eigenvalue Distribution

$\delta(\mathbf{x} - \mathbf{x}_0)$ is a unit point mass at \mathbf{x}_0 :

$$\int_{-\infty}^{\infty} f(\mathbf{x}) \delta(\mathbf{x} - \mathbf{x}_0) d\mathbf{x} = f(\mathbf{x}_0).$$

To each A , attach a probability measure:

$$\mu_{A,N}(\mathbf{x}) = \frac{1}{N} \sum_{i=1}^N \delta\left(\mathbf{x} - \frac{\lambda_i(A)}{2\sqrt{N}}\right)$$

$$\int_a^b \mu_{A,N}(\mathbf{x}) d\mathbf{x} = \frac{\#\left\{\lambda_i : \frac{\lambda_i(A)}{2\sqrt{N}} \in [a, b]\right\}}{N}$$

$$\text{k}^{\text{th}} \text{ moment} = \frac{\sum_{i=1}^N \lambda_i(A)^k}{2^k N^{\frac{k}{2}+1}} = \frac{\text{Trace}(A^k)}{2^k N^{\frac{k}{2}+1}}.$$

Density of States

Wigner's Semi-Circle Law

Wigner's Semi-Circle Law

$N \times N$ real symmetric matrices, entries i.i.d.r.v. from a fixed $p(x)$ with mean 0, variance 1, and other moments finite. Then for almost all A , as $N \rightarrow \infty$

$$\mu_{A,N}(x) \longrightarrow \begin{cases} \frac{2}{\pi} \sqrt{1-x^2} & \text{if } |x| \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

SKETCH OF PROOF: Correct Scale

$$\text{Trace}(A^2) = \sum_{i=1}^N \lambda_i(A)^2.$$

By the Central Limit Theorem:

$$\text{Trace}(A^2) = \sum_{i=1}^N \sum_{j=1}^N a_{ij} a_{ji} = \sum_{i=1}^N \sum_{j=1}^N a_{ij}^2 \sim N^2$$

$$\sum_{i=1}^N \lambda_i(A)^2 \sim N^2$$

Gives $N \text{Ave}(\lambda_i(A)^2) \sim N^2$ or $\text{Ave}(\lambda_i(A)) \sim \sqrt{N}$.

SKETCH OF PROOF: Averaging Formula

Recall k -th moment of $\mu_{A,N}(x)$ is $\text{Trace}(A^k)/2^k N^{k/2+1}$.

Average k -th moment is

$$\int \cdots \int \frac{\text{Trace}(A^k)}{2^k N^{k/2+1}} \prod_{i \leq j} p(a_{ij}) da_{ij}.$$

Proof by method of moments: Two steps

- Show average of k -th moments converge to moments of semi-circle as $N \rightarrow \infty$;
- Control variance (show it tends to zero as $N \rightarrow \infty$).

SKETCH OF PROOF: Averaging Formula for Second Moment

Substituting into expansion gives


$$\frac{1}{2^2 N^2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \sum_{i=1}^N \sum_{j=1}^N a_{ij}^2 \cdot p(a_{11}) da_{11} \cdots p(a_{NN}) da_{NN}$$

Integration factors as

$$\int_{a_{ij}=-\infty}^{\infty} a_{ij}^2 p(a_{ij}) da_{ij} \cdot \prod_{\substack{(k,l) \neq (i,j) \\ k < l}} \int_{a_{kl}=-\infty}^{\infty} p(a_{kl}) da_{kl} = 1.$$

Higher moments involve more advanced combinatorics (Catalan numbers).

Numerical example: Gaussian density




500 Matrices: Gaussian 400×400

$$p(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

Numerical example: Cauchy density $p(x) = 1/\pi(1 + x^2)$

Numerical example: Cauchy density $p(x) = 1/\pi(1+x^2)$



Spacings between events

GOE Conjecture

GOE Conjecture:

As $N \rightarrow \infty$, the probability density of the spacing b/w consecutive normalized eigenvalues approaches a limit independent of p .

GOE Conjecture

GOE Conjecture:


As $N \rightarrow \infty$, the probability density of the spacing b/w consecutive normalized eigenvalues approaches a limit independent of p .

Only known if p is a Gaussian.

$$\text{GOE}(x) \approx \frac{\pi}{2} x e^{-\pi x^2/4}.$$

Numerical Experiment: Uniform Distribution


Let $p(x) = \frac{1}{2}$ for $|x| \leq 1$.



5000: 300×300 uniform on $[-1, 1]$


Cauchy Distribution

$$\text{Let } p(x) = \frac{1}{\pi(1+x^2)}.$$




Cauchy Distribution

$$\text{Let } p(x) = \frac{1}{\pi(1+x^2)}.$$



Random Graphs



Degree of a vertex = number of edges leaving the vertex.
 Adjacency matrix: a_{ij} = number edges b/w Vertex i and Vertex j .


$$A = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 1 & 0 & 2 \\ 1 & 0 & 2 & 0 \end{pmatrix}$$

These are Real Symmetric Matrices.


McKay's Law (Kesten Measure) with $d = 3$

Density of Eigenvalues for d -regular graphs

$$f(x) = \begin{cases} \frac{d}{2\pi(d^2-x^2)} \sqrt{4(d-1)-x^2} & |x| \leq 2\sqrt{d-1} \\ 0 & \text{otherwise.} \end{cases}$$




McKay's Law (Kesten Measure) with $d = 6$



Fat Thin: fat enough to average, thin enough to get something different than Semi-circle.

3-Regular, 2000 Vertices and GOE

Spacings between eigenvalues of 3-regular graphs and the GOE:



Introduction to L -Functions

Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \operatorname{Re}(s) > 1.$$

Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1.$$

Unique Factorization: $n = p_1^{r_1} \cdots p_m^{r_m}$.

Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1.$$

Unique Factorization: $n = p_1^{r_1} \cdots p_m^{r_m}$.

$$\begin{aligned} \prod_p \left(1 - \frac{1}{p^s}\right)^{-1} &= \left[1 + \frac{1}{2^s} + \left(\frac{1}{2^s}\right)^2 + \cdots\right] \left[1 + \frac{1}{3^s} + \left(\frac{1}{3^s}\right)^2 + \cdots\right] \cdots \\ &= \sum_n \frac{1}{n^s}. \end{aligned}$$

Riemann Zeta Function (cont)

$$\zeta(s) = \sum_n \frac{1}{n^s} = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1$$

$$\pi(x) = \#\{p : p \text{ is prime}, p \leq x\}$$

Properties of $\zeta(s)$ and Primes:

Riemann Zeta Function (cont)

$$\zeta(s) = \sum_n \frac{1}{n^s} = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1$$

$$\pi(x) = \#\{p : p \text{ is prime}, p \leq x\}$$

Properties of $\zeta(s)$ and Primes:

- $\lim_{s \rightarrow 1^+} \zeta(s) = \infty, \pi(x) \rightarrow \infty.$

Riemann Zeta Function (cont)

$$\zeta(s) = \sum_n \frac{1}{n^s} = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1$$

$$\pi(x) = \#\{p : p \text{ is prime}, p \leq x\}$$

Properties of $\zeta(s)$ and Primes:

- $\lim_{s \rightarrow 1^+} \zeta(s) = \infty, \pi(x) \rightarrow \infty.$
- $\zeta(2) = \frac{\pi^2}{6}, \pi(x) \rightarrow \infty.$

Riemann Zeta Function

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \prod_{p \text{ prime}} \left(1 - \frac{1}{p^s}\right)^{-1}, \quad \text{Re}(s) > 1.$$

Functional Equation:


$$\xi(s) = \Gamma\left(\frac{s}{2}\right) \pi^{-\frac{s}{2}} \zeta(s) = \xi(1-s).$$

Riemann Hypothesis (RH):

All non-trivial zeros have $\text{Re}(s) = \frac{1}{2}$; can write zeros as $\frac{1}{2} + i\gamma$.

Observation: Spacings b/w zeros appear same as b/w eigenvalues of Complex Hermitian matrices $\overline{A}^T = A$.

Zeros of $\zeta(s)$ vs GUE



70 million spacings b/w adjacent zeros of $\zeta(s)$, starting at the 10^{20} th zero (from Odlyzko)

Bibliography



J. B. Conrey, *The Riemann hypothesis*, Notices of the AMS, **50** (2003), no. 3, 341–353.



B. Conrey, D. Farmer, P. Keating, M. Rubinstein and N. Snaith, *Integral Moments of L -Functions*, Proc. Lond. Math. Soc. **91** (2005), 33–104.



H. Davenport, *Multiplicative Number Theory, 3rd edition*, Graduate Texts in Mathematics **74**, Springer-Verlag, New York, 2000, revised by H. Montgomery.



N. Katz and P. Sarnak, *Zeros of zeta functions and symmetries*, Bull. AMS **36** (1999), 1–26.



J. P. Keating and N. C. Snaith, *Random matrices and L -functions*, Random matrix theory, J. Phys. A **36** (2003), 2859–2881.



M. Mehta, *Random Matrices*, 2nd edition, Academic Press, Boston, 1991.



S. J. Miller and R. Takloo-Bighash, *An Invitation to Modern Number Theory*, Princeton University Press, 2006.