

A Symplectic Test of the L -Functions Ratios Conjecture.

Steven J Miller
Brown University

`sjmiller@math.brown.edu`
`http://www.math.brown.edu/~sjmiller`

AMS Special Session, Courant, March 16th, 2008

History

- Farmer (1993): Considered

$$\int_0^T \frac{\zeta(\mathbf{s} + \alpha)\zeta(1 - \mathbf{s} + \beta)}{\zeta(\mathbf{s} + \gamma)\zeta(1 - \mathbf{s} + \delta)} dt,$$

conjectured (for appropriate values)

$$T \frac{(\alpha + \delta)(\beta + \gamma)}{(\alpha + \beta)(\gamma + \delta)} - T^{1-\alpha-\beta} \frac{(\delta - \beta)(\gamma - \alpha)}{(\alpha + \beta)(\gamma + \delta)}.$$

- Conrey-Farmer-Zirnbauer (2007): conjecture formulas for averages of products of L -functions over families:

$$R_{\mathcal{F}} = \sum_{f \in \mathcal{F}} \omega_f \frac{L\left(\frac{1}{2} + \alpha, f\right)}{L\left(\frac{1}{2} + \gamma, f\right)}.$$

Uses of the Ratios Conjecture

- **Applications:**
 - ◇ n -level correlations and densities;
 - ◇ mollifiers;
 - ◇ moments;
 - ◇ vanishing at the central point;
- **Advantages:**
 - ◇ RMT models often add arithmetic ad hoc;
 - ◇ predicts lower order terms, often to square-root level.

Inputs for 1-level density

- **Approximate Functional Equation:**

$$L(s, f) = \sum_{m \leq x} \frac{a_m}{m^s} + \epsilon \mathbb{X}_L(s) \sum_{n \leq y} \frac{a_n}{n^{1-s}};$$

- ◇ ϵ sign of the functional equation,
- ◇ $\mathbb{X}_L(s)$ ratio of Γ -factors from functional equation.

- **Explicit Formula:** g Schwartz test function,

$$\sum_{f \in \mathcal{F}} \omega_f \sum_{\gamma} g\left(\gamma \frac{\log N_f}{2\pi}\right) = \frac{1}{2\pi i} \int_{(c)} - \int_{(1-c)} R'_{\mathcal{F}}(\dots) g(\dots)$$

- ◇ $R'_{\mathcal{F}}(r) = \left. \frac{\partial}{\partial \alpha} R_{\mathcal{F}}(\alpha, \gamma) \right|_{\alpha=\gamma=r}$.

Procedure

- Use approximate functional equation to expand numerator.
- Expand denominator by generalized Mobius function: cusp form

$$\frac{1}{L(s, f)} = \sum_h \frac{\mu_f(h)}{h^s},$$

where $\mu_f(h)$ is the multiplicative function equaling 1 for $h = 1$, $-\lambda_f(p)$ if $h = p$, $\chi_0(p)$ if $h = p^2$ and 0 otherwise.

- Execute the sum over \mathcal{F} , keeping only main (diagonal) terms.
- Extend the m and n sums to infinity (complete the products).
- Differentiate with respect to the parameters.

Main Results

Symplectic Families

- Fundamental discriminants: d square-free and 1 modulo 4, or $d/4$ square-free and 2 or 3 modulo 4.
- Associated character χ_d :
 - ◇ $\chi_d(-1) = 1$ say d even;
 - ◇ $\chi_d(-1) = -1$ say d odd.
 - ◇ even (resp., odd) if $d > 0$ (resp., $d < 0$).

Will study following families:

- ◇ even fundamental discriminants at most X ;
- ◇ $\{8d : 0 < d \leq X, d \text{ an odd, positive square-free fundamental discriminant}\}$.

Prediction from Ratios Conjecture

$$\begin{aligned} \frac{1}{X^*} \sum_{d \leq X} \sum_{\gamma_d} g\left(\gamma_d \frac{\log X}{2\pi}\right) &= \frac{1}{X^* \log X} \int_{-\infty}^{\infty} g(\tau) \sum_{d \leq X} \left[\log \frac{d}{\pi} + \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{1}{4} \pm \frac{i\pi\tau}{\log X} \right) \right] d\tau \\ &+ \frac{2}{X^* \log X} \sum_{d \leq X} \int_{-\infty}^{\infty} g(\tau) \left[\frac{\zeta'}{\zeta} \left(1 + \frac{4\pi i\tau}{\log X} \right) + A'_D \left(\frac{2\pi i\tau}{\log X}; \frac{2\pi i\tau}{\log X} \right) \right. \\ &\left. - e^{-2\pi i\tau \log(d/\pi)/\log X} \frac{\Gamma\left(\frac{1}{4} - \frac{\pi i\tau}{\log X}\right)}{\Gamma\left(\frac{1}{4} + \frac{\pi i\tau}{\log X}\right)} \zeta \left(1 - \frac{4\pi i\tau}{\log X} \right) A_D \left(-\frac{2\pi i\tau}{\log X}; \frac{2\pi i\tau}{\log X} \right) \right] d\tau + O(X^{-\frac{1}{2}+\epsilon}), \end{aligned}$$

with

$$\begin{aligned} A_D(-r, r) &= \prod_p \left(1 - \frac{1}{(p+1)p^{1-2r}} - \frac{1}{p+1} \right) \cdot \left(1 - \frac{1}{p} \right)^{-1} \\ A'_D(r; r) &= \sum_p \frac{\log p}{(p+1)(p^{1+2r} - 1)}. \end{aligned}$$

Prediction from Ratios Conjecture

Main term is

$$\frac{1}{X^*} \sum_{d \leq X} \sum_{\gamma_d} g\left(\gamma_d \frac{\log X}{2\pi}\right) = \int_{-\infty}^{\infty} g(x) \left(1 - \frac{\sin(2\pi x)}{2\pi x}\right) dx$$

$$+ O\left(\frac{1}{\log X}\right),$$

which is the 1-level density for the scaling limit of $\mathrm{USp}(2N)$. If $\mathrm{supp}(\hat{g}) \subset (-1, 1)$, then the integral of $g(x)$ against $-\sin(2\pi x)/2\pi x$ is $-g(0)/2$.

Prediction from Ratios Conjecture

Assuming RH for $\zeta(s)$, for $\text{supp}(\widehat{g}) \subset (-\sigma, \sigma) \subset (-1, 1)$:

$$\begin{aligned} & \frac{-2}{X^* \log X} \sum_{d \leq X} \int_{-\infty}^{\infty} g(\tau) e^{-2\pi i \tau \frac{\log(d/\pi)}{\log X}} \frac{\Gamma\left(\frac{1}{4} - \frac{\pi i \tau}{\log X}\right)}{\Gamma\left(\frac{1}{4} + \frac{\pi i \tau}{\log X}\right)} \zeta\left(1 - \frac{4\pi i \tau}{\log X}\right) A_D\left(-\frac{2\pi i \tau}{\log X}; \frac{2\pi i \tau}{\log X}\right) d\tau \\ &= -\frac{g(0)}{2} + O\left(X^{-\frac{3}{4}(1-\sigma)+\epsilon}\right); \end{aligned}$$

the error term may be absorbed into the $O(X^{-1/2+\epsilon})$ error if $\sigma < 1/3$.


Main Results

Theorem (M– '07)

Let $\text{supp}(\widehat{g}) \subset (-\sigma, \sigma)$, assume RH for $\zeta(s)$. 1-Level Density agrees with prediction from Ratios Conjecture

- *up to $O(X^{-(1-\sigma)/2+\epsilon})$ for the family of quadratic Dirichlet characters with even fundamental discriminants at most X ;*
- *up to $O(X^{-1/2} + X^{-(1-\frac{3}{2}\sigma)+\epsilon} + X^{-\frac{3}{4}(1-\sigma)+\epsilon})$ for our sub-family. If $\sigma < 1/3$ then agrees up to $O(X^{-1/2+\epsilon})$.*

Numerics (J. Stopple): 1,003,083 negative fundamental discriminants $-d \in [10^{12}, 10^{12} + 3.3 \cdot 10^6]$



Histogram of normalized zeros ($\gamma \leq 1$, about 4 million).

- ◇ Red: main term.
- ◇ Blue: includes $O(1/\log X)$ terms.
- ◇ Green: all lower order terms.

Sketch of Proofs

Ratios Calculation

Hardest piece to analyze is

$$R(g; X) = -\frac{2}{X^* \log X} \sum_{d \leq X} \int_{-\infty}^{\infty} g(\tau) e^{-2\pi i \tau \frac{\log(d/\pi)}{\log X}} \frac{\Gamma\left(\frac{1}{4} - \frac{\pi i \tau}{\log X}\right)}{\Gamma\left(\frac{1}{4} + \frac{\pi i \tau}{\log X}\right)} \cdot \zeta\left(1 - \frac{4\pi i \tau}{\log X}\right) A_D\left(-\frac{2\pi i \tau}{\log X}; \frac{2\pi i \tau}{\log X}\right) d\tau,$$

$$A_D(-r, r) = \prod_p \left(1 - \frac{1}{(p+1)p^{1-2r}} - \frac{1}{p+1}\right) \cdot \left(1 - \frac{1}{p}\right)^{-1}.$$

Proof: shift contours, keep track of poles of ratios of Γ and zeta functions.

Ratios Calculation: Weaker result for $\text{supp}(\widehat{g}) \subset (-1, 1)$.

- d -sum is $X^* e^{-2\pi i(1 - \frac{\log \pi}{\log X})\tau} \left(1 - \frac{2\pi i\tau}{\log X}\right)^{-1} + O(X^{1/2})$;
- decay of g restricts τ -sum to $|\tau| \leq \log X$, Taylor expand everything but g : small error term and

$$\begin{aligned} & \int_{|\tau| \leq \log X} g(\tau) \sum_{n=-1}^N \frac{a_n}{\log^n X} (2\pi i\tau)^n e^{-2\pi i(1 - \frac{\log \pi}{\log X})\tau} d\tau \\ &= \sum_{n=-1}^N \frac{a_n}{\log^n X} \int_{|\tau| \leq \log X} (2\pi i\tau)^n g(\tau) e^{-2\pi i(1 - \frac{\log \pi}{\log X})\tau} d\tau; \end{aligned}$$

- from decay of g can extend the τ -integral to \mathbb{R} (essential that N is fixed and finite!), for $n \geq 0$ get the Fourier transform of $g^{(n)}$ (the n^{th} derivative of g) at $1 - \frac{\pi}{\log X}$, vanishes if $\text{supp}(\widehat{g}) \subset (-1, 1)$.

Number Theory Sums

$$\begin{aligned} S_{\text{even}} &= -\frac{2}{X^*} \sum_{d \leq X} \sum_{\ell=1}^{\infty} \sum_p \frac{\chi_d(p)^2 \log p}{p^\ell \log X} \widehat{g} \left(2 \frac{\log p^\ell}{\log X} \right) \\ S_{\text{odd}} &= -\frac{2}{X^*} \sum_{d \leq X} \sum_{\ell=0}^{\infty} \sum_p \frac{\chi_d(p) \log p}{p^{(2\ell+1)/2} \log X} \widehat{g} \left(\frac{\log p^{2\ell+1}}{\log X} \right). \end{aligned}$$

Number Theory Sums

Lemma

Let $\text{supp}(\hat{g}) \subset (-\sigma, \sigma) \subset (-1, 1)$. Then

$$\begin{aligned}
 S_{\text{even}} &= -\frac{g(0)}{2} + \frac{2}{\log X} \int_{-\infty}^{\infty} g(\tau) \frac{\zeta'}{\zeta} \left(1 + \frac{4\pi i\tau}{\log X}\right) d\tau \\
 &\quad + \frac{2}{\log X} \int_{-\infty}^{\infty} g(\tau) A'_D \left(\frac{2\pi i\tau}{\log X}; \frac{2\pi i\tau}{\log X}\right) + O(X^{-\frac{1}{2}+\epsilon}) \\
 S_{\text{odd}} &= O(X^{-\frac{1-\sigma}{2}} \log^6 X).
 \end{aligned}$$

If instead we consider the family of characters χ_{8d} for odd, positive square-free $d \in (0, X)$ (d a fundamental discriminant), then

$$S_{\text{odd}} = O(X^{-1/2+\epsilon} + X^{-(1-\frac{3}{2}\sigma)+\epsilon}).$$

Analysis of S_{even}

$\chi_d(p)^2 = 1$ except when $p|d$. Replace $\chi_d(p)^2$ with 1, and subtract off the contribution from when $p|d$:

$$\begin{aligned} S_{\text{even}} &= -2 \sum_{\ell=1}^{\infty} \sum_p \frac{\log p}{p^\ell \log X} \widehat{g} \left(2 \frac{\log p^\ell}{\log X} \right) \\ &\quad + \frac{2}{X^*} \sum_{d \leq X} \sum_{\ell=1}^{\infty} \sum_{p|d} \frac{\log p}{p^\ell \log X} \widehat{g} \left(2 \frac{\log p^\ell}{\log X} \right) \\ &= S_{\text{even};1} + S_{\text{even};2}. \end{aligned}$$

Lemma (Perron's Formula)

$$S_{\text{even};1} = -\frac{g(0)}{2} + \frac{2}{\log X} \int_{-\infty}^{\infty} g(\tau) \frac{\zeta'}{\zeta} \left(1 + \frac{4\pi i\tau}{\log X} \right) d\tau.$$

Analysis of S_{even} : $S_{\text{even};2}$

This piece gives us $\int g(\tau) A'_D(-\dots, \dots)$.

- Main ideas:
 - ◇ Restrict to $p \leq X^{1/2}$.
 - ◇ For $p < X^{1/2}$: $\sum_{d \leq X, p|d} 1 = \frac{X^*}{p+1} + O(X^{1/2})$.
 - ◇ Use Fourier Transform to expand \widehat{g} .

Analysis of S_{odd}

$$S_{\text{odd}} = -\frac{2}{X^*} \sum_{\ell=0}^{\infty} \sum_p \frac{\log p}{p^{(2\ell+1)/2} \log X} \widehat{g} \left(\frac{\log p^{2\ell+1}}{\log X} \right) \sum_{d \leq X} \chi_d(p).$$

Jutila's bound

$$\sum_{\substack{1 < n \leq N \\ n \text{ non-square}}} \left| \sum_{\substack{0 < d \leq X \\ d \text{ fund. disc.}}} \chi_d(n) \right|^2 \ll NX \log^{10} N.$$

Proof: Cauchy-Schwarz and Jutila: $p^{2\ell+1}$ non-square:

$$\left(\sum_{\ell=0}^{\infty} \sum_{p^{(2\ell+1)/2} \leq X^\sigma} \left| \sum_{d \leq X} \chi_d(p) \right|^2 \right)^{1/2} \ll X^{\frac{1+\sigma}{2}} \log^5 X.$$

Analysis of S_{odd} : Extending Support

More technical, replace Jutila's bound by applying Poisson Summation to character sums.

Lemma






Let $\text{supp}(\widehat{g}) \subset (-\sigma, \sigma) \subset (-1, 1)$. For family $\{8d : 0 < d \leq X, d \text{ an odd, positive square-free fundamental discriminant}\}$, $S_{\text{odd}} = O(X^{-\frac{1}{2}+\epsilon} + X^{-(1-\frac{3}{2}\sigma)+\epsilon})$.
In particular, if $\sigma < 1/3$ then $S_{\text{odd}} = O(X^{-1/2+\epsilon})$.






Conclusions






Conclusions






- Ratios Conjecture gives detailed predictions (up to $X^{1/2+\epsilon}$).
- Number Theory agrees with predictions for suitably restricted test functions.
- Numerics quite good.







References






-  M. V. Berry, *Semiclassical formula for the number variance of the Riemann zeros*, *Nonlinearity* **1** (1988), 399–407.
-  M. V. Berry and J. P. Keating, *The Riemann zeros and eigenvalue asymptotics*, *Siam Review* **41** (1999), no. 2, 236–266.
-  E. Bogomolny, O. Bohigas, P. Leboeuf and A. G. Monastera, *On the spacing distribution of the Riemann zeros: corrections to the asymptotic result*, *Journal of Physics A: Mathematical and General* **39** (2006), no. 34, 10743–10754.
-  E. B. Bogomolny and J. P. Keating, *Gutzwiller's trace formula and spectral statistics: beyond the diagonal approximation*, *Phys. Rev. Lett.* **77** (1996), no. 8, 1472–1475.
-  B. Conrey and D. Farmer, *Mean values of L-functions and symmetry*, *Internat. Math. Res. Notices* 2000, no. 17, 883–908.






-  B. Conrey, D. Farmer, P. Keating, M. Rubinstein and N. Snaith, *Integral moments of L-functions*, Proc. London Math. Soc. (3) **91** (2005), no. 1, 33–104.
-  J. B. Conrey, D. W. Farmer and M. R. Zirnbauer, *Autocorrelation of ratios of L-functions*, preprint.
<http://arxiv.org/abs/0711.0718>
-  J. B. Conrey, D. W. Farmer and M. R. Zirnbauer, *Howe pairs, supersymmetry, and ratios of random characteristic polynomials for the classical compact groups*, preprint.
<http://arxiv.org/abs/math-ph/0511024>
-  J. B. Conrey and N. C. Snaith, *Applications of the L-functions Ratios Conjecture*, Proc. Lon. Math. Soc. **93** (2007), no 3, 594–646.
-  J. B. Conrey and N. C. Snaith, *Triple correlation of the Riemann zeros*, preprint. <http://arxiv.org/abs/math/0610495>

-  H. Davenport, *Multiplicative Number Theory, 2nd edition*, Graduate Texts in Mathematics **74**, Springer-Verlag, New York, 1980, revised by H. Montgomery.
-  E. Dueñez, D. K. Huynh, J. P. Keating, S. J. Miller and N. C. Snaith, work in progress.
-  E. Dueñez and S. J. Miller, *The low lying zeros of a $GL(4)$ and a $GL(6)$ family of L -functions*, *Compositio Mathematica* **142** (2006), no. 6, 1403–1425.
-  E. Dueñez and S. J. Miller, *The effect of convolving families of L -functions on the underlying group symmetries*, preprint.
<http://arxiv.org/abs/math/0607688>
-  A. Erdélyi and F. G. Tricomi, *The asymptotic expansion of a ratio of gamma functions*, *Pacific J. Math.* **1** (1951), no. 1, 133–142.

-  E. Fouvry and H. Iwaniec, *Low-lying zeros of dihedral L-functions*, Duke Math. J. **116** (2003), no. 2, 189-217.
-  P. Gao, *N-level density of the low-lying zeros of quadratic Dirichlet L-functions*, Ph. D thesis, University of Michigan, 2005.
-  A. Güloğlu, *Low-Lying Zeros of Symmetric Power L-Functions*, Internat. Math. Res. Notices 2005, no. 9, 517-550.
-  G. Hardy and E. Wright, *An Introduction to the Theory of Numbers*, fifth edition, Oxford Science Publications, Clarendon Press, Oxford, 1995.
-  D. Hejhal, *On the triple correlation of zeros of the zeta function*, Internat. Math. Res. Notices 1994, no. 7, 294-302.

-  C. Hughes and S. J. Miller, *Low-lying zeros of L-functions with orthogonal symmetry*, Duke Math. J., **136** (2007), no. 1, 115–172.
-  C. Hughes and Z. Rudnick, *Linear Statistics of Low-Lying Zeros of L-functions*, Quart. J. Math. Oxford **54** (2003), 309–333.
-  D. K. Huynh, J. P. Keating and N. C. Snaith, work in progress.
-  H. Iwaniec, W. Luo and P. Sarnak, *Low lying zeros of families of L-functions*, Inst. Hautes Études Sci. Publ. Math. **91**, 2000, 55–131.
-  M. Jutila, *On character sums and class numbers*, Journal of Number Theory **5** (1973), 203–214.
-  M. Jutila, *On mean values of Dirichlet polynomials with real characters*, Acta Arith. **27** (1975), 191–198.

-  M. Jutila, *On the mean value of $L(1/2, \chi)$ for real characters*, *Analysis* **1** (1981), no. 2, 149–161.
-  N. Katz and P. Sarnak, *Random Matrices, Frobenius Eigenvalues and Monodromy*, AMS Colloquium Publications **45**, AMS, Providence, 1999.
-  N. Katz and P. Sarnak, *Zeros of zeta functions and symmetries*, *Bull. AMS* **36**, 1999, 1 – 26.
-  J. P. Keating, *Statistics of quantum eigenvalues and the Riemann zeros*, in *Supersymmetry and Trace Formulae: Chaos and Disorder*, eds. I. V. Lerner, J. P. Keating & D. E. Khmelnitskii (Plenum Press), 1–15.
-  J. P. Keating and N. C. Snaith, *Random matrix theory and $\zeta(1/2 + it)$* , *Comm. Math. Phys.* **214** (2000), no. 1, 57–89.

-  J. P. Keating and N. C. Snaith, *Random matrix theory and L-functions at $s = 1/2$* , Comm. Math. Phys. **214** (2000), no. 1, 91–110.
-  J. P. Keating and N. C. Snaith, *Random matrices and L-functions*, Random matrix theory, J. Phys. A **36** (2003), no. 12, 2859–2881.
-  S. J. Miller, *1- and 2-level densities for families of elliptic curves: evidence for the underlying group symmetries*, Compositio Mathematica **104** (2004), 952–992.
-  S. J. Miller, *Variation in the number of points on elliptic curves and applications to excess rank*, C. R. Math. Rep. Acad. Sci. Canada **27** (2005), no. 4, 111–120.
-  S. J. Miller, *Lower order terms in the 1-level density for families of holomorphic cuspidal newforms*, preprint.
<http://arxiv.org/abs/0704.0924>



H. Montgomery, *The pair correlation of zeros of the zeta function*, Analytic Number Theory, Proc. Sympos. Pure Math. **24**, Amer. Math. Soc., Providence, 1973, 181 – 193.



A. Odlyzko, *On the distribution of spacings between zeros of the zeta function*, Math. Comp. **48** (1987), no. 177, 273–308.









A. Odlyzko, *The 10^{22} -nd zero of the Riemann zeta function*, Proc. Conference on Dynamical, Spectral and Arithmetic Zeta-Functions, M. van Frankenhuysen and M. L. Lapidus, eds., Amer. Math. Soc., Contemporary Math. series, 2001,
<http://www.research.att.com/~amo/doc/zeta.html>.



A. E. Özlük and C. Snyder, *Small zeros of quadratic L-functions*, Bull. Austral. Math. Soc. **47** (1993), no. 2, 307–319.



A. E. Özlük and C. Snyder, *On the distribution of the nontrivial zeros of quadratic L-functions close to the real axis*, Acta Arith. **91** (1999), no. 3, 209–228.

-  G. Ricotta and E. Royer, *Statistics for low-lying zeros of symmetric power L-functions in the level aspect*, preprint.
<http://arxiv.org/abs/math/0703760>
-  E. Royer, *Petits zéros de fonctions L de formes modulaires*, Acta Arith. **99** (2001), no. 2, 147-172.
-  M. Rubinstein, *Low-lying zeros of L-functions and random matrix theory*, Duke Math. J. **109**, (2001), 147–181.
-  M. Rubinstein, *Computational methods and experiments in analytic number theory*. Pages 407–483 in Recent Perspectives in Random Matrix Theory and Number Theory, ed. F. Mezzadri and N. C. Snaith editors, 2005.
-  Z. Rudnick and P. Sarnak, *Zeros of principal L-functions and random matrix theory*, Duke Math. J. **81**, 1996, 269 – 322.
-  K. Soundararajan, *Nonvanishing of quadratic Dirichlet L-functions at $s = 1/2$* , Ann. of Math. (2) **152** (2000), 447–488.



J. Stopple, *The quadratic character experiment*, preprint.

<http://arxiv.org/abs/0802.4255>



M. Young, *Lower-order terms of the 1-level density of families of elliptic curves*, Internat. Math. Res. Notices 2005, no. 10, 587–633.



M. Young, *Low-lying zeros of families of elliptic curves*, J. Amer. Math. Soc. **19** (2006), no. 1, 205–250.