

ON NEAR PERFECT NUMBERS

PETER COHEN, KATHERINE CORDWELL, ALYSSA EPSTEIN, CHUNG HANG KWAN, ADAM LOTT,
AND STEVEN J. MILLER

ABSTRACT. The study of perfect numbers (numbers which equal the sum of their proper divisors) goes back to antiquity, and is responsible for some of the oldest and most popular conjectures in number theory. We investigate a generalization introduced by Pollack and Shevelev: k -near-perfect numbers. These are examples to the well-known pseudoperfect numbers first defined by Sierpiński, and are numbers such that the sum of all but at most k of its proper divisors equals the number. We establish their asymptotic order for all integers $k \geq 4$, as well as some properties of related quantities.

1. INTRODUCTION

Let $\sigma(n)$ be the sum of all positive divisors of n . A natural number n is *perfect* if $\sigma(n) = 2n$. Perfect numbers have played a prominent role in classical number theory for millennia. A well-known conjecture claims that there are infinitely many even, but no odd, perfect numbers. Despite the fact that these conjectures remain unproven, there has been significant progress on studying the distribution of perfect numbers [Vo, HoWi, Ka, Er1], as well as generalizations. One are the *pseudoperfect numbers*, which were introduced by Sierpiński [Si]. A natural number is pseudoperfect if it is a sum of some subset of its proper divisors. Erdős and Benkoski [Er2, BeEr] proved that the asymptotic density for pseudoperfect numbers, as well as that of abundant numbers that are not pseudoperfect (also called *weird numbers*), exist and are positive.

Pollack and Shevelev [PoSh] initiated the study of a subclass of pseudoperfect numbers called *near-perfect numbers*. A natural number is k -near-perfect if it is a sum of all of its proper divisors with *at most* k exceptions. Restriction on the number of exceptional divisors leads to asymptotic density 0. The number of 1-near-perfect numbers up to x is¹ at most $x^{3/4+o(1)}$, and in general for $k \geq 1$ the number of k -near-perfect numbers up to x is at most $x(\log \log x)^{k-1}/\log x$.

Our first result improves the count of k -near-perfect numbers.

Theorem 1.1. *For any non-negative integer k and real number $x \geq 1$, denote by $N(k; x)$ the set of k -near-perfect numbers up to x .*

For any $k \geq 4$, there exists a constant $x_0(k) > 0$ such that for $x \geq x_0(k)$, we have

$$\#N(k; x) \asymp_k \frac{x}{\log x} (\log \log x)^{\lfloor \frac{\log(k+4)}{\log 2} \rfloor - 3}. \quad (1.1)$$

Date: September 26, 2019.

2010 Mathematics Subject Classification. 11A25 (primary), 11N25, 11B83 (secondary).

Key words and phrases. Perfect numbers, Near-perfect numbers, Pseudoperfect numbers, Arithmetic functions, Sum of divisors function.

¹This is a result stated in [AnPoPo]. In the original paper of Pollack and Shevelev [PoSh], the upper bound was given by $x^{5/6+o(1)}$.

Our argument is based on a partition of the set $N(k; x)$ different from that of [PoSh] and this is described in Section 3. This allows us to carry out an inductive argument and reduces the count of $\#N(k; x)$ for large integers k to the determination of all k -near-perfect numbers for small integers k with a fixed number of positive divisors (see Lemma 2.10). When $4 \leq k \leq 11$, this even allows precise asymptotic formulae.

Theorem 1.2. *For $4 \leq k \leq 11$, there exists a constant $c_k > 0$ such that*

$$\#N(k; x) \sim c_k \frac{x}{\log x} \quad (1.2)$$

as $x \rightarrow \infty$.

Indeed, the computation of the constant c_k follows from Lemma 2.10 and

$$\begin{aligned} c_4 &= c_5 \approx 0.1667, \quad c_6 \approx 0.2024, \\ c_7 &= c_8 \approx 0.3913, \quad c_9 \approx 0.4968, \quad c_{10} \approx 0.5709, \quad c_{11} \approx 0.6274. \end{aligned}$$

Our last result is motivated by an open question raised in [BeEr]: can $\sigma(n)/n$ be arbitrarily large when n is a weird number? ² We replace ‘weirdness’ by ‘exact-perfectness’, where a natural number is k -exact-perfect if it is a sum of all of its proper divisors with exactly k exceptions. Note the result below is conditional on there being no odd perfect numbers.

Theorem 1.3. *Let $\epsilon \in (0, 2/5)$. Denote by $E(k)$ the set of all k -exact-perfect numbers, $E(k; x) := E(k) \cap [1, x]$ and $E_\epsilon(k; x) := \{n \leq x : n \in E(k), \sigma(n) \geq 2n + n^\epsilon\}$. Let M be the set of all natural numbers of the form $2q$, where q is a Mersenne prime³. If there are no odd perfect number, then for k sufficiently large and $k \notin M$, we have*

$$\lim_{x \rightarrow \infty} \frac{\#E_\epsilon(k; x)}{\#E(k; x)} = 1. \quad (1.3)$$

1.1. Outline. In Section 2 we introduce the necessary definitions and lemmata for our theorems. In Section 3, we set the stage for proving Theorem 1.1 and 1.2. In Section 4, 5 and 6, we prove Theorem 1.2, 1.1 and 1.3 and respectively.

1.2. Notations. We use the following notations and definitions.

- We write $f(x) \asymp g(x)$ if there exist positive constants c_1, c_2 such that $c_1 g(x) < f(x) < c_2 g(x)$ for all sufficiently large x .
- We write $f(x) \sim g(x)$ if $\lim_{x \rightarrow \infty} f(x)/g(x) = 1$.
- We write $f(x) = O(g(x))$ or $f(x) \ll g(x)$ if there exists a positive constant C such that $f(x) < Cg(x)$ for all sufficiently large x .
- We write $f(x) = o(g(x))$ if $\lim_{x \rightarrow \infty} f(x)/g(x) = 0$.
- In all cases, subscripts indicate dependence of implied constants on other parameters.
- Let $x \geq y \geq 2$. Denote by $\Phi(x, y)$ the set of all y -smooth numbers up to x and $\Phi_j(x, y) := \{n \leq x : n = p_1 \cdots p_j m_j, P^+(m_j) \leq y < p_j < \cdots < p_1\}$.
- We use p and p_i to denote primes, and $P^+(n)$ to denote the largest prime factor of n .
- Denote by $\tau(n)$ the number of positive divisors of n .
- Denote by $\Omega(n)$ the number of prime divisors of n counting multiplicities.

²A number is *weird* if the sum of its proper divisors is greater than itself, but no subset of these divisors sums to the original number.

³Mersenne primes are primes of the form $2^p - 1$ for some prime p .

- Denote by $N(k)$ the set of all k -near-perfect numbers and $N(k; x) := N(k) \cap [1, x]$.
- Denote by $E(k)$ the set of all k -exact-perfect numbers and $E(k; x) := E(k) \cap [1, x]$.

1.3. Acknowledgments. This work was supported in part by NSF Grants DMS1265673, DMS1347804, DMS1561945, and DMS1659037, the Williams SMALL REU Program, the Clare Boothe Luce Program, the COSINE Program of the Chinese University of Hong Kong and the Professor Charles K. Kao Research Exchange Scholarship 2015/16. We thank Kevin Ford, Charles Chun Che Li, Paul Pollack and Carl Pomerance for helpful discussions. We also thank the referee for his/her careful reading and valuable comments.

2. PREPARATIONS

In this section, we collect the necessary lemmata for our theorems. We begin with a well-known result of Landau regarding the arithmetic function $\Omega(n)$, the number of prime factors of n counting multiplicities, i.e., if $n = p_1^{a_1} \cdots p_r^{a_r}$, then $\Omega(n) = a_1 + \cdots + a_r$. We let

$$\Omega(s; x) := \{n \leq x : \Omega(n) = s\}. \quad (2.1)$$

Lemma 2.1. *Fix an integer $s \geq 1$. As $x \rightarrow \infty$, we have*

$$\#\Omega(s; x) \sim \#\{n \leq x : n = p_1 \cdots p_s, p_1 > \cdots > p_s\} \sim \frac{1}{(s-1)!} \frac{x}{\log x} (\log \log x)^{s-1}. \quad (2.2)$$

Proof. See [HaWr] Theorem 437 (Section 22.18). \square

Next, we state an elementary estimate of the number of y -smooth numbers up to x .⁴

Lemma 2.2. *Let*

$$u := \frac{\log x}{\log y} \quad (2.3)$$

and $\Phi(x, y)$ be the set of y -smooth numbers up to x . Then uniformly for $x \geq y \geq 2$, we have

$$\#\Phi(x, y) \ll x \exp(-u/2). \quad (2.4)$$

Proof. See Theorem 9.5 of [DeKLu]. \square

Our next lemma is a standard result from sieve theory.

Lemma 2.3. *Suppose A is a finite set of natural numbers, P is a set of primes, $z > 0$ and $P(z)$ is the product of primes in P not greater than z . Let*

$$S(A, P, z) := \{n \in A : (n, P(z)) = 1\}$$

and

$$A_d := \{a \in A : d \mid a\}.$$

Assume the following conditions.

⁴Let $y \geq 2$. A natural number n is said to be y -smooth if all of its prime factors are at most y .

(1) Suppose g is a multiplicative function satisfying

$$0 \leq g(p) < 1 \text{ for } p \in P \text{ and } g(p) = 0 \text{ for } p \notin P,$$

and there exists constants $B > 0$ and $\kappa \geq 0$ such that

$$\prod_{y \leq p \leq w} (1 - g(p))^{-1} \leq \left(\frac{\log w}{\log y} \right)^\kappa \exp \left(\frac{B}{\log y} \right)$$

for $2 \leq y < w$.

(2) Let $X > 0$. For any square-free number d with all of its prime factors in P , define

$$r_d := \#A_d - Xg(d).$$

Assume that r_d satisfies the inequality

$$\sum_{\substack{d|P(z) \\ d \leq X^\theta}} |r_d| \leq C \frac{x}{(\log x)^\kappa}.$$

for some $\theta > 0$.

Then for $2 \leq z \leq X$, we have

$$\#S(A, P, z) \ll_{\kappa, \theta, C, B} XV(z), \quad (2.5)$$

where

$$V(z) := \prod_{\substack{p \leq z \\ p \in P}} (1 - g(p)). \quad (2.6)$$

Proof. For example, see [FoHa]. \square

In the proof of Theorem 1.1, an estimate is needed for the size of the set

$$\Phi_j(x, y) := \{n \leq x : n = p_1 \cdots p_j m_j, P^+(m_j) \leq y < p_j < \cdots < p_1\}, \quad (2.7)$$

where $j \geq 1$, $x \geq y \geq 2$ and $P^+(m_j)$ denotes the largest prime factor of m_j . It follows from Lemmas 2.2 and 2.3.

Lemma 2.4. Suppose $j \geq 1$, $x \geq y \geq e$, and $y \leq x^{\frac{1}{4(j+1) \log \log x}}$. Then

$$\#\Phi_j(x, y) \ll_j \frac{x \log y}{\log x} (\log \log x)^{j-1}. \quad (2.8)$$

Proof. Introduce the following sets:

$$Q^{(0)}(x, y) := \{n \leq x : n = p_1 \cdots p_j m_j, P^+(m_j) \leq y < p_j < \cdots < p_1 \leq x^{\frac{1}{j+1}}\} \quad (2.9)$$

$$Q^{(j)}(x, y) := \{n \leq x : n = p_1 \cdots p_j m_j, P^+(m_j) \leq y < x^{\frac{1}{j+1}} < p_j < \cdots < p_1\}, \quad (2.10)$$

and for $1 \leq i \leq j-1$,

$$Q^{(i)}(x, y) := \{n \leq x : n = p_1 \cdots p_j m_j, P^+(m_j) \leq y < p_j < \cdots < p_{i+1} \leq x^{\frac{1}{j+1}} < p_i < \cdots < p_1\}. \quad (2.11)$$

Clearly, we have

$$\#\Phi_j(x, y) = \sum_{0 \leq i \leq j} Q^{(i)}(x, y). \quad (2.12)$$

(1) By (2.4) of Lemma 2.2, we estimate $\#Q^{(0)}(x, y)$ as follows:

$$\begin{aligned}
\#Q^{(0)}(x, y) &= \sum_{y < p_j < \dots < p_1 \leq x^{\frac{1}{j+1}}} \sum_{\substack{P^+(m_j) \leq y \\ m_j \leq x/(p_1 \dots p_j)}} 1 \\
&\ll \sum_{y < p_j < \dots < p_1 \leq x^{\frac{1}{j+1}}} \frac{x}{p_1 \dots p_j} \exp\left(-\frac{\log(x/p_1 \dots p_j)}{2 \log y}\right) \\
&\leq \sum_{y < p_j < \dots < p_1 \leq x^{\frac{1}{j+1}}} \frac{x}{p_1 \dots p_j} \exp\left(-\frac{1}{2(j+1)} \frac{\log x}{\log y}\right) \\
&\leq x \exp\left(-\frac{1}{2(j+1)} \frac{\log x}{\log y}\right) \left(\sum_{p \leq x^{\frac{1}{j+1}}} \frac{1}{p}\right)^j \\
&\ll_j x (\log \log x)^j \exp\left(-\frac{1}{2(j+1)} \frac{\log x}{\log y}\right) \tag{2.13}
\end{aligned}$$

We have to make sure that (2.13) is of acceptable size. Indeed, since

$$y \leq x^{\frac{1}{4(j+1) \log \log x}},$$

it follows that

$$\begin{aligned}
\frac{\log x}{\log y} \log \log x &= \exp(\log \log x + \log \log \log x - \log \log y) \\
&\leq \exp\left(\frac{4(j+1)(\log y)(\log \log x)}{2(j+1)(\log y)}\right) \\
&\leq \exp\left(\frac{\log x}{2(j+1)(\log y)}\right)
\end{aligned}$$

and

$$x (\log \log x)^j \exp\left(-\frac{1}{2(j+1)} \frac{\log x}{\log y}\right) \leq \frac{x \log y}{\log x} (\log \log x)^{j-1}.$$

(2) In order to estimate $\#Q^{(j)}(x, y)$, we apply Lemma 2.3 with A being the set of all natural numbers up to x , P being the set of primes in $(y, x^{1/(j+1)}]$, $z := x^{1/(j+1)}$, $X := x$ and $g(d) := 1/d$. Then $S(A, P, z)$ is the set of all natural numbers up to x whose prime factors are at most y or at least $x^{\frac{1}{j+1}}$. (Note that there are at most j prime factors can be larger than $x^{\frac{1}{j+1}}$.)

By Mertens' estimates, we can see that all of the assumptions of Lemma 2.3 are satisfied with $\kappa = B = C = 1$ and $\theta = 1/2$. Hence, we have

$$\#Q^{(j)}(x) \leq \#S(A, P, z) \ll \frac{x \log y}{\log x}. \tag{2.14}$$

(3) For $1 \leq i \leq j-1$, we estimate $\#Q^{(i)}(x, y)$ also by using Lemma 2.3. For any choices of primes p_{i+1}, \dots, p_j such that $y < p_j < \dots < p_{i+1} \leq x^{\frac{1}{j+1}}$, we choose

$$X := \frac{x}{p_{i+1} \dots p_j},$$

A being the set of all natural numbers up to X , P being the set of primes in $(y, x^{1/(j+1)}]$, $z := x^{1/(j+1)}$ and $g(d) := 1/d$. Hence,

$$\begin{aligned}
\#Q^{(i)}(x, y) &= \sum_{y < p_j < \dots < p_{i+1} \leq x^{\frac{1}{j+1}}} \sum_{\substack{P^+(m_j) \leq y \\ p_1 > \dots > p_i > x^{\frac{1}{j+1}} \\ p_1 \dots p_i m_j \leq x / (p_{i+1} \dots p_j)}} 1 \\
&\ll \sum_{y < p_j < \dots < p_{i+1} \leq x^{\frac{1}{j+1}}} \frac{x}{p_{i+1} \dots p_j} \frac{\log y}{\log x} \\
&\leq \frac{x \log y}{\log x} \left(\sum_{p \leq x^{\frac{1}{j+1}}} \frac{1}{p} \right)^{j-i} \ll_j \frac{x \log y}{\log x} (\log \log x)^{j-i}. \tag{2.15}
\end{aligned}$$

The result now follows from (2.12), (2.13), (2.14) and (2.15). \square

Remark 2.5. Since

$$\{n \leq x : n = p_1 \dots p_j m_j, m_j \leq y < p_j < \dots < p_1\} \subset \Phi_j(x, y), \tag{2.16}$$

it follows from Lemma 2.1 that

$$\begin{aligned}
\#\Phi_j(x, y) &\geq \sum_{m_j \leq y} \sum_{\substack{n_j \leq \frac{x}{m_j} \\ n_j = p_1 \dots p_j \\ \text{for some } p_1 > \dots > p_j > y}} 1 \\
&\gg \sum_{m_j \leq y} \frac{x/m_j}{\log(x/m_j)} \left(\log \log \frac{x}{m_j} \right)^{j-1} \\
&\geq \frac{x}{\log x} \left(\log \log \frac{x}{y} \right)^{j-1} \sum_{m_j \leq y} \frac{1}{m_j} \\
&\gg \frac{x \log y}{\log x} \left(\log \log \frac{x}{y} \right)^{j-1}. \tag{2.17}
\end{aligned}$$

Below we state some elementary observations about near-perfect numbers.

Lemma 2.6. Prime powers cannot be k -near-perfect for any integer $k \geq 0$.

Proof. This follows directly from the definition of near-perfect numbers and the uniqueness of q -ary representation. \square

Lemma 2.7 (Euclid-Euler). All even perfect numbers are of the form $2^{p-1}(2^p - 1)$, where p is a Mersenne prime, i.e., a prime p such that $2^p - 1$ is also a prime.

Lemma 2.8. An odd perfect number has at least 4 distinct prime factors.

In fact, it is now known that an odd perfect number must have at least 10 distinct prime factors. This is due to Nielsen [Niel]. The proof of an odd perfect has at least 4 distinct prime factors is completely elementary.

The following lemma resembles the aforementioned theorem of Euclid-Euler and it serves as a complete classification of 1-near-perfect numbers with two distinct prime factors. This is helpful in reducing the number of cases to be considered in Lemma 2.10.

Lemma 2.9. *A 1-near-perfect number which is not perfect and has two distinct prime factors is of the form*

- (1) $2^{t-1}(2^t - 2^k - 1)$, where $2^t - 2^k - 1$ is prime,
- (2) $2^{2p-1}(2^p - 1)$, where p is a Mersenne prime.
- (3) $2^{p-1}(2^p - 1)^2$, where p is a Mersenne prime.
- (4) 40.

Proof. See [ReCh]. □

Upon carrying out the recursive process as described in Section 3 and 5, it boils down to prove the following lemma which can be done by explicit computation.

Lemma 2.10. *Let $\tau(m)$ be the number of positive divisors of the positive integer m .*

- (1) *If $\tau(m)$ is prime, then m cannot be k -near-perfect for any integer $k \geq 0$.*
- (2) *Suppose $\tau(m) = 10$. Then*
 - (a) *if m is perfect, then $m = 496$.*
 - (b) *if m is 1-near-perfect, then $m \in \{496, 368, 464\}$.*
- (3) *Suppose $\tau(m) = 9$. Then*
 - (a) *m cannot be perfect.*
 - (b) *if m is 1-near-perfect, then $m = 196$.*
 - (c) *if m is 2-near-perfect, then $m \in \{196, 36\}$.*
- (4) *Suppose $\tau(m) = 8$. Then*
 - (a) *m cannot be perfect.*
 - (b) *if m is 1-near-perfect, then $m \in \{24, 40, 56, 88, 104\}$.*
 - (c) *if m is 2-near-perfect, then $m \in \{24, 40, 56, 88, 104, 30, 54, 66\}$.*
 - (d) *if m is 3-near-perfect, then $m \in \{24, 40, 56, 88, 104, 30, 54, 66, 42\}$.*
- (5) *Suppose $\tau(m) = 6$. Then if m is k -near-perfect for some $k \geq 0$, then $m \in \{28, 12, 18, 20\}$.*
- (6) *Suppose $\tau(m) = 4$. Then if m is k -near-perfect for some $k \geq 0$, then $m = 6$.*

Proof.

- (1) Follows immediately from Lemma 2.6.
- (2) Suppose m is a 1-near-perfect and $\tau(m) = 10$. Since $\tau(m) = 10$, m is of the form q^9 or q^4r , where q, r are distinct primes. The first case cannot happen by Lemma 2.6.

Now suppose the second case. If m is perfect, by Lemma 2.8, it must be even. Then by Lemma 2.7, $m = q^4r = 2^{p-1}(2^p - 1)$ for some Mersenne prime p . It follows that

$$q = 2, p - 1 = 4 \text{ and } r = 2^p - 1,$$

i.e., $(q, r) = (2, 31)$ and $p = 5$. Note that q, r are distinct primes and p is a Mersenne prime. Thus, we have $m = 2^4 \cdot 31 = 496$.

If m is 1-near-perfect but not perfect, we use Lemma 2.9 instead and similarly, we have $m = 2^4 \cdot 23 = 368$, $2^4 \cdot 29 = 464$.

Thus, all the possible m 's are 368, 464, 496.

- (3) Suppose m is a 2-near-perfect and $\tau(m) = 9$. Since $\tau(m) = 9$, m is of the form q^8 or q^2r^2 , where q, r are distinct primes. The first case cannot happen by Lemma 2.6. If $m = q^2r^2$ is 1-near-perfect, as before by Lemma 2.7, 2.8 and 2.9, the only possibility is $m = 2^2 \cdot 7^2 = 196$.

Now suppose $m = q^2 r^2$ is 2-near-perfect but not 1-near-perfect. It suffices to consider the following 16 equations by observing the symmetry of q and r in $q^2 r^2$:

$$\begin{aligned} & (1 + q + q^2)(1 + r + r^2) - 2q^2 r^2 \\ = & 1 + q, 1 + q^2, 1 + qr, 1 + q^2 r, q + q^2, q + r, q^2 + r, q + qr, q + q^2 r, \\ & r + q^2 r, q^2 + r^2, q^2 + qr, q^2 + q^2 r, q^2 + qr^2, qr + q^2 r, q^2 r + qr^2. \end{aligned} \quad (2.18)$$

Given any $q, r \geq 2$, it is clear that $1 + q$ is the smallest among the 16 expressions on the right side of (2.18). We claim that if $q \geq 7$ and $r \geq 2$, then

$$(1 + q + q^2)(1 + r + r^2) - 2q^2 r^2 < 1 + q,$$

i.e.,

$$f_q(r) := (q^2 - q - 1)r^2 - (1 + q + q^2)r - q^2 > 0.$$

This is simply a quadratic polynomial inequality in r . Note that $q^2 - q - 1 > 0$ and

$$\begin{aligned} \Delta(q) &:= (1 + q + q^2)^2 + 4q^2(q^2 - q - 1) = 5q^4 - 2q^3 - q^2 + 2q + 1 \\ &> 0. \end{aligned}$$

Thus, if

$$r > \frac{(1 + q + q^2) + \sqrt{\Delta(q)}}{2(q^2 - q - 1)} \quad (2.19)$$

then

$$f_q(r) > 0.$$

The inequality (2.19) is satisfied with $q \geq 7$ and $r \geq 2$ since for $q \geq 7$, we have

$$2 > \frac{(1 + q + q^2) + \sqrt{\Delta(q)}}{2(q^2 - q - 1)}.$$

The claim follows.

Thus by this claim, the left side of (2.18) is strictly less than each of the 16 possibilities of the right side of (2.18) when $q \geq 7$ and $r \geq 2$. Now, it suffices to solve the 16 equations in r with $q = 2, 3, 5$. The only solution is $(q, r) = (3, 2)$ (i.e., $m = 36$), which comes from the equation

$$(1 + q + q^2)(1 + r + r^2) = 2q^2 r^2 + 1 + q^2 r.$$

- (4) Suppose m is 3-near-perfect and $\tau(m) = 8$. Since $\tau(m) = 8$, m is of the form q^7 , $q^3 r$ or qrs , where q, r, s are distinct primes. Once again by Lemma 2.6, the first case is impossible.

- (a) Suppose $m = q^3 r$. As we have done in (2) and (3), if m is 1-near-perfect, then

$$\begin{aligned} m &= 2^3 \cdot 7 = 56, 2^3 \cdot 11 = 88, 2^3 \cdot 13 = 104, \\ &2^3 \cdot 3 = 24, 2^3 \cdot 5 = 40, \end{aligned}$$

by Lemmas 2.7, 2.8 and 2.9.

Suppose $m = q^3 r$ is 3-near-perfect but not 1-near-perfect. Then it suffices to consider the $\binom{7}{2} + \binom{7}{2} = 56$ equations formed by all of the possible pairs or

triples distinct proper divisors. Following the steps in (3), out of the sums of these pairs or triples, the smallest ones are $1 + q$ or $1 + r$.

When $q \geq 5$ and $r \geq 2$, we have

$$r \geq 2 > \frac{q^3 + q^2}{q^3 - q^2 - q - 1} \quad \text{and} \quad r \geq 2 > \frac{q^2 + q + 1}{q^2 - q - 1}.$$

These imply that

$$(1 + q + q^2 + q^3)(1 + r) - 2q^3r < 1 + q, 1 + r. \quad (2.20)$$

The same inequality is valid by replacing the right side of (2.20) by the sum of any of the 56 possible pairs or triples of proper divisors of $m = q^3r$, when $q \geq 5$ and $r \geq 2$.

It remains to solve the 56 equations in r with $q = 2, 3$. Only the following equations are solvable:

- (i) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = 1 + r$; $(q, r) = (2, 7)$ (i.e., $m = 56$),
- (ii) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = q + q^2$; $(q, r) = (3, 2)$ (i.e., $m = 54$),
- (iii) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = q + q^3$; $(q, r) = (2, 5)$ (i.e., $m = 40$),
- (iv) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = q^2 + q^3$; $(q, r) = (2, 3)$ (i.e., $m = 24$),
- (v) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = 1 + q^2 + r$; $(q, r) = (2, 5)$ (i.e., $m = 40$), $(q, r) = (3, 2)$ (i.e., $m = 54$),
- (vi) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = 1 + q^3 + r$; $(q, r) = (2, 3)$ (i.e., $m = 24$),
- (vii) $(1 + q + q^2 + q^3)(1 + r) - 2q^3r = q + q^2 + rq$; $(q, r) = (2, 3)$ (i.e., $m = 24$).

- (b) Suppose $m = qrs$. It cannot be perfect. We shall use a similar strategy as above. By symmetry, it suffices to solve the following 19 equations one-by-one:

$$\begin{aligned} & (1 + q)(1 + r)(1 + s) - 2qrs \\ &= 1, q, qr, 1 + q, 1 + qr, q + r, q + qr, q + rs, qr + qs, \\ & 1 + q + r, 1 + q + qr, 1 + q + rs, 1 + qr + qs, q + r + s, \\ & q + r + qr, q + qr + s, q + qr + rs, q + qr + qs, qr + rs + qs. \end{aligned} \quad (2.21)$$

We claim that

$$(1 + q)(1 + r)(1 + s) - 2qrs < 1$$

for $q \geq 11, r \geq 5$ and $s \geq 2$.

This can be verified as follows. Since $s \geq 2$, we have

$$\frac{12s + 11}{10s - 2} < 5.$$

Then by $r \geq 5$, we have

$$r > \frac{12s + 11}{10s - 2}.$$

This implies that

$$11(rs - r - s - 1) > rs + r + s.$$

By $rs - r - s - 1 > 0$ and $q \geq 11$, we have

$$q(rs - r - s - 1) > rs + r + s,$$

and the claim follows.

Thus, it suffices to solve the equations with $q = 2, 3, 5, 7$ or $r = 2, 3$. This reduces the 19 three-variable equations in (2.21) to two-variable ones.

Out of these equations, only the following equations are solvable:

- (i) $(1+q)(1+r)(1+s) - 2qrs = 1+q$; $(q, r, s) = (11, 2, 3), (11, 3, 2)$ ($m = 66$),
 - (ii) $(1+q)(1+r)(1+s) - 2qrs = q+qr$; $(q, r, s) = (2, 5, 3)$ ($m = 30$)
 - (iii) $(1+q)(1+r)(1+s) - 2qrs = 1+q+rs$; $(q, r, s) = (5, 2, 3), (5, 3, 2)$ ($m = 30$)
 - (iv) $(1+q)(1+r)(1+s) - 2qrs = q+r+s$; $(q, r, s) = (2, 3, 7), (2, 7, 3), (3, 2, 7), (3, 7, 2), (7, 2, 3), (7, 3, 2)$ (i.e., $m = 42$).
- (5) Suppose m is a k -near-perfect number for some $k \geq 0$ and $\tau(m) = 6$. Then $k \in \{0, 1, 2, 3, 4, 5\}$ and Lemma 2.6 implies that m is of the form q^2r with q, r being distinct primes.

When $k = 0$, $m = 28$. When $k = 1$, by Lemma 2.9 we have $m \in \{12, 18, 20\}$. For $2 \leq k \leq 5$, consider the following Diophantine equations:

$$\begin{aligned} \sigma(m) - 2m &= (1+q+q^2)(1+r) - 2q^2r \\ &\in \{1+q, 1+q^2, 1+r, 1+qr, q+q^2, q+r, q+qr, q^2+r, q^2+qr, \\ &\quad r+qr, \quad (k=2) \\ &\quad 1+q+q^2, 1+q+r, 1+q+qr, 1+q^2+r, 1+q^2+qr, 1+r+qr, \\ &\quad q+q^2+r, q+q^2+qr, q+r+qr, q^2+r+qr, \quad (k=3) \\ &\quad 1+q+q^2+r, 1+q+q^2+qr, 1+q+r+qr, 1+q^2+r+qr, \\ &\quad q+q^2+r+qr, \quad (k=4) \\ &\quad 1+q+q^2+r+qr \quad (k=5)\}. \end{aligned}$$

We may express r in terms of q easily:

$$\begin{aligned} r &= 1 + \frac{q+1}{q^2-q-1}, \frac{q}{q^2-q-1}, 1 + \frac{2}{q-1}, 1 + \frac{1}{q-1}, \frac{1}{q^2-q-1}, 1 - \frac{q+1}{q^2+1}, \\ &\quad 1 + \frac{2}{q^2-1}, \frac{q+1}{q^2-q}, \frac{1}{q-1}, 1 + \frac{1+q}{q^2} \quad (k=2), \\ &\quad 0, 1 + \frac{1}{q-1}, 1 + \frac{1}{q^2-1}, \frac{1}{q-1}, \frac{q}{q^2-1}, 1 + \frac{1}{q}, \frac{1}{q^2-q}, \frac{1}{q^2-1}, 1 + \frac{1}{q^2}, \frac{q+1}{q^2}, \\ &\quad (k=3), \\ &\quad 0, 0, 1, \frac{1}{q}, \frac{1}{q^2} \quad (k=4), \\ &\quad 0 \quad (k=5). \end{aligned}$$

The solvabilities of the equations are now apparent as only for small q 's the expressions are possibly integral. Also recall the restriction that q, r have to be distinct primes. Thus, only

$$(1+q+q^2)(1+r) - 2q^2r = 1+r$$

has solutions and $(q, r) = (2, 3), (3, 2)$, which correspond to $m = 12, 18$.

- (6) Suppose m is a k -near-perfect number for some $k \geq 0$ and $\tau(m) = 4$. Then $k \in \{0, 1, 2, 3\}$ and Lemma 2.6 implies that m is of the form qr , where q, r are

distinct primes. By also noting the symmetry of q and r , it suffices to consider the following Diophantine equations:

$$\begin{aligned}\sigma(m) - 2m &= (1+q)(1+r) - 2qr \\ &\in \{0, 1, q, 1+q, q+r, 1+q+r\}.\end{aligned}$$

Simply expand the above equations, we have

$$1+q+r = qr, \quad r = q(r-1), \quad 1 = r(q-1), \quad r = 2qr, \quad 1 = qr, \quad qr = 0$$

respectively. Each of these equations are now straight-forward to solve and the only possible solution is $m = 6$.

□

The proof of Theorem 1.3 rests on the study of the equation $\sigma(n) = \ell n + k$ which is carried out by a number of authors in the past decades; for more detail, see [AnPoPo, Po1, Po2, Po3, PoPo, PoSh, PoPoTh]. In this article, we only need the case of $\ell = 2$ and adopt following definitions from the aforementioned literature.

Definition 2.11 (Regular / Sporadic Solutions). *The solutions of $\sigma(n) = 2n + k$ of the form*

$$n = pm', \text{ where } p \nmid m', \sigma(m') = 2m', \sigma(m') = k, \quad (2.22)$$

are called regular. All other solutions are called sporadic.

Lemma 2.12. *Let $x \geq 3$ and k be an integer. The number of sporadic solutions $n \leq x$ to $\sigma(n) = 2n + k$ is at most $x^{3/5+o(1)}$ as $x \rightarrow \infty$, uniformly in k .*

Proof. See [PoPoTh] Theorem 4.4.

□

3. OUTLINE OF THEOREM 1.1 AND 1.2

Let us first recall the settings in [PoSh]. In order to estimate the size of the set $N(k; x)$, one may partition it into the following three subsets and estimate each respectively:

$$\begin{aligned}N_1(k; x) &:= \{n \in N(k; x) : P^+(n) \leq y\}, \\ N_2(k; x) &:= \{n \in N(k; x) : P^+(n) > y \text{ and } P^+(n)^2 | n\}, \\ N_3(k; x) &:= \{n \in N(k; x) : P^+(n) > y \text{ and } P^+(n) \nmid n\},\end{aligned} \quad (3.1)$$

where we shall remark on the choice of $y = y(x)$ at the end of this section.

In [PoSh] they further partitioned $N_3(k; x)$ according to whether $\tau(m)$ is at most k or not. They bounded the contribution from $\tau(m) \leq k$ simply by $\frac{x}{\log x} (\log \log x)^{k-1}$, i.e., Lemma 2.1. Instead, if one considers the normal order of $\log \tau(n)$, which is $(\log 2) \log \log n$, one obtains the bound $\frac{x}{\log x} (\log \log x)^{\lfloor \frac{\log k}{\log 2} \rfloor}$ for $N(k; x)$. More work is needed, though, as this is still not the correct order for $\#N(k; x)$; we thus have to partition $N_3(k; x)$ more carefully. This is explained as follows.

Definition 3.1. *Suppose $n = pm$ with $p > P^+(m)$. For k -near-perfect number n , there exists a set of proper divisors D_n of n with $\#D_n \leq k$ such that*

$$\sigma(n) = 2n + \sum_{d \in D_n} d. \quad (3.2)$$

We define the following associated sets:

$$\begin{aligned} D_n^{(1)} &:= \{d \in D_n : p \nmid d\}, \\ D_n^{(2)} &:= \{d/p : d \in D_n, p \mid d\}. \end{aligned} \quad (3.3)$$

It is clear that $D_n^{(1)}$ and $D_n^{(2)}$ consists of positive divisors and proper divisors of m respectively.

Proposition 3.2. *With the same settings in Definition 3.1, n is k -near-perfect and the set $D_n^{(1)}$ consists of all positive divisors of m if and only if $\tau(m) \leq k$ and $m \in N(k - \tau(m))$.*

Proof. (\Rightarrow): Immediately from the assumptions,

$$\begin{aligned} (1+p)\sigma(m) &= \sigma(pm) = 2pm + \sum_{d \in D_n^{(1)}} d + p \sum_{d \in D_n^{(2)}} d \\ &= 2pm + \sigma(m) + p \sum_{d \in D_n^{(2)}} d. \end{aligned} \quad (3.4)$$

This implies

$$\sigma(m) = 2m + \sum_{d \in D_n^{(2)}} d. \quad (3.5)$$

Since $\#D_n^{(1)} = \tau(m)$ and $\#D_n^{(1)} + \#D_n^{(2)} = \#D_n \leq k$, we have $\tau(m) \leq k$, $\#D_n^{(2)} \leq k - \tau(m)$ and $m \in N(k - \tau(m))$.

(\Leftarrow): There exists a set of proper divisors D_m of m with $\#D_m \leq k - \tau(m)$ such that

$$\sigma(m) = 2m + \sum_{d \in D_m} d.$$

Then

$$\sigma(n) = (1+p)\sigma(m) = \sum_{d|m} d + p \left(2m + \sum_{d \in D_m} d \right) = 2n + \sum_{d|m} d + p \sum_{d \in D_m} d. \quad (3.6)$$

Now,

$$\{d \mid m\} \cup \{pd : d \in D_m\}$$

is a set of proper divisors of n with at most $\tau(m) + (k - \tau(m)) = k$ elements. Thus, n is a k -near-perfect number. Also, $D_n^{(1)} = \{d \mid m\}$ and $D_n^{(2)} = D_m$. \square

To facilitate discussion that follows, we introduce the following notations:

Definition 3.3.

$$\begin{aligned} N_3^{(1)}(k; x) &:= \{n \leq x : n = pm, p > \max\{y, P^+(m)\}, \tau(m) \leq k \\ &\quad \text{and } m \in N(k - \tau(m))\}, \end{aligned} \quad (3.7)$$

$$N_3^{(2)}(k; x) := N_3(k; x) \setminus N_3^{(1)}(k; x), \quad (3.8)$$

$$M(k) := \left\{ n \in N(k) : n = pm, p > P^+(m), D_n^{(1)} \subsetneq \{d \mid m\} \right\} \quad (3.9)$$

$$M(k; x) := M(k) \cap [1, x]. \quad (3.10)$$

We carry out the above partition into $N_1, N_2, N_3^{(1)}, N_3^{(2)}$ recursively in Section 5. At each step, we show that the contributions from $N_1, N_2, N_3^{(2)}$ are of acceptable sizes,⁵ whereas the description for $N_3^{(1)}$ allows us to move onto the next step in the recursive process. After this is done, we only have to apply Lemma 2.10, i.e., the determination of k -near-perfect numbers for small integers k with a fixed number of positive divisors. In this way, we improve upon the bound $\frac{x}{\log x}(\log \log x)^{\lfloor \frac{\log k}{\log 2} \rfloor}$ and establish Theorem 1.1. For more detail, see Section 5.1. As a by-product, we are able to deduce the precise asymptotic formulae for $4 \leq k \leq 11$ in Theorem 1.2.

The proof of Theorem 1.2 is simpler than that of Theorem 1.1. It follows quite directly from the partition as in Theorem 1.1 without encountering complications of the recursive process. We shall start with its proof and briefly recall the essential estimates done in [PoSh] in the next section (Section 4).

Finally, we would also like to make a remark on the choice of the parameter y . In [PoSh], they chose $y = x^{\frac{1}{4 \log \log x}}$ for their applications. However, this is not admissible in the proof of Theorem 1.1. Firstly, it is clear that the choice of [PoSh] does not satisfy the conditions in Lemma 2.4 (j will be chosen in terms of k in Section 5.5 and j grows with k). Secondly, in order to make sure the estimate in Lemma 2.4 is of acceptable sizes with respect to Theorem 1.1, i.e., smaller than $\frac{x}{\log x}(\log \log x)^{\lfloor \frac{\log(k+4)}{\log 2} \rfloor - 3}$, it is essential to choose $y = (\log x)^\alpha$ for some $\alpha > 0$. Thirdly, α needs to be large enough so that the contribution of $N_3^{(2)}$ is acceptable, see Section 5.4. We shall see $\alpha = 3k + 10$ is good enough. We shall stick with this choice of y in Section 4.2 and 5. For Section 4.3, however, we must choose a different y there for better estimates.

4. PROOF OF THEOREM 1.2

4.1. We shall review the argument of [PoSh] in this subsection for the convenience of readers.

The estimations for $N_1(k; x)$ and $N_2(k; x)$ are straight-forward. Indeed,

$$\#N_1(k; x) \leq \#\Phi(x, y) \quad (4.1)$$

and

$$\#N_2(k; x) \leq \#\{n \leq x : P^+(n) > y, P^+(n)^2 \mid n\} \leq \sum_{p > y} \frac{x}{p^2} \ll \frac{x}{y}. \quad (4.2)$$

Suppose $n = pm \in N_3^{(2)}(k; x)$. For the counting argument below, we shall also assume that

$$\tau(m) \leq (\log x)^3. \quad (4.3)$$

This is acceptable because

$$\#\{n \leq x : \tau(n) > 2(\log x)^3\} \ll \frac{x}{(\log x)^2}. \quad (4.4)$$

This follows from $2\tau(m) = \tau(n)$ and the crude estimate

$$2(\log x)^3 \cdot \#\{n \leq x : \tau(n) > 2(\log x)^3\} \leq \sum_{n \leq x} \tau(n) \ll x \log x.$$

⁵We say that the size of a quantity is *acceptable* if it is not greater than that of the main term, e.g., $\frac{x}{\log x}(\log \log x)^{\lfloor \frac{\log(k+4)}{\log 2} \rfloor - 3}$ in Theorem 1.1 and $x/\log x$ in Theorem 1.2.

In the following, we count the number of possible p 's such that $pm \in N_3^{(2)}(k; x)$ for each $m \leq x/y$. Since $n = pm$ is k -near-perfect,

$$(1+p)\sigma(m) = 2pm + \sum_{d \in D_n^{(1)}} d + p \sum_{d \in D_n^{(2)}} d, \quad (4.5)$$

where the sets $D_n^{(1)}$ and $D_n^{(2)}$ are defined in Definition 3.1. Reducing both sides (mod p) yields

$$p \mid \left(\sigma(m) - \sum_{d \in D_n^{(1)}} d \right). \quad (4.6)$$

By (3.8) and Proposition 3.2,

$$\sigma(m) - \sum_{d \in D_n^{(1)}} d > 0.$$

Moreover,

$$\sigma(m) - \sum_{d \in D_n^{(1)}} d \leq \sigma(m) \ll m \log \log m \leq x \log \log x.$$

Thus, the number of prime factors of $\left(\sigma(m) - \sum_{d \in D_n^{(1)}} d \right)$ is

$$O(\log x). \quad (4.7)$$

Since $D_n^{(1)} \subset \{d \mid m\}$ and $\#D_n^{(1)} \leq k$, the number of possible values for $\left(\sigma(m) - \sum_{d \in D_n^{(1)}} d \right)$ is

$$\leq (1 + \tau(m))^k \ll_k (\log x)^{3k} \quad (4.8)$$

by (4.3). As a result, from (4.6), (4.7) and (4.8), the number of possible p 's is

$$\ll_k (\log x)^{3k+1}. \quad (4.9)$$

From this, we conclude that

$$\#N_3^{(2)}(k; x) \ll_k \frac{x}{y} (\log x)^{3k+1} + \frac{x}{(\log x)^2}. \quad (4.10)$$

4.2. Throughout this subsection, we take $y := (\log x)^{3k+10}$.

By (2.4) of Lemma 2.2, there exists a constant $x_1(k) > 0$ such that for any $x \geq x_1(k)$,

$$\#N_1(k; x) \ll x \exp \left(-\frac{1}{2(3k+10)} \frac{\log x}{\log \log x} \right) \leq \frac{x}{(\log x)^2}. \quad (4.11)$$

Immediately from (4.2) and (4.10),

$$\#N_2(k; x), \#N_3^{(2)}(k; x) \ll_k \frac{x}{(\log x)^2}. \quad (4.12)$$

Thus, the contributions from $N_1(k; x)$, $N_2(k; x)$ and $N_3^{(2)}(k; x)$ are acceptable.

It remains to consider $n \in N_3^{(1)}(k; x)$, i.e., $n = pm \leq x$, $p > \max\{y, P^+(m)\}$ and $m \in N(k - \tau(m))$. When $4 \leq k \leq 11$, there are only finitely many such m and they

have been completely determined in Lemma 2.10. Thus, by the Prime Number Theorem, we have

$$\#N_3^{(1)}(k; x) \sim \sum_{r=4}^k \sum_{\substack{\tau(m)=r \\ m \in N(k-r)}} \pi(x/m) \sim \left(\sum_{r=4}^k \sum_{\substack{\tau(m)=r \\ m \in N(k-r)}} \frac{1}{m} \right) \frac{x}{\log x}$$

as $x \rightarrow \infty$. Explicitly, the constant

$$c_k := \sum_{r=4}^k \sum_{\substack{\tau(m)=r \\ m \in N(k-r)}} \frac{1}{m} \quad (4.13)$$

is equal to

$$\begin{aligned} c_4 &= c_5 = \frac{1}{6}, \quad c_6 = \frac{17}{84} = \frac{1}{6} + \frac{1}{28} \approx 0.2024, \\ c_7 &= c_8 = \frac{493}{1260} = \frac{1}{6} + \frac{1}{12} + \frac{1}{18} + \frac{1}{20} + \frac{1}{28} \approx 0.3913, \\ c_9 &= \frac{1}{6} + \frac{1}{12} + \frac{1}{18} + \frac{1}{20} + \frac{1}{28} + \frac{1}{24} + \frac{1}{40} + \frac{1}{56} + \frac{1}{88} + \frac{1}{104} \\ &= \frac{179017}{360360} \approx 0.4968, \\ c_{10} &= \frac{1}{6} + \frac{1}{12} + \frac{1}{18} + \frac{1}{20} + \frac{1}{28} + \frac{1}{24} + \frac{1}{40} + \frac{1}{56} + \frac{1}{88} + \frac{1}{104} \\ &\quad + \frac{1}{30} + \frac{1}{54} + \frac{1}{66} + \frac{1}{196} + \frac{1}{496} \\ &= \frac{267857123}{469188720} \approx 0.5709, \\ c_{11} &= \frac{1}{6} + \frac{1}{12} + \frac{1}{18} + \frac{1}{20} + \frac{1}{28} + \frac{1}{24} + \frac{1}{40} + \frac{1}{56} + \frac{1}{88} + \frac{1}{104} \\ &\quad + \frac{1}{30} + \frac{1}{54} + \frac{1}{66} + \frac{1}{196} + \frac{1}{496} + \frac{1}{42} + \frac{1}{36} + \frac{1}{368} + \frac{1}{464} \\ &= \frac{196329752441}{312948876240} \approx 0.6274. \end{aligned}$$

This completes the proof of Theorem 1.2.

4.3. Before we end this section, we would like to follow-up on a remark of [PoSh] (pp. 3044) where they claimed the result

$$\#N(k; x) \ll x \exp(-(c_k + o(1))\sqrt{\log x \log \log x}), \quad (4.14)$$

for $k = 2, 3$, where $c_2 = \sqrt{6}/6 \approx 0.4082$ and $c_3 = \sqrt{2}/4 \approx 0.3535$. In view of the discussion in Section 3, the reason for a much smaller estimates for $k = 2, 3$ lies in the the nonexistence of near perfect numbers of the form p or p^2 where p is a prime (see Lemma 2.6). This implies $N_3^{(1)}(3; x)$ (see 3.7) is an empty set .

Since a complete argument for (4.14) was not given in [PoSh], we supply more detail here and hope it will be helpful for the interested readers. The argument below actually shows that one can take c_2 and c_3 to be $1/\sqrt{2} \approx 0.7071$, but this improvement is not substantial.

Here, it is essential to apply a more precise count of $\#\Phi(x, y)$ than the one given in Lemma 2.2. From Theorem 9.15 and Corollary 9.18 of [DeKLu], we have

$$\#\Phi(x, y) = x \exp(-u \log u + O(u \log \log u)) \quad (4.15)$$

and this is uniform for

$$(\log x)^3 \leq y \leq x, \quad (4.16)$$

where

$$u := \frac{\log x}{\log y}. \quad (4.17)$$

In this subsection, we shall choose a different $y = y(x)$ from the one taken in the rest of this article (i.e., $y = (\log x)^{3k+10}$).

We modify the estimations for $N_3^{(2)}(k; x)$ sketched in Section 4.1 slightly:

$$\begin{aligned} & \#N_3^{(2)}(k; x) \\ & \leq \sum_{m \leq \frac{x}{y}} \# \left\{ p \leq \frac{x}{m} : p \mid \left(\sigma(m) - \sum_{d \in D_{pm}^{(1)}} d \right) \text{ and } \sigma(m) - \sum_{d \in D_{pm}^{(1)}} d > 0 \right\} \\ & \ll_k \sum_{m \leq \frac{x}{y}} (\log x) \tau(m)^k \\ & \ll \frac{x}{y} (\log x)^{2^k}. \end{aligned} \quad (4.18)$$

For a proof of the last estimate, see [MV] eq. (2.31), pp. 61. Also, compare this with (4.10). We shall see shortly it is a better estimate in the case of $k = 2, 3$ with a new choice of y .

Therefore for $k = 2, 3$, by (4.15), (4.1), (4.2), (4.18) and the fact that $N_3^{(2)}(k; x) = \emptyset$, we have

$$\begin{aligned} \#N(k; x) &= \#N_1(k; x) + \#N_2(k; x) + \#N_3^{(2)}(k; x) \\ &\ll_k x \exp(-u \log u + O(u \log \log u)) + \frac{x}{y} (\log x)^{2^k}. \end{aligned} \quad (4.19)$$

We optimize the last estimate by setting

$$u \log u = \log y - 2^k \log \log x. \quad (4.20)$$

A good approximation for u satisfying (4.20) is

$$u^2 = 2 \frac{\log x}{\log \log x}, \quad (4.21)$$

i.e.,

$$\log y = \frac{1}{\sqrt{2}} \sqrt{\log x \log \log x}. \quad (4.22)$$

It is to see it satisfies the requirement (4.16). Plugging this into (4.19), we have

$$\#N(k; x) \ll x \exp \left(-\frac{1}{\sqrt{2}} \left\{ 1 + O \left(\frac{\log \log \log x}{\log \log x} \right) \right\} \sqrt{\log x \log \log x} \right) \quad (4.23)$$

for $k = 2, 3$.

5. PROOF OF THEOREM 1.1

5.1. Outline. Throughout this section, we fix $y := (\log x)^{3k+10}$ and denote by $T_r(x)$ the set of natural numbers in $[1, x]$ of the form $p_1 \cdots p_r m_r$ with $p_1 > \cdots > p_r > \max\{y, P^+(m_r)\}$.

The estimates required in Step 0 are sketched in Section 4.1:

$$\#N_1(k; x), \#N_2(k; x), \#N_3^{(2)}(k; x) \ll_k \frac{x}{(\log x)^2}, \quad (5.1)$$

which are of acceptable sizes. Therefore by (3.7), it now suffices to consider the set

$$\left\{ n \in T_1(x) : \tau(m_1) \leq k, m_1 \in N \left(k - \tau(m_1); \frac{x}{y} \right) \right\}. \quad (5.2)$$

In Step 1, we estimate the size of the set 5.2 by repeating the partition to m_1 , i.e., consider

$$\left\{ n \in T_1(x) : \tau(m_1) \leq k, m_1 \in R \left(k - \tau(m_1); \frac{x}{y} \right) \right\} \quad (5.3)$$

for $R = N_1, N_2, N_3^{(1)}, N_3^{(2)}$.

- When $R = N_1, N_2, N_3^{(2)}$, the sets (5.3) will be shown to be of acceptable sizes $O(\frac{x}{\log x} \log \log x)$.
- When $R = N_3^{(1)}$, recall from (3.7) that the condition $m_1 \in N_3^{(1)}(k - \tau(m_1); x/y)$ refers to

$$\begin{aligned} m_1 &= p_2 m_2 \leq \frac{x}{y}, \quad p_2 > \max\{y, P^+(m_2)\}, \\ \tau(m_2) &\leq k - \tau(m_1), \quad m_2 \in N(k - \tau(m_1 - \tau(m_2))). \end{aligned} \quad (5.4)$$

Thus, the set (5.3) is indeed equal to

$$\left\{ n \in T_2(x) : \tau(m_2) \leq \frac{k}{3}, m_2 \in N \left(k - 3\tau(m_2); \frac{x}{y^2} \right) \right\}. \quad (5.5)$$

In Step 2, we estimate the size of (5.5) by repeating the partition to m_2 , so on and so forth. More generally at Step $j-1$, we arrive at the tasks of showing the sizes of the sets

$$\left\{ n \in T_{j-1}(x) : \tau(m_{j-1}) \leq \frac{k}{2^{j-1}-1}, m_{j-1} \in R \left(k - (2^{j-1}-1)\tau(m_{j-1}); \frac{x}{y^{j-1}} \right) \right\} \quad (5.6)$$

being $O_j \left(\frac{x}{\log x} (\log \log x)^{j-1} \right)$ for $R = N_1, N_2, N_3^{(2)}$. This will be done in Sections 5.2, 5.3 and 5.4.

The recursion ends once we hit an k -admissible integer j_0 .

Definition 5.1. An integer $j_0 \geq 1$ is said to be k -admissible if

$$1 \leq \# \left\{ m_{j_0} \in \mathbb{N} : \tau(m_{j_0}) \leq \frac{k}{2^{j_0}-1}, m_{j_0} \in N(k - (2^{j_0}-1)\tau(m_{j_0})) \right\} < \infty. \quad (5.7)$$

Remark 5.2. In view of Lemma 2.10(1)(6), (5.7) is equivalent to

$$\# \left\{ m_{j_0} \in \mathbb{N} : 4 \leq \tau(m_{j_0}) \leq \frac{k}{2^{j_0}-1}, m_{j_0} \in N(k - (2^{j_0}-1)\tau(m_{j_0})) \right\} < \infty. \quad (5.8)$$

It follows from Lemma 2.1 that

$$\begin{aligned}
& \# \left\{ n \in T_{j_0-1}(x) : \tau(m_{j_0-1}) \leq \frac{k}{2^{j_0-1}-1}, \right. \\
& \quad \left. m_{j_0-1} \in N_3^{(1)} \left(k - (2^{j_0-1}-1)\tau(m_{j_0-1}); \frac{x}{y^{j_0-1}} \right) \right\} \\
& \leq \# \left\{ n \in T_{j_0}(x) : \tau(m_{j_0}) \leq \frac{k}{2^{j_0}-1}, m_{j_0} \in N(k - (2^{j_0}-1)\tau(m_{j_0})) \right\} \\
& \ll_{j_0} \frac{x}{\log x} (\log \log x)^{j_0-1}
\end{aligned} \tag{5.9}$$

for an k -admissible integer j_0 . By showing that

$$j_0 = j_0(k) := \left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 2 \tag{5.10}$$

is k -admissible in Section 5.5, and together with the estimates (5.13), (5.14), (5.18) and (5.9), we have the upper bound in Theorem 1.1.

The lower bound simply follows from the observation

$$6p_1 \cdots p_s = p_1 \cdots p_s + 2p_1 \cdots p_s + 3p_1 \cdots p_s,$$

where $p_1 > \cdots > p_s > 3$ are primes. Thus, $6p_1 \cdots p_s$ is a k_s -near-perfect numbers with

$$k_s := \tau(6p_1 \cdots p_s) - 1 - 3 = 2^{s+2} - 4.$$

Fix any integer $k \geq 4$. Take the largest integer $s \geq 1$ such that $k_s \leq k$, i.e.,

$$s = \left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 2.$$

Then by Lemma 2.1 one has

$$\begin{aligned}
\#N(k; x) & \geq \# \{ 6p_1 \cdots p_s \leq x : p_1 > \cdots > p_s > 3 \} \\
& \gg_k \frac{x}{\log x} (\log \log x)^{\left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 3}.
\end{aligned} \tag{5.11}$$

5.2. Estimation for $R = N_1$. In view of our claim that

$$j_0 = \left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 2$$

is an k -admissible integer, we apply Lemma 2.4 for

$$2 \leq j \leq j_0.$$

In order to meet the assumptions of Lemma 2.4, we restrict to $x \geq x_0(k)$, where $x_0(k) > 0$ is a large constant such that for $x \geq x_0(k)$,

$$(\log x)^{3k+10} \leq x^{\frac{1}{4j_0 \log \log x}}. \tag{5.12}$$

Hence for $x \geq x_0(k)$ and $2 \leq j \leq j_0$, we have

$$\begin{aligned}
& \# \left\{ n \in T_{j-1}(x) : \tau(m_{j-1}) \leq \frac{k}{2^{j-1}-1}, \right. \\
& \quad \left. m_{j-1} \in N_1 \left(k - (2^{j-1}-1)\tau(m_{j-1}); \frac{x}{y^{j-1}} \right) \right\}
\end{aligned}$$

being bounded by

$$\#\Phi_{j-1}(x, y) \ll_j \frac{x}{\log x} (\log \log x)^{j-1} \ll_k \frac{x}{\log x} (\log \log x)^{j_0-1}. \quad (5.13)$$

5.3. Estimation for $R = N_2$. From our previous analysis, for $2 \leq j \leq j_0$ we have

$$\begin{aligned} & \# \left\{ n \in T_{j-1}(x) : m_{j-1} \in N_2 \left(k - (2^{j-1} - 1)\tau(m_{j-1}); \frac{x}{y^{j-1}} \right) \right\} \\ & \leq \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq x}} \sum_{\substack{m_{j-1} \leq x/p_1 \dots p_{j-1} \\ P^+(m_{j-1})^2 | m_{j-1} \\ P^+(m_{j-1}) > y}} 1 \\ & \leq \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq x}} \frac{x}{y} \frac{1}{p_1 \dots p_{j-1}} \\ & \leq \frac{x}{y} \left(\sum_{p \leq x} \frac{1}{p} \right)^{j-1} \ll \frac{x}{y} (\log \log x)^{j-1} \ll \frac{x}{(\log x)^{3k+10}} (\log \log x)^{j_0-1}. \quad (5.14) \end{aligned}$$

5.4. Estimation of $R = N_3^{(2)}$. Recall the notations $M(k)$ and $M(k; x)$ introduced in Section 3. From the argument sketched in Section 4.1, we have

$$\#\{n \leq x : n \in M(k), P^+(n) > y\} \ll_k \frac{x}{y} (\log x)^{3k+1}. \quad (5.15)$$

Then

$$\begin{aligned} \#M(k; x) &= \#\{n \leq x : n \in M(k), P^+(n) \leq y\} + \#\{n \leq x : n \in M(k), P^+(n) > y\} \\ &\ll_k \#\Phi(x, y) + \frac{x}{y} (\log x)^{3k+1} \ll_k \frac{x}{(\log x)^2}. \quad (5.16) \end{aligned}$$

It follows from partial summation that

$$\sum_{n \in M(k)} \frac{1}{n} < \infty. \quad (5.17)$$

Therefore for $2 \leq j \leq j_0$, by applying (5.16), (5.17) and Lemma 2.1, we have

$$\begin{aligned}
& \# \left\{ n \in T_{j-1}(x) : m_{j-1} \in N_3^{(2)} \left(k - (2^{j-1} - 1)\tau(m_{j-1}); \frac{x}{y^{j-1}} \right) \right\} \\
& \leq \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq \sqrt{x}}} \sum_{\substack{m_{j-1} \leq x/p_1 \dots p_{j-1} \\ m_{j-1} \in M(k)}} 1 + \sum_{\substack{m_{j-1} \leq \sqrt{x} \\ m_{j-1} \in M(k)}} \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq x/m_{j-1}}} 1 \\
& \ll_k \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq \sqrt{x}}} \frac{\frac{x}{p_1 \dots p_{j-1}}}{(\log \frac{x}{p_1 \dots p_{j-1}})^2} + \sum_{\substack{m_{j-1} \leq \sqrt{x} \\ m_{j-1} \in M(k)}} \frac{\frac{x}{m_{j-1}}}{\log \frac{x}{m_{j-1}}} \left(\log \log \frac{x}{m_{j-1}} \right)^{j-2} \\
& \ll \frac{x}{(\log x)^2} \sum_{\substack{p_1 > \dots > p_{j-1} > y \\ p_1 \dots p_{j-1} \leq \sqrt{x}}} \frac{1}{p_1 \dots p_{j-1}} + \frac{x}{\log x} (\log \log x)^{j-2} \sum_{\substack{m_{j-1} \leq \sqrt{x} \\ m_{j-1} \in M(k)}} \frac{1}{m_{j-1}} \\
& \ll \frac{x}{(\log x)^2} \left(\sum_{p \leq \sqrt{x}} \frac{1}{p} \right)^{j-1} + \frac{x}{\log x} (\log \log x)^{j-2} \\
& \ll \frac{x}{\log x} (\log \log x)^{j_0-2}. \tag{5.18}
\end{aligned}$$

5.5. Analyzing $R = N_3^{(1)}$. We consider the following cases:

(1)

$$4 \cdot (2^s - 1) \leq k < 8 \cdot (2^s - 1) \tag{5.19}$$

for some $s > 1$,

(2)

$$k = 8 \cdot 2^s - \ell \tag{5.20}$$

with either

(a) $s \geq 3$ and $\ell \in \{5, 6, 7, 8\}$, or

(b) $s = 2$ and $\ell \in \{6, 7, 8\}$.

(3)

$$k = 27 \tag{5.21}$$

(i.e., $s = 2$ and $\ell = 5$ in (5.20)).

It is clear that the above covers all integers $k \geq 12$.⁶ In any case, we have

$$s = \left\lfloor \frac{\log(\frac{k}{4} + 1)}{\log 2} \right\rfloor = \left\lfloor \frac{\log(k + 4)}{\log 2} \right\rfloor - 2. \tag{5.22}$$

Indeed for Case (1),

$$\frac{\log(\frac{k}{8} + 1)}{\log 2} < s \leq \frac{\log(\frac{k}{4} + 1)}{\log 2}$$

and

$$0 < \frac{\log(\frac{k}{4} + 1)}{\log 2} - \frac{\log(\frac{k}{8} + 1)}{\log 2} < 1.$$

Hence,

$$s = \left\lfloor \frac{\log(\frac{k}{4} + 1)}{\log 2} \right\rfloor = \left\lfloor \frac{\log(k + 4)}{\log 2} \right\rfloor - 2. \tag{5.23}$$

⁶Note that the cases $4 \leq k \leq 11$ have been settled in Theorem 1.2.

For Cases (2) and (3), we have

$$s + 2 < \frac{\log(k+4)}{\log 2} < s + 3,$$

and

$$s = \left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 2. \quad (5.24)$$

Thus, the upper bound in Theorem 1.1 would follow if we establish the claim that s is an k -admissible integer in each case.

(1) Suppose

$$4 \cdot (2^s - 1) \leq k < 8 \cdot (2^s - 1) \quad (5.25)$$

for some $s > 1$. From Lemma 2.10(5)(6), the facts that

$$4 \leq \tau(m_s) \leq \frac{k}{2^s - 1} < 8$$

and m_s being near-perfect, we have $m_s \in \{6, 12, 28, 20, 28\}$. Thus, s is an k -admissible integer.

(2) Suppose

$$k = 8 \cdot 2^s - \ell \quad (5.26)$$

with either

(a) $s \geq 3$ and $\ell \in \{5, 6, 7, 8\}$; or

(b) $s = 2$ and $\ell \in \{6, 7, 8\}$.

In both cases, we have

$$4 \leq \tau(m_s) \leq \frac{k}{2^s - 1} < 9.$$

If $4 \leq \tau(m_s) < 8$, then $m_s \in \{6, 12, 28, 20, 28\}$ as in the previous case. Now suppose $\tau(m_s) = 8$. Then

$$\begin{aligned} m_s &\in N(k - (2^s - 1)\tau(m_s)) \\ &= N((2^s - 1)(8 - \tau(m_s)) + (8 - \ell)) \\ &= N(8 - \ell) \subset N(3). \end{aligned}$$

By Lemma 2.10 (4), $m_s \in \{24, 40, 56, 88, 104, 30, 54, 66, 42\}$. As a result, s is an k -admissible integer.

(3) Suppose $k = 27$. Then

$$4 \leq \tau(m_2) \leq 9.$$

If $4 \leq \tau(m_2) \leq 8$, then $m_2 \in \{6, 12, 28, 20, 28\} \cup \{24, 40, 56, 88, 104, 30, 54, 66, 42\}$ as in the previous cases. When $\tau(m_2) = 9$, $m_2 \in N(0)$ and there is no such m_2 by Lemma 2.10 (3). Hence, 2 is 27-admissible.

This completes the proof of

$$N(k; x) \ll_k \frac{x}{\log x} (\log \log x)^{\left\lfloor \frac{\log(k+4)}{\log 2} \right\rfloor - 3} \quad (5.27)$$

for any $k \geq 4$ and hence the proof of Theorem 1.1.

6. PROOF OF THEOREM 1.3

Let $\epsilon \in (0, 2/5)$. By Lemma 2.12,

$$\begin{aligned} \#(E(k; x) \setminus E_\epsilon(k; x)) &\leq \#\{n \leq x : n \in E(k), n = pm', p \nmid m', \sigma(m') = 2m'\} \\ &\quad + O(x^{3/5+\epsilon+o(1)}). \end{aligned} \quad (6.1)$$

For $n \in E(k)$ with $n = pm'$, $p \nmid m'$ and $\sigma(m') = 2m'$, we have

$$pm' = \sum_{d_1 \in D_1} d_1 + p \sum_{d_2 \in D_2} d_2, \quad (6.2)$$

where D_1 is a subset of positive divisors of m' , D_2 is a subset of proper divisors of m' with $\#D_1 + \#D_2 = \tau(pm') - 1 - k = 2\tau(m') - 1 - k$.

Suppose that $D_1 \neq \emptyset$. Then

$$1 \leq \sum_{d_1 \in D_1} d_1 \leq \sigma(m') = 2m'. \quad (6.3)$$

Reducing (6.2) modulo p , we have

$$p \mid \sum_{d_1 \in D_1} d_1. \quad (6.4)$$

The number of possible values for p is $O(\log 2m') = O(\log x)$. Thus the number of possible values for such n is $O(x^{o(1)} \log x)$ by the Hornfeck-Wirsing Theorem ([HoWi]), which is acceptable.

Now suppose that $D_1 = \emptyset$. Then $\#D_2 = 2\tau(m') - 1 - k$ and

$$m' = \sum_{d_2 \in D_2} d_2. \quad (6.5)$$

Since $\sigma(m') = 2m'$, we have $\#D_2 = \tau(m') - 1$. Therefore, $\tau(m') - 1 = 2\tau(m') - 1 - k$, i.e., $\tau(m') = k$.

By the hypothesis of non-existence of odd perfect number and the Euclid-Euler Theorem, we have $m' = 2^{q'-1}(2^{q'} - 1)$ for some Mersenne prime q' . So $k = \tau(m') = 2q' \in M$. Hence if $k \notin M$, then we have a contradiction and

$$\#(E(k; x) \setminus E_\epsilon(k; x)) = O(x^{o(1)} \log x) + O(x^{3/5+\epsilon+o(1)}) = O(x^{3/5+\epsilon+o(1)}). \quad (6.6)$$

It was shown in [PoSh], by using a form of the Prime Number Theorem of Drmota, Mauduit and Rivat, that for all large k the number of k -exactly-perfect numbers up to x is $\gg_k x / \log x$. Therefore

$$\frac{\#(E(k; x) \setminus E_\epsilon(k; x))}{\#E(k; x)} \ll_k \frac{\log x}{x^{2/5-\epsilon-o(1)}} \quad (6.7)$$

and

$$\lim_{x \rightarrow \infty} \frac{\#E_\epsilon(k; x)}{\#E(k; x)} = 1. \quad (6.8)$$

Remark 6.1. Suppose $k \in M$. Then $k = 2q$ for some Mersenne prime q . Let $m = 2^{q-1}(2^q - 1)$. Then $q' = q$ and so $m' = m$ in the above argument. By the Prime Number Theorem,

$$\limsup_{x \rightarrow \infty} \frac{\#(E(k; x) \setminus E_\epsilon(k; x))}{x / \log x} \leq \frac{1}{m}. \quad (6.9)$$

On the other hand, since m is perfect, the number of proper divisors of m is $\tau(m) - 1 = 2q - 1$. Hence pm is a sum of $2q - 1$ of its proper divisors. The number of proper divisors of pm is $\tau(pm) - 1 = 4q - 1$. So, pm is a sum of all of its proper divisors with exactly $(4q - 1) - (2q - 1) = 2q$ exceptions, i.e., $pm \in E(k)$. Clearly $\sigma(pm) - 2pm < (pm)^\epsilon$ if $p > (2m^{1-\epsilon})^{1/\epsilon}$ and $p \nmid m$. It follows that

$$\liminf_{x \rightarrow \infty} \frac{\#(E(k; x) \setminus E_\epsilon(k; x))}{x / \log x} \geq \frac{1}{m}. \quad (6.10)$$

As a result,

$$\lim_{x \rightarrow \infty} \frac{\#(E(k; x) \setminus E_\epsilon(k; x))}{x / \log x} = \frac{1}{m}. \quad (6.11)$$

REFERENCES

- [AnPoPo] A. Anavi and P. Pollack and C. Pomerance, *On congruences of the form $\sigma(n) \equiv a \pmod{n}$* , Int. J. Number Theory **9** (2012), 115-124.
- [BeEr] S. J. Benkoski and P. Erdős, *On weird and pseudoperfect numbers*, Math. Comp. **28** (1974), no.126, 617-623.
- [DeKLu] J-M De Koninck, F. Luca *Analytic Number Theory: Exploring the Anatomy of Integers*, Graduate Studies in Mathematics Vol. 134, American Mathematical Society.
- [Er1] P. Erdős, *On perfect and multiply perfect numbers*, Annali di Matematica Pura ed Applicata, **42** (1956), no. 1, 253-258.
- [Er2] P. Erdős, *Some Extremal Problems in Combinatorial Number Theory*, Mathematical Essays Dedicated to A. J. Macintyre, 123–133, Ohio Univ. Press, Athens, Ohio, 1970.
- [FoHa] K. Ford, H. Halberstam *The Brun-Hooley Sieve*, Journal of Number Theory, **81** (2000), 335-350.
- [Gu] R. K. Guy, *Unsolved problems in number theory*, third ed., Springer (2004), 74-75, 78.
- [HaWr] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, fourth ed., Oxford University Press, Oxford (1975), pp. 368-370.
- [HoWi] B. Hornfeck and E. Wirsing, *Über die Häufigkeit vollkommener Zahlen*, Math. Ann. **133** (1957), 431-438.
- [Ka] H. J. Kanold, *Über die Verteilung der vollkommene Zahlen und allgemeinerer Zahlenmengen*, Math Ann. **132** (1957), 442-450.
- [MV] H. L. Montgomery, R. C. Vaughan, *Multiplicative Number Theory I. Classical Theory*, Cambridge Studies in Advanced Mathematics no. 97 (2007), pp. 60-62
- [Niel] P. P. Nielsen, *Odd Perfect Numbers, Diophantine Equations, and Upper Bounds*, Math. Comp. **84** (2015), 2549-2567
- [Po1] C. Pomerance, *On composite n for which $\phi(n) \mid n - 1$* , Acta Arith. **28** (1976), 387-389.
- [Po2] C. Pomerance, *On composite n for which $\phi(n) \mid n - 1$, II*, Pacific J. Math. **69** (1977), 177-186.
- [Po3] C. Pomerance, *On the congruences $\sigma(n) \equiv a \pmod{n}$ and $n \equiv a \pmod{\phi(n)}$* , Acta Arith. **26** (1975), 265-272.
- [Pol] P. Pollack, *Not always buried deep: A second course in elementary number theory*, AMS, Providence (2009), 249, 258-259.
- [PoPo] P. Pollack, C. Pomerance, *On the distribution of some integers related to perfect and amicable numbers*, Colloq. Math. **30** (2013), 169-182.
- [PoPoTh] P. Pollack, C. Pomerance, L. Thompson, *Divisor-sum fibres*, Mathematika **64** (2018), 330-342.
- [PoSh] P. Pollack and V. Shevelev, *On perfect and near-perfect numbers*, J. Number Theory **132** (2012), 3037-3046.
- [ReCh] X.-Z. Ren and Y.-G. Chen *On near-perfect numbers with two distinct prime factors*, Bull. Aust. Math. Soc. **88** (2013), 520-524.
- [Sc] A. Schinzel, *On functions $\phi(n)$ and $\sigma(n)$* , Bull. Acad. Pol. Sci. Cl. III **3** (1955), 415-419.
- [Si] W. Sierpiński, *Sur les nombres pseudoparfais*, Mat. Vesnik **17** (1965), 212-213.
- [Ten] G. Tenenbaum, *Introduction to Analytic and Probabilistic Number Theory*, 3rd Ed., Graduate Studies in Mathematics **163**, American Mathematical Society (2015).
- [Vo] B. Volkmann, *A theorem on the set of perfect numbers*, Bull. A.M.S. **62** (1956), Abstract 180.

Email address: petercohen33@gmail.com

DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139

Email address: kcordwel@cs.cmu.edu

DEPARTMENT OF COMPUTER SCIENCE, CARNEGIE MELLON UNIVERSITY, PA 15213

Email address: wtgalyssa@gmail.com

STANFORD LAW SCHOOL, CA 94305

Email address: ck2854@math.columbia.edu

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK, NY 10027

Email address: adamlott99@math.ucla.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA 90095

Email address: sjml@williams.edu, Steven.Miller.MC.96@aya.yale.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267