

Benford's Law: Why the IRS might care about the $3x + 1$ problem and $\zeta(s)$

Steven J. Miller

`sjml@williams.edu,`

`Steven.Miller.MC.96@aya.yale.edu`

`http://web.williams.edu/Mathematics/sjmiller/public_html/`


UCF: 5-1-2026

Introduction

- A. Berger and T. P. Hill, *An Introduction to Benford's Law*, Princeton University Press, Princeton, 2015. See also <http://www.benfordonline.net/>.
- A. E. Kossovsky, *Benford's Law: Theory, the General Law of Relative Quantities, and Forensic Fraud Detection Applications*, WSPC, 2014.
- S. J. Miller (editor), *Theory and Applications of Benford's Law*, Princeton University Press, 2015.
- M. Nigrini, *Benford's Law: Applications for Forensic Accounting, Auditing, and Fraud Detection*, 1st Edition, Wiley, 2014.

Interesting Question


Motivating Question: For a nice data set, such as the Fibonacci numbers, stock prices, street addresses of college employees and students, ..., what percent of the leading digits are 1?



Natural guess: 10% (but immediately correct to 11%!).

Interesting Question


Motivating Question: For a nice data set, such as the Fibonacci numbers, stock prices, street addresses of college employees and students, ..., what percent of the leading digits are 1?




Answer: Benford's law!

Examples with First Digit Bias


Fibonacci numbers




Twitter users by # followers



Most common iPhone passcodes



Distance of stars from Earth



Summary

- Explain Benford's Law.
- Discuss examples and applications.
- Sketch proofs.
- Describe open problems.

Caveats!

- A math test indicating fraud is *not* proof of fraud: unlikely events, alternate reasons.



Examples

- recurrence relations
- special functions (such as $n!$)
- iterates of power, exponential, rational maps
- products of random variables
- L -functions, characteristic polynomials
- iterates of the $3x + 1$ map
- differences of order statistics
- hydrology and financial data
- many hierarchical Bayesian models

Applications


- Analyzing round-off errors.
- Determining the optimal way to store numbers.
- Detecting tax and image fraud, and data integrity.

General Theory

Benford's Law: Newcomb (1881), Benford (1938)

Statement

For many data sets, probability of observing a first digit of d base B is $\log_B \left(\frac{d+1}{d} \right)$; base 10 about 30% are 1's ($\log_{10}(2) \approx .3010 \dots$).



Benford's Law (probabilities)

Background Material

- Modulo: $a = b \pmod c$ if $a - b$ is an integer times c ; thus $17 = 5 \pmod{12}$, and $4.5 = .5 \pmod{1}$.
- Significand: $x = S_{10}(x) \cdot 10^k$, k integer, $1 \leq S_{10}(x) < 10$. Thus $2020.1701 = 2.0201701 \cdot 10^3$.
- $S_{10}(a) = S_{10}(b)$ if and only if a and b have the same leading digits. Note $\log_{10} a = \log_{10} S_{10}(b) + k$.
- **Key observation:** $\log_{10}(x) = \log_{10}(\tilde{x}) \pmod{1}$ if and only if x and \tilde{x} have the same leading digits.

Thus often study $y = \log_{10} x \pmod{1}$.

Advanced: $e^{2\pi i u} = e^{2\pi i(u \pmod{1})}$.

Equidistribution and Benford's Law


Equidistribution

$\{y_n\}_{n=1}^{\infty}$ is equidistributed modulo 1 if probability $y_n \bmod 1 \in [a, b]$ tends to $b - a$:

$$\frac{\#\{n \leq N : y_n \bmod 1 \in [a, b]\}}{N} \rightarrow b - a.$$


- Thm: $\beta \notin \mathbb{Q}$, $n\beta$ is equidistributed mod 1.
- Examples: $\log_{10} 2$, $\log_{10} \left(\frac{1+\sqrt{5}}{2} \right) \notin \mathbb{Q}$.

Example of Equidistribution: $n\sqrt{\pi} \bmod 1$




$n\sqrt{\pi} \bmod 1$ for $n \leq 10$

Example of Equidistribution: $n\sqrt{\pi} \bmod 1$




$n\sqrt{\pi} \bmod 1$ for $n \leq 100$

Example of Equidistribution: $n\sqrt{\pi} \bmod 1$



$n\sqrt{\pi} \bmod 1$ for $n \leq 1000$

Example of Equidistribution: $n\sqrt{\pi} \bmod 1$



$n\sqrt{\pi} \bmod 1$ for $n \leq 10,000$

Logarithms and Benford's Law

Fundamental Equivalence

Data set $\{x_i\}$ is Benford base B if $\{y_i\}$ is equidistributed mod 1, where $y_i = \log_B x_i$.


Logarithms and Benford's Law

Fundamental Equivalence

Data set $\{x_i\}$ is Benford base B if $\{y_i\}$ is equidistributed mod 1, where $y_i = \log_B x_i$.

$$x = S_{10}(x) \cdot 10^k \text{ then}$$


$$\log_{10} x = \log_{10} S_{10}(x) + k = \log_{10} S_{10}x \text{ mod } 1.$$




Logarithms and Benford's Law

$$\begin{aligned} \text{Prob}(\text{leading digit } d) &= \log_{10}(d+1) - \log_{10}(d) \\ &= \log_{10}\left(\frac{d+1}{d}\right) \\ &= \log_{10}\left(1 + \frac{1}{d}\right). \end{aligned}$$

Have Benford's law \leftrightarrow
mantissa of logarithms
of data are uniformly
distributed



The Power of the Right Perspective



The Power of the Right Perspective



Examples

- 2^n is Benford base 10 as $\log_{10} 2 \notin \mathbb{Q}$.

Examples

- **Fibonacci numbers are Benford base 10.**

$$F_{n+1} = F_n + F_{n-1}.$$

Guess $F_n = r^n$: $r^{n+1} = r^n + r^{n-1}$ or $r^2 = r + 1$.

Roots $r = (1 \pm \sqrt{5})/2$.

General solution: $F_n = c_1 r_1^n + c_2 r_2^n$.

$$\text{Binet: } F_n = \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n.$$

- **Most linear recurrence relations Benford:**

Examples

- **Fibonacci numbers are Benford base 10.**

$$F_{n+1} = F_n + F_{n-1}.$$

Guess $F_n = r^n$: $r^{n+1} = r^n + r^{n-1}$ or $r^2 = r + 1$.

Roots $r = (1 \pm \sqrt{5})/2$.

General solution: $F_n = c_1 r_1^n + c_2 r_2^n$.

$$\text{Binet: } F_n = \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n.$$

- **Most linear recurrence relations Benford:**

$$\diamond a_{n+1} = 2a_n$$

Examples

- **Fibonacci numbers are Benford base 10.**

$$F_{n+1} = F_n + F_{n-1}.$$

Guess $F_n = r^n$: $r^{n+1} = r^n + r^{n-1}$ or $r^2 = r + 1$.

Roots $r = (1 \pm \sqrt{5})/2$.

General solution: $F_n = c_1 r_1^n + c_2 r_2^n$.

Binet: $F_n = \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n$.

- **Most linear recurrence relations Benford:**

$$\diamond a_{n+1} = 2a_n - a_{n-1}$$

Examples

- **Fibonacci numbers are Benford base 10.**

$$F_{n+1} = F_n + F_{n-1}.$$

Guess $F_n = r^n$: $r^{n+1} = r^n + r^{n-1}$ or $r^2 = r + 1$.

Roots $r = (1 \pm \sqrt{5})/2$.

General solution: $F_n = c_1 r_1^n + c_2 r_2^n$.

$$\text{Binet: } F_n = \frac{1}{\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n.$$

- **Most linear recurrence relations Benford:**

◇ $a_{n+1} = 2a_n - a_{n-1}$

◇ take $a_0 = a_1 = 1$ or $a_0 = 0, a_1 = 1$.

Digits of 2^n

First 60 values of 2^n (only displaying 30)

			digit	#	Obs Prob	Benf Prob
1	1024	1048576				
2	2048	2097152	1	18	.300	.301
4	4096	4194304	2	12	.200	.176
8	8192	8388608	3	6	.100	.125
16	16384	16777216	4	6	.100	.097
32	32768	33554432	5	6	.100	.079
64	65536	67108864	6	4	.067	.067
128	131072	134217728	7	2	.033	.058
256	262144	268435456	8	5	.083	.051
512	524288	536870912	9	1	.017	.046

Digits of 2^n

First 60 values of 2^n (only displaying 30)

			digit	#	Obs Prob	Benf Prob
1	1024	1048576				
2	2048	2097152	1	18	.300	.301
4	4096	4194304	2	12	.200	.176
8	8192	8388608	3	6	.100	.125
16	16384	16777216	4	6	.100	.097
32	32768	33554432	5	6	.100	.079
64	65536	67108864	6	4	.067	.067
128	131072	134217728	7	2	.033	.058
256	262144	268435456	8	5	.083	.051
512	524288	536870912	9	1	.017	.046

Digits of 2^n

First 60 values of 2^n (only displaying 30): $2^{10} = 1024 \approx 10^3$.

			digit	#	Obs Prob	Benf Prob
1	1024	1048576				
2	2048	2097152	1	18	.300	.301
4	4096	4194304	2	12	.200	.176
8	8192	8388608	3	6	.100	.125
16	16384	16777216	4	6	.100	.097
32	32768	33554432	5	6	.100	.079
64	65536	67108864	6	4	.067	.067
128	131072	134217728	7	2	.033	.058
256	262144	268435456	8	5	.083	.051
512	524288	536870912	9	1	.017	.046

Digits of 2^n

First 1000 values of 2^n (only displaying 30): $2^{10} = 1024 \approx 10^3$.

			digit	#	Obs Prob	Benf Prob
1	1024	1048576				
2	2048	2097152	1	301	0.301	.301
4	4096	4194304	2	176	0.176	.176
8	8192	8388608	3	125	0.125	.125
16	16384	16777216	4	97	0.097	.097
32	32768	33554432	5	79	0.079	.079
64	65536	67108864	6	69	0.069	.067
128	131072	134217728	7	56	0.056	.058
256	262144	268435456	8	52	0.052	.051
512	524288	536870912	9	45	0.045	.046


Logarithms and Benford's Law

χ^2 values for α^n , $1 \leq n \leq N$ (5% 15.5).

N	$\chi^2(\gamma)$	$\chi^2(e)$	$\chi^2(\pi)$
100	0.72	0.30	46.65
200	0.24	0.30	8.58
400	0.14	0.10	10.55
500	0.08	0.07	2.69
700	0.19	0.04	0.05
800	0.04	0.03	6.19
900	0.09	0.09	1.71
1000	0.02	0.06	2.90


Logarithms and Benford's Law: Base 10 (5%: $\log(\chi^2) \approx 2.74$)

$\log(\chi^2)$ vs N for π^n (red) and e^n (blue),
 $n \in \{1, \dots, N\}$.



Logarithms and Benford's Law: Base 10 (5%: $\log(\chi^2) \approx 2.74$)

$\log(\chi^2)$ vs N for π^n (red) and e^n (blue),
 $n \in \{1, \dots, N\}$. Note $\pi^{175} \approx 1.0028 \cdot 10^{87}$.



Applications


Detecting Fraud

Bank Fraud

- Audit of a bank revealed huge spike of numbers starting with 48 and 49, most due to one person.
- Write-off limit of \$5,000. Officer had friends applying for credit cards, ran up balances just under \$5,000 then he would write the debts off.

Benford's Law and Election Data

Philadelphia Precincts Biden 2020 Base 10



Data for Philadelphia Precincts Biden 2020 Base 10


Chisquare first digit: 27.1094. Chisquare second digit: 6.81388.

Chisquare thresholds (8 degrees of freedom): 15.5 (95%), 20.1 (99%)

Chisquare thresholds (9 degrees of freedom): 16.9 (95%), 21.7 (99%)

First digit test fails Benford at the 99% level.

Second digit test passes at the 95% confidence level.



Data for Philadelphia Precincts Biden 2020 Base 3

Chisquare first digit: 0.120108. Chisquare second digit: 1.57438.

Chisquare thresholds (1 degrees of freedom): 3.84 (95%), 6.63 (99%)

Chisquare thresholds (2 degrees of freedom): 5.99 (95%), 9.21 (99%)

First digit test passes at the 95% confidence level.

Second digit test passes at the 95% confidence level.

Can you see the cat in the tree?



Transmitting Images

How to transmit an image?

- Have an $L \times W$ grid with LW pixels.
- Each pixel a triple: (Red, Green, Blue).
- Often each value in $\{0, 1, 2, 3, \dots, 2^n - 1\}$.
- $n = 8$ gives 256 choices for each, or 16,777,216 possibilities.

Steganography

Steganography: Concealing a message in another message: <https://en.wikipedia.org/wiki/Steganography>.

Steganography

Steganography: Concealing a message in another message: <https://en.wikipedia.org/wiki/Steganography>.

Take one of the colors, say **red**, a number from 0 to 255.

Write in binary: $r_7 2^7 + r_6 2^6 + \dots + r_1 2 + r_0$.

If change just the last or last two digits, very minor change to image.

Can you see the cat in the tree?



Can you see the cat in the tree?




Why Benford's Law?

Streets

Not all data sets satisfy Benford's Law.

- Long street $[1, L]$: $L = 199$ versus $L = 999$.
- Oscillates b/w $1/9$ and $5/9$ with first digit 1.




Probability first digit 1 versus street length L .

Streets

Not all data sets satisfy Benford's Law.

- Long street $[1, L]$: $L = 199$ versus $L = 999$.
- Oscillates b/w $1/9$ and $5/9$ with first digit 1.




Probability first digit 1 versus $\log(\text{street length } L)$.

Streets

Not all data sets satisfy Benford's Law.

- Long street $[1, L]$: $L = 199$ versus $L = 999$.
- Oscillates b/w $1/9$ and $5/9$ with first digit 1.




Probability first digit 1 versus $\log(\text{street length } L)$.

What if we have many streets of different lengths?

Amalgamating Streets



All houses: 1000 Streets,
each from 1 to 10000.



First digit and first two digits vs Benford.

Amalgamating Streets



All houses: 1000 Streets,
each from 1 to rand(10000).



First digit and first two digits vs Benford.

Amalgamating Streets

All houses: 1000 Streets,
each 1 to $\text{rand}(\text{rand}(10000))$.





First digit and first two digits vs Benford.

Conclusion: More processes, closer to Benford.

Amalgamating Streets

All houses: 1000 Streets,
each 1 to $\text{rand}(\text{rand}(\text{rand}(10000)))$.




First digit and first two digits vs Benford.

Conclusion: More processes, closer to Benford.

Central Limit Theorem

Normal $N(\mu, \sigma^2)$: $p(x) = e^{-(x-\mu)^2/2\sigma^2} / \sqrt{2\pi\sigma^2}$.



Theorem


If X_1, X_2, \dots independent, identically distributed random variables (mean μ , variance σ^2 , finite moments) then

$$S_N := \frac{X_1 + \dots + X_N - N\mu}{\sigma\sqrt{N}} \text{ converges to } N(0, 1).$$

Central Limit Theorem: Sums of Uniform Random Variables

$X_i \sim \text{Unif}(-1/2, 1/2)$ (adjusted to mean 0, variance 1)


$Y_1 = X_1/\sigma_{X_1}$ vs $N(0, 1)$.



Central Limit Theorem: Sums of Uniform Random Variables

$X_i \sim \text{Unif}(-1/2, 1/2)$ (adjusted to mean 0, variance 1)


$Y_2 = (X_1 + X_2)/\sigma_{X_1+X_2}$ vs $N(0, 1)$.



Central Limit Theorem: Sums of Uniform Random Variables

$X_i \sim \text{Unif}(-1/2, 1/2)$ (adjusted to mean 0, variance 1)


$$Y_4 = (X_1 + X_2 + X_3 + X_4) / \sigma_{X_1+X_2+X_3+X_4} \text{ vs } N(0, 1).$$



Central Limit Theorem: Sums of Uniform Random Variables

$X_i \sim \text{Unif}(-1/2, 1/2)$ (adjusted to mean 0, variance 1)

$$Y_8 = (X_1 + \cdots + X_8) / \sigma_{X_1 + \cdots + X_8} \text{ vs } N(0, 1).$$



Central Limit Theorem: Sums of Uniform Random Variables

$X_i \sim \text{Unif}(-1/2, 1/2)$ (adjusted to mean 0, variance 1)

Density of $Y_4 = (X_1 + \dots + X_4)/\sigma_{X_1+\dots+X_4}$.


$$\left\{ \begin{array}{ll} \frac{1}{27} (18 + 9\sqrt{3}y - \sqrt{3}y^3) & y = 0 \\ \frac{1}{18} (12 - 6y^2 - \sqrt{3}y^3) & -\sqrt{3} < y < 0 \\ \frac{1}{54} (72 - 36\sqrt{3}y + 18y^2 - \sqrt{3}y^3) & \sqrt{3} < y < 2\sqrt{3} \\ \frac{1}{54} (18\sqrt{3}y - 18y^2 + \sqrt{3}y^3) & y = \sqrt{3} \\ \frac{1}{18} (12 - 6y^2 + \sqrt{3}y^3) & 0 < y < \sqrt{3} \\ \frac{1}{54} (72 + 36\sqrt{3}y + 18y^2 + \sqrt{3}y^3) & -2\sqrt{3} < y \leq -\sqrt{3} \\ 0 & \text{True} \end{array} \right.$$

$$\sqrt{3}$$

(Don't even think of asking to see Y_8 's!)

Normal Distributions Mod 1


As $\sigma \rightarrow \infty$, $N(0, \sigma^2) \bmod 1 \rightarrow \text{Unif}(0, 1)$.



Variance is .01.

Normal Distributions Mod 1


As $\sigma \rightarrow \infty$, $N(0, \sigma^2) \bmod 1 \rightarrow \text{Unif}(0, 1)$.



Variance is .1.

Normal Distributions Mod 1

As $\sigma \rightarrow \infty$, $N(0, \sigma^2) \bmod 1 \rightarrow \text{Unif}(0, 1)$.



Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

X_1, X_2, \dots nice, $W_N = X_1 \cdot X_2 \cdots X_N$.

Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

X_1, X_2, \dots nice, $W_N = X_1 \cdot X_2 \cdots X_N$.

$Y_i = \log_{10} X_i$, $V_N := \log_{10} W_N$.

Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

X_1, X_2, \dots nice, $W_N = X_1 \cdot X_2 \cdots X_N$.

$Y_i = \log_{10} X_i$, $V_N := \log_{10} W_N$.

$$V_N = \log_{10}(X_1 \cdot X_2 \cdots X_N)$$

Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

X_1, X_2, \dots nice, $W_N = X_1 \cdot X_2 \cdots X_N$.

$Y_i = \log_{10} X_i$, $V_N := \log_{10} W_N$.

$$\begin{aligned} V_N &= \log_{10}(X_1 \cdot X_2 \cdots X_N) \\ &= \log_{10} X_1 + \log_{10} X_2 + \cdots + \log_{10} X_N \end{aligned}$$

Products and Benford's Law

Pavlovian Response: See a product, take a logarithm.

X_1, X_2, \dots nice, $W_N = X_1 \cdot X_2 \cdots X_N$.

$Y_i = \log_{10} X_i$, $V_N := \log_{10} W_N$.

$$\begin{aligned} V_N &= \log_{10}(X_1 \cdot X_2 \cdots X_N) \\ &= \log_{10} X_1 + \log_{10} X_2 + \cdots + \log_{10} X_N \\ &= Y_1 + Y_2 + \cdots + Y_N. \end{aligned}$$

Need distribution of $V_N \bmod 1$, which by CLT becomes uniform, implying Benfordness!

Stick Decomposition

- T. Becker, D. Burt, T. C. Corcoran, A. Greaves-Tunnell, J. R. Iafrate, J. Jing, S. J. Miller, J. D. Porfilio, R. Ronan, J. Samranvedhya, F. W. Strauch and B. Talbut, *Benford's Law and Continuous Dependent Random Variables*, *Annals of Physics* **388** (2018), 350–381.
- J. Iafrate, S. J. Miller and F. W. Strauch, *Equipartitions and a distribution for numbers: A statistical model for Benford's law*, *Physical Review E* **91** (2015), no. 6, 062138 (6 pages).
- *Benford Behavior in Stick Fragmentation Problems*, Bruce Fang, Ava Irons, Ella Lippelman, and Steven J. Miller, *Stats* (Special Issue on Benford's Law(s) and Applications, second edition).
<https://doi.org/10.3390/stats8040091>.
- *Behavior Resulting From Stick and Box Fragmentation Processes* (Bruce Fang, Steven J. Miller), preprint (2025).
<https://arxiv.org/abs/2508.12915>.

Fixed Proportion Decomposition Process

Decomposition Process

- 1 Consider a stick of length \mathcal{L} .

Fixed Proportion Decomposition Process

Decomposition Process

- 1 Consider a stick of length \mathcal{L} .
- 2 Uniformly choose a proportion $p \in (0, 1)$.

Fixed Proportion Decomposition Process

Decomposition Process


- 1 Consider a stick of length \mathcal{L} .
- 2 Uniformly choose a proportion $p \in (0, 1)$.
- 3 Break the stick into two pieces—lengths $p\mathcal{L}$ and $(1 - p)\mathcal{L}$.

Fixed Proportion Decomposition Process

Decomposition Process


- 1 Consider a stick of length \mathcal{L} .
- 2 Uniformly choose a proportion $p \in (0, 1)$.
- 3 Break the stick into two pieces—lengths $p\mathcal{L}$ and $(1 - p)\mathcal{L}$.
- 4 Repeat N times (using the same proportion).

Fixed Proportion Decomposition Process




Fixed Proportion Conjecture (Joy Jing '13)

Conjecture: The above decomposition process is Benford as $N \rightarrow \infty$ for any $p \in (0, 1)$, $p \neq \frac{1}{2}$.




(B) $p = 0.51$ and $N = 10000$.




(B) $p = 0.99$ and $N = 50000$. Benford distribution overlaid.

Fixed Proportion Conjecture (Joy Jing '13)

Conjecture: The above decomposition process is Benford as $N \rightarrow \infty$ for any $p \in (0, 1)$, $p \neq \frac{1}{2}$.



(B) $p = 0.51$ and $N = 10000$.



(B) $p = 0.99$ and $N = 50000$. Benford distribution overlaid.

Counterexample (SMALL REU '13): $p = \frac{1}{11}$, $1 - p = \frac{10}{11}$.

Benford Analysis

At N^{th} level,

- 2^N sticks
- $N + 1$ distinct lengths: write $p^{N-j}(1 - p)^j$ as

$p^N \left(\frac{1 - p}{p} \right)^j$, $j \in \{0, \dots, N\}$, have $\binom{N}{j}$ times.

Benford Analysis

At N^{th} level,

- 2^N sticks
- $N + 1$ distinct lengths: write $p^{N-j}(1 - p)^j$ as

$$p^N \left(\frac{1-p}{p} \right)^j, \quad j \in \{0, \dots, N\}, \quad \text{have } \binom{N}{j} \text{ times.}$$

(Weighted) Geometric with ratio $\frac{1-p}{p} = 10^y$;
behavior depends on irrationality of y !

Benford Analysis

At N^{th} level,

- 2^N sticks
- $N + 1$ distinct lengths: write $p^{N-j}(1 - p)^j$ as

$$p^N \left(\frac{1 - p}{p} \right)^j, \quad j \in \{0, \dots, N\}, \text{ have } \binom{N}{j} \text{ times.}$$

(Weighted) Geometric with ratio $\frac{1-p}{p} = 10^y$;
behavior depends on irrationality of y !

Theorem: Benford if and only if y irrational.

Benford Analysis (cont)


Say $\frac{1-p}{p} = 10^{r/q}$ for r, q integers.

All terms with index $j \bmod q$ have same leading digit; probability index $j \bmod q$ is

$$\begin{aligned} \frac{1}{2^N} \left[\binom{N}{j} + \binom{N}{j+q} + \binom{N}{j+2q} + \dots \right] &= \frac{1}{q} \sum_{s=0}^{q-1} \left(\cos \frac{\pi s}{q} \right)^N \cos \frac{\pi(N-2j)s}{q} \\ &= \frac{1}{q} \left(1 + \sum_{s=1}^{q-1} \left(\cos \frac{\pi s}{q} \right)^N \cos \frac{\pi(N-2j)s}{q} \right) \\ &= \frac{1}{q} \left(1 + \text{Err} \left[(q-1) \left(\cos \frac{\pi}{q} \right)^N \right] \right), \end{aligned}$$



where $\text{Err}[X]$ indicates an absolute error of size at most X

Examples




$p = 3/11$, 1000 levels; $y = \log_{10}(8/3) \notin \mathbb{Q}$
(irrational)

Examples



$p = 1/11$, 1000 levels; $y = 1 \in \mathbb{Q}$
(rational)

Examples



$p = 1/(1 + 10^{33/10})$, 1000 levels; $y = 33/10 \in \mathbb{Q}$
(rational)

Random Cuts




Figure: Unrestricted Decomposition: Breaking L into pieces, $N = 3$.

Fixed Proportion Decomposition Process

- Consider a stick of length L .
- Proportions p_1, p_2, \dots, p_{m-1} in $(0, 1)$.
- Cut at p_1, p_2, \dots, p_{m-1} , create m pieces.
- Do N iterations, which results in
 - Te $\diamond m^N$ sticks
 - $\diamond \binom{m+N-1}{N}$ distinct lengths for generic p_i .

Theorem (Irons-Lippelman-Miller)

Consider the above described decomposition process. Choose y_1, y_2, \dots, y_{m-1} such that $10^{y_1} = p_1/p_2$, $10^{y_2} = p_2/p_3$, \dots , $10^{y_{m-1}} = p_{m-1}/p_m$, which are the ratios of adjacent lengths. The decomposition process results in a distribution of stick lengths that do not follow Benford's Law if all

$$y_1, y_2, \dots, y_{m-1} \in \mathbb{Q}.$$

Theorem (Fang-Miller)

Choose y_1, y_2, \dots, y_{m-1} such that $10^{y_1} = p_1/p_2$, $10^{y_2} = p_2/p_3, \dots, 10^{y_{m-1}} = p_{m-1}/p_m$, which is the ratio of adjacent lengths. If at least one of y_1, y_2, \dots, y_{m-1} is irrational, then the distribution of stick lengths follow Benford's Law.

Benford Good Processes

- A. Kontorovich and S. J. Miller, *Benford's Law, values of L-functions and the $3x + 1$ problem*, Acta Arithmetica **120** (2005), no. 3, 269–297.

Poisson Summation and Benford's Law: Definitions

- Feller, Pinkham (often exact processes)
- data $Y_{T,B} = \log_B \vec{X}_T$ (discrete/continuous):

$$\mathbb{P}(A) = \lim_{T \rightarrow \infty} \frac{\#\{n \in A : n \leq T\}}{T}$$

- Poisson Summation Formula: f nice:

$$\sum_{l=-\infty}^{\infty} f(l) = \sum_{l=-\infty}^{\infty} \hat{f}(l),$$

$$\text{Fourier transform } \hat{f}(\xi) = \int_{-\infty}^{\infty} f(x) e^{-2\pi i x \xi} dx.$$

Benford Good Process

X_T is **Benford Good** if there is a nice f st

$$\text{CDF}_{\vec{Y}_{T,B}}(y) = \int_{-\infty}^y \frac{1}{T} f\left(\frac{t}{T}\right) dt + E_T(y) := G_T(y)$$

and monotonically increasing h ($h(|T|) \rightarrow \infty$):

- **Small tails:** $G_T(\infty) - G_T(Th(T)) = o(1)$,
 $G_T(-Th(T)) - G_T(-\infty) = o(1)$.

- **Decay of the Fourier Transform:**

$$\sum_{\ell \neq 0} \left| \frac{\widehat{f}(T\ell)}{\ell} \right| = o(1).$$

- **Small translated error:** $\mathcal{E}(a, b, T) =$

$$\sum_{|\ell| \leq Th(T)} [E_T(b + \ell) - E_T(a + \ell)] = o(1).$$

Main Theorem

Theorem (Kontorovich and M–, 2005)

X_T converging to X as $T \rightarrow \infty$ (think spreading Gaussian). If X_T is Benford good, then X is Benford.

- **Examples**

- ◇ L -functions
- ◇ characteristic polynomials (RMT)
- ◇ $3x + 1$ problem
- ◇ geometric Brownian motion.

Sketch of the proof

- **Structure Theorem:**
 - ◇ main term is something nice spreading out
 - ◇ apply Poisson summation
- **Control translated errors:**
 - ◇ hardest step
 - ◇ techniques problem specific

Sketch of the proof (continued)

$$\sum_{l=-\infty}^{\infty} \mathbb{P} \left(a + l \leq \vec{Y}_{T,B} \leq b + l \right)$$

$$= \sum_{|l| \leq Th(T)} [G_T(b + l) - G_T(a + l)] + o(1)$$

Sketch of the proof (continued)


$$\begin{aligned}
 & \sum_{l=-\infty}^{\infty} \mathbb{P} \left(a + l \leq \vec{Y}_{T,B} \leq b + l \right) \\
 = & \sum_{|\ell| \leq Th(T)} [G_T(b + \ell) - G_T(a + \ell)] + o(1) \\
 = & \int_a^b \sum_{|\ell| \leq Th(T)} \frac{1}{T} f \left(\frac{t + \ell}{T} \right) dt + \mathcal{E}(a, b, T) + o(1)
 \end{aligned}$$

Sketch of the proof (continued)

$$\begin{aligned}
 & \sum_{l=-\infty}^{\infty} \mathbb{P} \left(a + l \leq \vec{Y}_{T,B} \leq b + l \right) \\
 = & \sum_{|l| \leq Th(T)} [G_T(b + l) - G_T(a + l)] + o(1) \\
 = & \int_a^b \sum_{|l| \leq Th(T)} \frac{1}{T} f \left(\frac{t + l}{T} \right) dt + \mathcal{E}(a, b, T) + o(1) \\
 = & \hat{f}(0) \cdot (b - a) + \sum_{l \neq 0} \hat{f}(Tl) \frac{e^{2\pi i b l} - e^{2\pi i a l}}{2\pi i l} + o(1).
 \end{aligned}$$

Riemann Zeta Function

$$\left| \zeta \left(\frac{1}{2} + i \frac{k}{4} \right) \right|, k \in \{0, 1, \dots, 65535\}.$$



The $3x + 1$ Problem
and
Benford's Law

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- 7

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11$

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17$

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17 \rightarrow_2 13$

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17 \rightarrow_2 13 \rightarrow_3 5$

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17 \rightarrow_2 13 \rightarrow_3 5 \rightarrow_4 1$

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17 \rightarrow_2 13 \rightarrow_3 5 \rightarrow_4 1 \rightarrow_2 1$,

3x + 1 Problem

- Kakutani (conspiracy), Erdős (not ready).
- x odd, $T(x) = \frac{3x+1}{2^k}$, $2^k \parallel 3x + 1$.
- Conjecture: for some $n = n(x)$, $T^n(x) = 1$.
- $7 \rightarrow_1 11 \rightarrow_1 17 \rightarrow_2 13 \rightarrow_3 5 \rightarrow_4 1 \rightarrow_2 1$,
2-path (1, 1), 5-path (1, 1, 2, 3, 4).
 m -path: (k_1, \dots, k_m) .

Heuristic Proof of $3x + 1$ Conjecture

$$\begin{aligned}
 a_{n+1} &= T(a_n) \\
 \mathbb{E}[\log a_{n+1}] &\approx \sum_{k=1}^{\infty} \frac{1}{2^k} \log \left(\frac{3a_n}{2^k} \right) \\
 &= \log a_n + \log 3 - \log 2 \sum_{k=1}^{\infty} \frac{k}{2^k} \\
 &= \log a_n + \log \left(\frac{3}{4} \right).
 \end{aligned}$$

Geometric Brownian Motion, drift $\log(3/4) < 1$.

3x + 1 and Benford

Theorem (Kontorovich and M–, 2005)

As $m \rightarrow \infty$, $x_m / (3/4)^m x_0$ is Benford.

Theorem (Lagarias-Soundararajan, 2006)

$X \geq 2^N$, for all but at most $c(B)N^{-1/36} X$ initial seeds the distribution of the first N iterates of the $3x + 1$ map are within $2N^{-1/36}$ of the Benford probabilities.

Structure Theorem: Sinai, Kontorovich-Sinai

$$\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{\#\{n \leq N: n \equiv 1, 5 \pmod{6}, n \in A\}}{\#\{n \leq N: n \equiv 1, 5 \pmod{6}\}}.$$

(k_1, \dots, k_m) : two full arithm progressions:

$$6 \cdot 2^{k_1 + \dots + k_m} p + q.$$

Theorem (Sinai, Kontorovich-Sinai)

k_i -values are i.i.d.r.v. (geometric, 1/2):

Structure Theorem: Sinai, Kontorovich-Sinai

$$\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{\#\{n \leq N : n \equiv 1, 5 \pmod{6}, n \in A\}}{\#\{n \leq N : n \equiv 1, 5 \pmod{6}\}}.$$

(k_1, \dots, k_m) : two full arithm progressions:

$$6 \cdot 2^{k_1 + \dots + k_m} p + q.$$

Theorem (Sinai, Kontorovich-Sinai)

k_i -values are i.i.d.r.v. (geometric, 1/2):

$$\mathbb{P} \left(\frac{\log_2 \left[\frac{x_m}{\left(\frac{3}{4}\right)^m x_0} \right]}{\sqrt{2m}} \leq a \right) = \mathbb{P} \left(\frac{S_m - 2m}{\sqrt{2m}} \leq a \right)$$

Structure Theorem: Sinai, Kontorovich-Sinai

$$\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{\#\{n \leq N : n \equiv 1, 5 \pmod{6}, n \in A\}}{\#\{n \leq N : n \equiv 1, 5 \pmod{6}\}}.$$

(k_1, \dots, k_m) : two full arithm progressions:
 $6 \cdot 2^{k_1 + \dots + k_m} p + q.$

Theorem (Sinai, Kontorovich-Sinai)

k_i -values are i.i.d.r.v. (geometric, $1/2$):

$$\mathbb{P} \left(\frac{\log_2 \left[\frac{x_m}{\left(\frac{3}{4}\right)^m x_0} \right]}{(\log_2 B) \sqrt{2m}} \leq a \right) = \mathbb{P} \left(\frac{S_m - 2m}{(\log_2 B) \sqrt{2m}} \leq a \right)$$

Structure Theorem: Sinai, Kontorovich-Sinai

$$\mathbb{P}(A) = \lim_{N \rightarrow \infty} \frac{\#\{n \leq N : n \equiv 1, 5 \pmod{6}, n \in A\}}{\#\{n \leq N : n \equiv 1, 5 \pmod{6}\}}.$$

(k_1, \dots, k_m) : two full arithm progressions:

$$6 \cdot 2^{k_1 + \dots + k_m} p + q.$$

Theorem (Sinai, Kontorovich-Sinai)

k_i -values are i.i.d.r.v. (geometric, $1/2$):

$$\mathbb{P} \left(\frac{\log_B \left[\frac{x_m}{\left(\frac{3}{4}\right)^m x_0} \right]}{\sqrt{2m}} \leq a \right) = \mathbb{P} \left(\frac{\frac{(S_m - 2m)}{\log_2 B}}{\sqrt{2m}} \leq a \right)$$

Sketch of the proof of Benfordness

- Failed Proof: lattices, bad errors.

- CLT: $(S_m - 2m)/\sqrt{2m} \rightarrow N(0, 1)$:

$$\mathbb{P}(S_m - 2m = k) = \frac{\eta(k/\sqrt{m})}{\sqrt{m}} + O\left(\frac{1}{g(m)\sqrt{m}}\right).$$

- Quantified Equidistribution:

$$I_\ell = \{\ell M, \dots, (\ell + 1)M - 1\}, \quad M = m^c, \quad c < 1/2$$

$$k_1, k_2 \in I_\ell: \left| \eta\left(\frac{k_1}{\sqrt{m}}\right) - \eta\left(\frac{k_2}{\sqrt{m}}\right) \right| \text{ small}$$

$$C = \log_B 2 \text{ of irrationality type } \kappa < \infty:$$

$$\#\{k \in I_\ell : \overline{kC} \in [a, b]\} = M(b-a) + O(M^{1+\epsilon-1/\kappa}).$$

Irrationality Type

Irrationality type

α has irrationality type κ if κ is the supremum of all γ with

$$\liminf_{q \rightarrow \infty} q^{\gamma+1} \min_p \left| \alpha - \frac{p}{q} \right| = 0.$$

- Algebraic irrationals: type 1 (Roth's Thm).
- Theory of Linear Forms: $\log_B 2$ of finite type.

Linear Forms

Theorem (Baker)

$\alpha_1, \dots, \alpha_n$ algebraic numbers height $A_j \geq 4$,
 $\beta_1, \dots, \beta_n \in \mathbb{Q}$ with height at most $B \geq 4$,

$$\Lambda = \beta_1 \log \alpha_1 + \dots + \beta_n \log \alpha_n.$$

If $\Lambda \neq 0$ then $|\Lambda| > B^{-C\Omega \log \Omega'}$, with
 $d = [\mathbb{Q}(\alpha_i, \beta_j) : \mathbb{Q}]$, $C = (16nd)^{200n}$,
 $\Omega = \prod_j \log A_j$, $\Omega' = \Omega / \log A_n$.

Gives $\log_{10} 2$ of finite type, with $\kappa < 1.2 \cdot 10^{602}$:

$$|\log_{10} 2 - p/q| = |q \log 2 - p \log 10| / q \log 10.$$

Quantified Equidistribution

Theorem (Erdős-Turan)

$$D_N = \frac{\sup_{[a,b]} |N(b-a) - \#\{n \leq N : x_n \in [a,b]\}|}{N}$$

There is a C such that for all m:

$$D_N \leq C \cdot \left(\frac{1}{m} + \sum_{h=1}^m \frac{1}{h} \left| \frac{1}{N} \sum_{n=1}^N e^{2\pi i h x_n} \right| \right)$$

Proof of Erdős-Turan

Consider special case $x_n = n\alpha$, $\alpha \notin \mathbb{Q}$.

- Exponential sum $\leq \frac{1}{|\sin(\pi h\alpha)|} \leq \frac{1}{2||h\alpha||}$.
- Must control $\sum_{h=1}^m \frac{1}{h||h\alpha||}$, see irrationality type enter.
- type κ , $\sum_{h=1}^m \frac{1}{h||h\alpha||} = O(m^{\kappa-1+\epsilon})$, take $m = \lfloor N^{1/\kappa} \rfloor$.

3x + 1 Data: random 10,000 digit number, $2^k \parallel 3x + 1$

80,514 iterations ($(4/3)^n = a_0$ predicts 80,319);
 $\chi^2 = 13.5$ (5% 15.5).

Digit	Number	Observed	Benford
1	24251	0.301	0.301
2	14156	0.176	0.176
3	10227	0.127	0.125
4	7931	0.099	0.097
5	6359	0.079	0.079
6	5372	0.067	0.067
7	4476	0.056	0.058
8	4092	0.051	0.051
9	3650	0.045	0.046

3x + 1 Data: random 10,000 digit number, 2|3x + 1

241,344 iterations, $\chi^2 = 11.4$ (5% 15.5).

Digit	Number	Observed	Benford
1	72924	0.302	0.301
2	42357	0.176	0.176
3	30201	0.125	0.125
4	23507	0.097	0.097
5	18928	0.078	0.079
6	16296	0.068	0.067
7	13702	0.057	0.058
8	12356	0.051	0.051
9	11073	0.046	0.046

5x + 1 Data: random 10,000 digit number, $2^k \parallel 5x + 1$

27,004 iterations, $\chi^2 = 1.8$ (5% 15.5).

Digit	Number	Observed	Benford
1	8154	0.302	0.301
2	4770	0.177	0.176
3	3405	0.126	0.125
4	2634	0.098	0.097
5	2105	0.078	0.079
6	1787	0.066	0.067
7	1568	0.058	0.058
8	1357	0.050	0.051
9	1224	0.045	0.046

5x + 1 Data: random 10,000 digit number, 2|5x + 1


241,344 iterations, $\chi^2 = 3 \cdot 10^{-4}$ (5% 15.5).

Digit	Number	Observed	Benford
1	72652	0.301	0.301
2	42499	0.176	0.176
3	30153	0.125	0.125
4	23388	0.097	0.097
5	19110	0.079	0.079
6	16159	0.067	0.067
7	13995	0.058	0.058
8	12345	0.051	0.051
9	11043	0.046	0.046

The Riemann Zeta Function $\zeta(s)$
and
Benford's Law

The Riemann Zeta Function and Benford's Law

$$\left| \zeta \left(\frac{1}{2} + i \frac{k}{4} \right) \right|, k \in \{0, 1, \dots, 65535\}.$$



First digits of $\left| \zeta \left(\frac{1}{2} + i \frac{k}{4} \right) \right|$ versus Benford's law.

Proof Sketch: 'Good' L -Functions

We say an L -function is *good* if:

- Euler product:

$$L(s, f) = \sum_{n=1}^{\infty} \frac{a_f(n)}{n^s} = \prod_p \prod_{j=1}^d (1 - \alpha_{f,j}(p)p^{-s})^{-1}.$$

- $L(s, f)$ has a meromorphic continuation to \mathbb{C} , is of finite order, and has at most finitely many poles (all on the line $\operatorname{Re}(s) = 1$).
- Functional equation:

$$e^{i\omega} G(s)L(s, f) = e^{-i\omega} \overline{G(1 - \bar{s})L(1 - \bar{s})},$$

where $\omega \in \mathbb{R}$ and

$$G(s) = Q^s \prod_{i=1}^h \Gamma(\lambda_i s + \mu_i)$$

with $Q, \lambda_i > 0$ and $\operatorname{Re}(\mu_i) \geq 0$.

Proof Sketch: 'Good' L-Functions (cont)

- For some $\varkappa > 0$, $c \in \mathbb{C}$, $x \geq 2$ we have

$$\sum_{p \leq x} \frac{|a_f(p)|^2}{p} = \varkappa \log \log x + c + O\left(\frac{1}{\log x}\right).$$

- The $\alpha_{f,j}(p)$ are (Ramanujan-Petersson) tempered: $|\alpha_{f,j}(p)| \leq 1$.
- If $N(\sigma, T)$ is the number of zeros ρ of $L(s)$ with $\operatorname{Re}(\rho) \geq \sigma$ and $\operatorname{Im}(\rho) \in [0, T]$, then for some $\beta > 0$ we have

$$N(\sigma, T) = O\left(T^{1-\beta\left(\sigma-\frac{1}{2}\right)} \log T\right).$$

Known in some cases, such as $\zeta(s)$ and Hecke cuspidal forms of full level and even weight $k > 0$.

Log-Normal Law (Hejhal, Laurinćikas, Selberg)

Log-Normal Law

$$\frac{\mu(\{t \in [T, 2T] : \log |L(\sigma + it, f)| \in [a, b]\})}{T} =$$

$$\frac{1}{\sqrt{\psi(\sigma, T)}} \int_a^b e^{-\pi u^2 / \psi(\sigma, T)} du + \text{Error}$$

$$\psi(\sigma, T) = \aleph \log \left[\min \left(\log T, \frac{1}{\sigma - \frac{1}{2}} \right) \right] + O(1)$$

$$\frac{1}{2} \leq \sigma \leq \frac{1}{2} + \frac{1}{\log^\delta T}, \quad \delta \in (0, 1).$$

Result: Values of L -functions and Benford's Law

Theorem (Kontorovich and M–, 2005)

$L(s, f)$ a good L -function, as $T \rightarrow \infty$,
 $L(\sigma_T + it, f)$ is Benford.

Ingredients

- Approximate $\log L(\sigma_T + it, f)$ with $\sum_{n \leq x} \frac{c(n)\Lambda(n)}{\log n} \frac{1}{n^{\sigma_T + it}}$.
- study moments $\int_T^{2T} |\cdot|, k \leq \log^{1-\delta} T$.
- Montgomery-Vaughan: $\int_T^{2T} \sum a_n n^{-it} \overline{\sum b_m m^{-it}} dt = H \sum a_n \bar{b}_n + O(1) \sqrt{\sum n |a_n|^2 \sum n |b_n|^2}$.

Results: Explicit L -Function Statement

Theorem (Kontorovich-Miller '05)

Let $L(s, f)$ be a good L -function. Fix a $\delta \in (0, 1)$. For each T , let $\sigma_T = \frac{1}{2} + \frac{1}{\log^\delta T}$. Then as $T \rightarrow \infty$

$$\frac{\mu \{t \in [T, 2T] : M_B(|L(\sigma_T + it, f)|) \leq \tau\}}{T} \rightarrow \log_B \tau$$

Thus the values of the L -function satisfy Benford's Law in the limit for any base B .

Conclusions and References

Conclusions and Future Investigations

- See many different systems exhibit Benford behavior.
- Ingredients of proofs (logarithms, equidistribution).
- Applications to fraud detection / data integrity.

-  A. K. Adhikari, *Some results on the distribution of the most significant digit*, Sankhyā: The Indian Journal of Statistics, Series B **31** (1969), 413–420.
-  A. K. Adhikari and B. P. Sarkar, *Distribution of most significant digit in certain functions whose arguments are random variables*, Sankhyā: The Indian J. of Statistics, Series B **30** (1968), 47–58.
-  R. N. Bhattacharya, *Speed of convergence of the n -fold convolution of a probability measure on a compact group*, Z. Wahrscheinlichkeitstheorie verw. Geb. **25** (1972), 1–10.
-  F. Benford, *The law of anomalous numbers*, Proceedings of the American Philosophical Society **78** (1938), 551–572.
http://www.jstor.org/stable/984802?seq=1#page_scan_tab_contents.

-  A. Berger, L. A. Bunimovich and T. Hill, *One-dimensional dynamical systems and Benford's Law*, Trans. AMS **357** (2005), no. 1, 197–219. <http://www.ams.org/journals/tran/2005-357-01/S0002-9947-04-03455-5/>.
-  A. Berger and T. Hill, *Newton's method obeys Benford's law*, The Amer. Math. Monthly **114** (2007), no. 7, 588-601. http://digitalcommons.calpoly.edu/cgi/viewcontent.cgi?article=1058&context=rgp_rsr.
-  A. Berger and T. Hill, *Benford on-line bibliography*, <http://www.benfordonline.net/>.
-  J. Boyle, *An application of Fourier series to the most significant digit problem* Amer. Math. Monthly **101** (1994), 879–886. http://www.jstor.org/stable/2975136?seq=1#page_scan_tab_contents.
-  J. Brown and R. Duncan, *Modulo one uniform distribution of the sequence of logarithms of certain recursive sequences*, Fibonacci Quarterly **8** (1970) 482–486.



P. Diaconis, *The distribution of leading digits and uniform distribution mod 1*, Ann. Probab. **5** (1979), 72–81.

<http://statweb.stanford.edu/~cgates/PERSI/papers/digits.pdf>.



W. Feller, *An Introduction to Probability Theory and its Applications, Vol. II*, second edition, John Wiley & Sons, Inc., 1971.



R. W. Hamming, *On the distribution of numbers*, Bell Syst. Tech. J. **49** (1970), 1609-1625.

<https://archive.org/details/bstj49-8-1609>.



T. Hill, *The first-digit phenomenon*, American Scientist **86** (1996), 358–363. <http://www.americanscientist.org/issues/feature/1998/4/the-first-digit-phenomenon/99999>.



T. Hill, *A statistical derivation of the significant-digit law*, Statistical Science **10** (1996), 354–363.

<https://projecteuclid.org/euclid.ss/1177009869>.

- 
 P. J. Holewijn, *On the uniform distribuiton of sequences of random variables*, *Z. Wahrscheinlichkeitstheorie verw. Geb.* **14** (1969), 89–92.

- 
 W. Hurlimann, *Benford's Law from 1881 to 2006: a bibliography*, <http://arxiv.org/abs/math/0607168>.

- 
 D. Jang, J. Kang, A. Kruckman, J. Kudo & S. J. Miller, *Chains of distributions, hierarchical Bayesian models and Benford's Law*, *Journal of Algebra, Number Theory: Advances and Applications*, volume 1, number 1 (March 2009), 37–60. <http://arxiv.org/abs/0805.4226>.

- 
 E. Janvresse and T. de la Rue, *From uniform distribution to Benford's law*, *Journal of Applied Probability* **41** (2004) no. 4, 1203–1210. http://www.jstor.org/stable/4141393?seq=1#page_scan_tab_contents.

- 
 A. Kontorovich and S. J. Miller, *Benford's Law, Values of L-functions and the $3x + 1$ Problem*, *Acta Arith.* **120** (2005), 269–297. <http://arxiv.org/pdf/math/0412003.pdf>.

-  D. Knuth, *The Art of Computer Programming, Volume 2: Seminumerical Algorithms*, Addison-Wesley, third edition, 1997.
-  J. Lagarias and K. Soundararajan, *Benford's Law for the $3x + 1$ Function*, J. London Math. Soc. (2) **74** (2006), no. 2, 289–303.
<http://arxiv.org/pdf/math/0509175.pdf>.
-  S. Lang, *Undergraduate Analysis*, 2nd edition, Springer-Verlag, New York, 1997.
-  P. Levy, *L'addition des variables aléatoires définies sur une circonférence*, Bull. de la S. M. F. **67** (1939), 1–41.
-  E. Ley, *On the peculiar distribution of the U.S. Stock Indices Digits*, The American Statistician **50** (1996), no. 4, 311–313.
http://www.jstor.org/stable/2684926?seq=1#page_scan_tab_contents.

- 
 R. M. Loynes, *Some results in the probabilistic theory of asymptotic uniform distributions modulo 1*, *Z. Wahrscheinlichkeitstheorie verw. Geb.* **26** (1973), 33–41.

- 
 S. J. Miller, *Benford's Law: Theory and Applications*, Princeton University Press, 2015. http://web.williams.edu/Mathematics/sjmiller/public_html/benford/.

- 
 S. J. Miller and M. Nigrini, *The Modulo 1 Central Limit Theorem and Benford's Law for Products*, *International Journal of Algebra* **2** (2008), no. 3, 119–130.
<http://arxiv.org/pdf/math/0607686v2>.

- 
 S. J. Miller and M. Nigrini, *Order Statistics and Benford's law*, *International Journal of Mathematics and Mathematical Sciences*, Volume 2008 (2008), Article ID 382948, 19 pages.
<http://arxiv.org/pdf/math/0601344v5.pdf>.



S. J. Miller and R. Takloo-Bighash, *An Invitation to Modern Number Theory*, Princeton University Press, Princeton, NJ, 2006.
http://web.williams.edu/Mathematics/sjmiller/public_html/book/index.html.



S. Newcomb, *Note on the frequency of use of the different digits in natural numbers*, Amer. J. Math. **4** (1881), 39-40.
http://www.jstor.org/stable/2369148?seq=1#page_scan_tab_contents.



M. Nigrini, *Digital Analysis and the Reduction of Auditor Litigation Risk*. Pages 69–81 in *Proceedings of the 1996 Deloitte & Touche / University of Kansas Symposium on Auditing Problems*, ed. M. Ettredge, University of Kansas, Lawrence, KS, 1996.



M. Nigrini, *The Use of Benford's Law as an Aid in Analytical Procedures*, Auditing: A Journal of Practice & Theory, **16** (1997), no. 2, 52–67.

-  M. Nigrini and S. J. Miller, *Benford's Law applied to hydrology data – results and relevance to other geophysical data*, *Mathematical Geology* **39** (2007), no. 5, 469–490. <http://link.springer.com/article/10.1007%2Fs11004-007-9109-5?LI=true>.
-  M. Nigrini and S. J. Miller, *Data diagnostics using second order tests of Benford's Law*, *Auditing: A Journal of Practice and Theory* **28** (2009), no. 2, 305–324. http://accounting.uwaterloo.ca/uwcisa/symposiums/symposium_2007/AdvancedBenfordsLaw7.pdf.
-  R. Pinkham, *On the Distribution of First Significant Digits*, *The Annals of Mathematical Statistics* **32**, no. 4 (1961), 1223-1230.
-  R. A. Raimi, *The first digit problem*, *Amer. Math. Monthly* **83** (1976), no. 7, 521–538.



H. Robbins, *On the equidistribution of sums of independent random variables*, Proc. Amer. Math. Soc. **4** (1953), 786–799.
http://projecteuclid.org/download/pdf_1/euclid.aoms/1177704862.



H. Sakamoto, *On the distributions of the product and the quotient of the independent and uniformly distributed random variables*, Tôhoku Math. J. **49** (1943), 243–260.



P. Schatte, *On sums modulo 2π of independent random variables*, Math. Nachr. **110** (1983), 243–261.



P. Schatte, *On the asymptotic uniform distribution of sums reduced mod 1*, Math. Nachr. **115** (1984), 275–281.

-  P. Schatte, *On the asymptotic logarithmic distribution of the floating-point mantissas of sums*, Math. Nachr. **127** (1986), 7–20.
-  E. Stein and R. Shakarchi, *Fourier Analysis: An Introduction*, Princeton University Press, 2003.
-  M. D. Springer and W. E. Thompson, *The distribution of products of independent random variables*, SIAM J. Appl. Math. **14** (1966) 511–526. http://www.jstor.org/stable/2946226?seq=1#page_scan_tab_contents.
-  K. Stromberg, *Probabilities on a compact group*, Trans. Amer. Math. Soc. **94** (1960), 295–309. http://www.jstor.org/stable/1993313?seq=1#page_scan_tab_contents.
-  P. R. Turner, *The distribution of leading significant digits*, IMA J. Numer. Anal. **2** (1982), no. 4, 407–412.