

# Completeness of Positive Linear Recurrence Sequences

Elżbieta Bołdyriew (eboldyriew@colgate.edu)

John Haviland (haviwj@umich.edu)

Phuc Lam (plam6@u.rochester.edu)

John Lentfer (jlentfer@hmc.edu)

Fernando Trejos Suárez (fernando.trejos@yale.edu)

Joint work with Steven J. Miller

The Nineteenth International Conference on Fibonacci Numbers  
and Their Applications  
07/21/2020

# Introduction

# Positive Linear Recurrence Sequences

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  of positive integers is a **Positive Linear Recurrence Sequence (PLRS)** if the following properties hold:

# Positive Linear Recurrence Sequences

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  of positive integers is a **Positive Linear Recurrence Sequence (PLRS)** if the following properties hold:

- (Recurrence relation) There are non-negative integers  $L, c_1, \dots, c_L$  such that

$$H_{n+1} = c_1 H_n + \dots + c_L H_{n+1-L}$$

with  $L, c_1, c_L$  positive.

# Positive Linear Recurrence Sequences

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  of positive integers is a **Positive Linear Recurrence Sequence (PLRS)** if the following properties hold:

- (Recurrence relation) There are non-negative integers  $L, c_1, \dots, c_L$  such that

$$H_{n+1} = c_1 H_n + \dots + c_L H_{n+1-L}$$

with  $L, c_1, c_L$  positive.

- (Initial conditions)  $H_1 = 1$ , and for  $1 \leq n < L$ ,

$$H_{n+1} = c_1 H_n + \dots + c_n H_1 + 1$$

# Positive Linear Recurrence Sequences

- We write  $[c_1, \dots, c_L]$  for  $H_{n+1} = c_1 H_n + \dots + c_L H_{n-L+1}$ .

# Positive Linear Recurrence Sequences

- We write  $[c_1, \dots, c_L]$  for  $H_{n+1} = c_1 H_n + \dots + c_L H_{n-L+1}$ .
- For example, for the Fibonacci numbers, we write  $[1, 1]$ . This definition gives initial conditions  $F_1 = 1, F_2 = 2$ .

# Positive Linear Recurrence Sequences

- We write  $[c_1, \dots, c_L]$  for  $H_{n+1} = c_1 H_n + \dots + c_L H_{n-L+1}$ .
- For example, for the Fibonacci numbers, we write  $[1, 1]$ . This definition gives initial conditions  $F_1 = 1, F_2 = 2$ .
- Despite satisfying positive linear recurrences, the Lucas and Pell numbers are not PLRS, since their initial conditions do not meet the definition.

# Introduction to Completeness

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  is called **complete** if every positive integer is a sum of its terms, using each term at most once.

# Introduction to Completeness

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  is called **complete** if every positive integer is a sum of its terms, using each term at most once.

- The sequence with the recurrence  $[1, 3]$  is *not* complete. Its terms are  $\{1, 2, 5, 11, \dots\}$ ; you cannot get 4 or 9 as the sequence grows too quickly.

# Introduction to Completeness

## Definition

A sequence  $\{H_i\}_{i \geq 1}$  is called **complete** if every positive integer is a sum of its terms, using each term at most once.

- The sequence with the recurrence  $[1, 3]$  is *not* complete. Its terms are  $\{1, 2, 5, 11, \dots\}$ ; you cannot get 4 or 9 as the sequence grows too quickly.
- The Fibonacci sequence  $F_{n+1} = F_n + F_{n-1}$ , with initial conditions  $F_1 = 1, F_2 = 2$ , is complete (follows from Zeckendorf's Theorem).

## The Doubling Sequence

The PLRS [2], which has the recurrence  $H_{n+1} = 2H_n$ , has terms  $H_n = 2^{n-1}$  and is complete because every integer has a binary representation.

# The Doubling Sequence

The PLRS [2], which has the recurrence  $H_{n+1} = 2H_n$ , has terms  $H_n = 2^{n-1}$  and is complete because every integer has a binary representation.

## Theorem (Brown)

*The complete sequence with maximal terms is  $H_n = 2^{n-1}$ .*

# The Doubling Sequence

The PLRS [2], which has the recurrence  $H_{n+1} = 2H_n$ , has terms  $H_n = 2^{n-1}$  and is complete because every integer has a binary representation.

## Theorem (Brown)

*The complete sequence with maximal terms is  $H_n = 2^{n-1}$ .*

Any PLRS of the form  $[1, \dots, 1, 2]$  has the same terms as [2], i.e.,  $H_n = 2^{n-1}$ .

# Brown's Criterion

## Theorem (Brown)

*A nondecreasing sequence  $\{H_i\}_{i \geq 1}$  is complete if and only if  $H_1 = 1$  and for every  $n \geq 1$ ,*

$$H_{n+1} \leq 1 + \sum_{i=1}^n H_i.$$

# Brown's Criterion

## Theorem (Brown)

*A nondecreasing sequence  $\{H_i\}_{i \geq 1}$  is complete if and only if  $H_1 = 1$  and for every  $n \geq 1$ ,*

$$H_{n+1} \leq 1 + \sum_{i=1}^n H_i.$$

Can we bound where a sequence must fail Brown's Criterion?

# Brown's Criterion

## Theorem (Brown)

*A nondecreasing sequence  $\{H_i\}_{i \geq 1}$  is complete if and only if  $H_1 = 1$  and for every  $n \geq 1$ ,*

$$H_{n+1} \leq 1 + \sum_{i=1}^n H_i.$$

Can we bound where a sequence must fail Brown's Criterion? We think so!

## Conjecture (SMALL 2020)

*If a PLRS  $H_{n+1} = c_1 H_n + \cdots + c_L H_{n+1-L}$  is incomplete, then it fails Brown's criterion before the  $2L$ -th term.*

# Families of Sequences

# Analyzing Families of Sequences

## Theorem (SMALL 2020)

- ①  $[1, \underbrace{0, \dots, 0}_k, N]$ , is complete if and only if

$$N \leq \left\lfloor \frac{(k+2)(k+3)}{4} + \frac{1}{2} \right\rfloor.$$

- ②  $[1, 1, \underbrace{0, \dots, 0}_k, N]$ , is complete if and only if

$$N \leq \left\lfloor \frac{F_{k+6} - (k+5)}{4} \right\rfloor,$$

where  $F_k$  is the  $k$ th Fibonacci number.

# Proof Sketch

## Theorem (SMALL 2020)

- ①  $[1, 0, \dots, 0, N]$ , with  $k$  zeros, is complete if and only if
- $$N \leq \left\lfloor \frac{(k+2)(k+3)}{4} + \frac{1}{2} \right\rfloor.$$

*Partial Proof.* We sketch that if  $N_{\max} = \left\lfloor \frac{(k+2)(k+3)}{4} + \frac{1}{2} \right\rfloor$ , then the sequence is complete. It is similar for  $N < N_{\max}$ .

# Proof Sketch

## Theorem (SMALL 2020)

- ①  $[1, 0, \dots, 0, N]$ , with  $k$  zeros, is complete if and only if
- $$N \leq \left\lfloor \frac{(k+2)(k+3)}{4} + \frac{1}{2} \right\rfloor.$$

*Partial Proof.* We sketch that if  $N_{\max} = \left\lfloor \frac{(k+2)(k+3)}{4} + \frac{1}{2} \right\rfloor$ , then the sequence is complete. It is similar for  $N < N_{\max}$ . With the recurrence relation and Brown's Criterion,

$$\begin{aligned} H_{n+1} &= H_n + N_{\max} H_{n-k-1} \\ &\leq H_n + (N_{\max} - 1)H_{n-k-1} + H_{n-k-2} + \dots + H_1 + 1 \end{aligned}$$

By induction,  $(N_{\max} - 1)H_{n-k-1} \leq H_{n-1} + \dots + H_{n-k-1}$ , so

$$\leq H_n + \dots + H_1 + 1.$$

## Example for $L = 6$

By the previous theorem,  $[1, 0, 0, 0, 0, N]$  is complete for  $N \leq 11$ .

## Example for $L = 6$

By the previous theorem,  $[1, 0, 0, 0, 0, N]$  is complete for  $N \leq 11$ .

### Question

Does there exist a complete PLRS of length  $L = 6$  with  $N > 11$ ?

## Example for $L = 6$

Here are the maximal last terms for preserving completeness for several other sequences of length  $L = 6$ :

## Example for $L = 6$

Here are the maximal last terms for preserving completeness for several other sequences of length  $L = 6$ :

- $[1, 0, 0, 0, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 1, 0, 0, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 0, 1, 0, 0, N]$  is complete for  $N \leq 12$ .
- $[1, 0, 0, 1, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 0, 0, 0, 1, N]$  is complete for  $N \leq 10$ .

## Example for $L = 6$

Here are the maximal last terms for preserving completeness for several other sequences of length  $L = 6$ :

- $[1, 0, 0, 0, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 1, 0, 0, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 0, 1, 0, 0, N]$  is complete for  $N \leq 12$ .
- $[1, 0, 0, 1, 0, N]$  is complete for  $N \leq 11$ .
- $[1, 0, 0, 0, 1, N]$  is complete for  $N \leq 10$ .

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

## Example for $L = 6$

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

## Example for $L = 6$

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

- $[1, 0, 0, 0, 0, 12]$  has terms  $\{1, 2, 3, 4, 5, 6, 18, 42, \dots\}$  and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 11, 16, 22, 40, \dots\}$

## Example for $L = 6$

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

- $[1, 0, 0, 0, 0, 12]$  has terms  $\{1, 2, 3, 4, 5, 6, 18, 42, \dots\}$   
and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 11, 16, 22, 40, \dots\}$
- $[1, 0, 1, 0, 0, 12]$  has terms  $\{1, 2, 3, 5, 8, 12, 29, 61, \dots\}$   
and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 12, 20, 32, 61, \dots\}$

## Example for $L = 6$

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

- $[1, 0, 0, 0, 0, 12]$  has terms  $\{1, 2, 3, 4, 5, 6, 18, 42, \dots\}$  and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 11, 16, 22, 40, \dots\}$
- $[1, 0, 1, 0, 0, 12]$  has terms  $\{1, 2, 3, 5, 8, 12, 29, 61, \dots\}$  and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 12, 20, 32, 61, \dots\}$
- $[1, 1, 1, 0, 0, 12]$  has terms  $\{1, 2, 4, 8, 15, 28, 63, \dots\}$  and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 8, 16, 31, 59, \dots\}$

## Example for $L = 6$

Why is  $[1, 0, 1, 0, 0, 12]$  complete, but  $[1, 0, 0, 0, 0, 12]$  is not complete?

- $[1, 0, 0, 0, 0, 12]$  has terms  $\{1, 2, 3, 4, 5, 6, 18, 42, \dots\}$   
and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 11, 16, 22, 40, \dots\}$
- $[1, 0, 1, 0, 0, 12]$  has terms  $\{1, 2, 3, 5, 8, 12, 29, 61, \dots\}$   
and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 7, 12, 20, 32, 61, \dots\}$
- $[1, 1, 1, 0, 0, 12]$  has terms  $\{1, 2, 4, 8, 15, 28, 63, \dots\}$   
and so computing the sums  $\sum_{i=1}^n H_i + 1$  we see  $\{2, 4, 8, 16, 31, 59, \dots\}$

# Sequences of Initial Ones

## Theorem (SMALL 2020)

If a sequence  $[1, \dots, 1, \underbrace{0, \dots, 0}_k, N]$  is complete with  $m \geq 3$ , then

$$N \leq \frac{1}{2} \left( 1 + \sum_{i=1}^{k+1} F_i^{(m)} + \sum_{i=1}^{k+1-m} F_i^{(m)} + \dots + \sum_{i=1}^{(k+1) \bmod m} F_i^{(m)} \right)$$

where  $F_i^{(m)}$  is the  $m$ -bonacci sequence,  $[1, \dots, 1]_m$ .

# Theorem on Adding Ones

## Theorem (SMALL 2020)

- For  $L \geq 6$ , consider the sequence  $\{H_n\}$  given by  $[1, 0, \dots, 0, 1, 0, \dots, 0, M]$ . Then, if  $M$  is maximal such that  $\{H_n\}$  is complete, and  $N$  is maximal such that  $[1, 0, \dots, 0, N]$  is complete, we have  $M \geq N$ .
- For a fixed length  $L$ , the sequence  $[1, \underbrace{0, \dots, 0}_k, \underbrace{1, \dots, 1}_m, N]$  with  $m$  ones has a lower bound on  $N$  than the sequence  $[1, \underbrace{0, \dots, 0}_{k-1}, \underbrace{1, \dots, 1}_{m+1}, N]$ .

In particular, if  $m < \frac{L}{2}$ , the bound is precisely

$$N \leq \left\lfloor \frac{(L-m)(L+m+1)}{4} + \frac{1}{48}m(m+1)(m+2)(m+3) + \frac{1-2m}{2} \right\rfloor.$$

# Modifying Sequences

## Modifying Coefficients of a PLRS

When studying a PLRS, what modifications to the recurrence coefficients preserve completeness or incompleteness?

## Modifying Coefficients of a PLRS

When studying a PLRS, what modifications to the recurrence coefficients preserve completeness or incompleteness?

### Theorem (SMALL 2020)

- If a sequence  $[c_1, \dots, c_{L-1}, c_L]$  is complete, then so is  $[c_1, \dots, c_{L-1}, d_L]$  for any  $d_L \leq c_L$ .

*Remark. This is not true for  $c_i$  in any position.*

- If a sequence  $[1, \dots, 1, 0, \dots, 0, c_L]$  is complete and

$c_L = 2^{k+1} - 1$ ,  $[1, \dots, 1, 0, \dots, 0, c_L + j]$  is incomplete

for any positive integer  $j$ .

## Modifying Lengths of a PLRS

### Theorem (SMALL 2020)

- *If a sequence  $[c_1, \dots, c_L]$  is incomplete, then so is  $[c_1, \dots, c_{L-1} + c_L]$ .*
- *If a sequence  $[c_1, \dots, c_L]$  is incomplete, then so is  $[c_1, \dots, c_L, c_{L+1}]$  for any  $c_{L+1} > 0$ .*

# Modifying Lengths of a PLRS

## Theorem (SMALL 2020)

- If a sequence  $[c_1, \dots, c_L]$  is incomplete, then so is  $[c_1, \dots, c_{L-1} + c_L]$ .
- If a sequence  $[c_1, \dots, c_L]$  is incomplete, then so is  $[c_1, \dots, c_L, c_{L+1}]$  for any  $c_{L+1} > 0$ .

## Conjecture (SMALL 2020)

If a sequence  $[1, \dots, 1, \underbrace{0, \dots, 0}_m, c_L]$  is complete, then so is  $[\underbrace{1, \dots, 1}_{m+j}, \underbrace{0, \dots, 0}_k, c_L]$  for any positive integer  $j$ .

# Principal Roots

# Principal Roots

## Theorem (Binet's Formula)

*If  $r_1, \dots, r_k$  are the distinct roots of the characteristic polynomial of a PLRS  $\{H_n\}$ , then there exist polynomials  $q_1, \dots, q_k$  such that  $H_n = q_1(n)r_1^n + \dots + q_k(n)r_k^n$ .*

# Principal Roots

## Theorem (Binet's Formula)

*If  $r_1, \dots, r_k$  are the distinct roots of the characteristic polynomial of a PLRS  $\{H_n\}$ , then there exist polynomials  $q_1, \dots, q_k$  such that  $H_n = q_1(n)r_1^n + \dots + q_k(n)r_k^n$ .*

For PLRS, the characteristic polynomial has a unique positive root  $r_1$  which is the largest in absolute value, called the *principal root*.

# Principal Roots

## Theorem (Binet's Formula)

*If  $r_1, \dots, r_k$  are the distinct roots of the characteristic polynomial of a PLRS  $\{H_n\}$ , then there exist polynomials  $q_1, \dots, q_k$  such that  $H_n = q_1(n)r_1^n + \dots + q_k(n)r_k^n$ .*

For PLRS, the characteristic polynomial has a unique positive root  $r_1$  which is the largest in absolute value, called the *principal root*.

## Theorem (SMALL 2020)

*If  $H_n$  is a complete PLRS and  $r_1$  is its principal root, then  $r_1 \leq 2$ .*

## Bounding Principal Roots

- If a sequence is complete,  $r_1 \leq 2$ .

## Bounding Principal Roots

- If a sequence is complete,  $r_1 \leq 2$ .
- There exists a second bound  $1 < B_L < 2$  on the principal roots, so that if a sequence is incomplete, the its principal root  $r_1$  satisfies  $r_1 \geq B_L$ . This bound is dependent on the length of the generating sequence  $[c_1, \dots, c_L]$ . We conjecture the following:

### Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \left\lceil \frac{L(L+1)}{4} \right\rceil + 1]$ .*

## Bounding Principal Roots

- If a sequence is complete,  $r_1 \leq 2$ .
- There exists a second bound  $1 < B_L < 2$  on the principal roots, so that if a sequence is incomplete, the its principal root  $r_1$  satisfies  $r_1 \geq B_L$ . This bound is dependent on the length of the generating sequence  $[c_1, \dots, c_L]$ . We conjecture the following:

### Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \left\lceil \frac{L(L+1)}{4} \right\rceil + 1]$ .*

- If this holds, then for large  $L$ , we would have  $B_L \approx (L/2)^{2/L}$ . In particular,  $\lim_{L \rightarrow \infty} B_L = 1$ .

# Root-Bounding Proof Sketch

## Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \left\lceil \frac{L(L+1)}{4} \right\rceil + 1]$ .*

# Root-Bounding Proof Sketch

## Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \lceil \frac{L(L+1)}{4} \rceil + 1]$ .*

Suppose  $[c_1, \dots, c_L]$  is an incomplete sequence.

Case 1:  $\sum_{k=1}^L c_k \geq 2 + \lceil \frac{L(L+1)}{4} \rceil$

We combine the following two invariant arguments:

# Root-Bounding Proof Sketch

## Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \lceil \frac{L(L+1)}{4} \rceil + 1]$ .*

Suppose  $[c_1, \dots, c_L]$  is an incomplete sequence.

Case 1:  $\sum_{k=1}^L c_k \geq 2 + \lceil \frac{L(L+1)}{4} \rceil$

We combine the following two invariant arguments:

- The principal root of  $[c_1, \dots, c_L]$  is strictly greater than that of  $[c_1, \dots, c_k - 1, \dots, c_L + 1]$ , for any  $k$ .

# Root-Bounding Proof Sketch

## Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \lceil \frac{L(L+1)}{4} \rceil + 1]$ .*

Suppose  $[c_1, \dots, c_L]$  is an incomplete sequence.

Case 1:  $\sum_{k=1}^L c_k \geq 2 + \lceil \frac{L(L+1)}{4} \rceil$

We combine the following two invariant arguments:

- The principal root of  $[c_1, \dots, c_L]$  is strictly greater than that of  $[c_1, \dots, c_k - 1, \dots, c_L + 1]$ , for any  $k$ .
- The principal root of  $[1, 0, \dots, 0, S]$  is strictly greater than that of  $[1, 0, \dots, 0, S - 1]$ .

# Root-Bounding Proof Sketch

## Conjecture (SMALL 2020)

*For any given  $L$ , the incomplete sequence of length  $L$  with the lowest principal root is  $[1, 0, \dots, 0, \lceil \frac{L(L+1)}{4} \rceil + 1]$ .*

Suppose  $[c_1, \dots, c_L]$  is an incomplete sequence.

Case 1:  $\sum_{k=1}^L c_k \geq 2 + \lceil \frac{L(L+1)}{4} \rceil$

We combine the following two invariant arguments:

- The principal root of  $[c_1, \dots, c_L]$  is strictly greater than that of  $[c_1, \dots, c_k - 1, \dots, c_L + 1]$ , for any  $k$ .
- The principal root of  $[1, 0, \dots, 0, S]$  is strictly greater than that of  $[1, 0, \dots, 0, S - 1]$ .

Combining these two, any sequence with large sum can be "reduced" to  $[1, 0, \dots, 0, \lceil \frac{L(L+1)}{4} \rceil + 1]$ .

## Root-Bounding Proof Sketch

Case 2:  $\sum_{k=1}^L c_k \leq 1 + \left\lceil \frac{L(L+1)}{4} \right\rceil$

It can be shown any “counterexample” would fulfill:

- $\forall 1 \leq k \leq L + 1,$

$$\sum_{i=2}^k c_i \leq \left\lceil \frac{k(k+1)}{4} \right\rceil.$$

- $\sum_{i=2}^L c_i (\lambda_{L+1}^{L+1-i} - \lambda_L^{L-i}) < \frac{L+2}{2}$ , where  $\lambda_L$  is the root of  $[1, 0, \dots, 0, \lceil L(L+1)/4 \rceil + 1$ .

This forces the coefficients of  $[c_1, \dots, c_L]$  to be small enough to force a contradiction; for example, an analytical argument shows the first 32.5% or so must be 0.

## Future Directions

# Future Directions

- Extend analysis of the bound of  $N$  in  $[\underbrace{1, \dots, 1}_m, 0, \dots, 0, N]$ , which involves the  $m$ -bonacci numbers, defined by  $[\underbrace{1, \dots, 1}_m]$ .

## Future Directions

- Extend analysis of the bound of  $N$  in  $[\underbrace{1, \dots, 1}_m, 0, \dots, 0, N]$ , which involves the  $m$ -bonacci numbers, defined by  $[\underbrace{1, \dots, 1}_m]$ .
- Find the bound  $N$  for arbitrary coefficients  $c_2, \dots, c_{L-1}$  in  $[1, c_2, \dots, c_{L-1}, N]$ .

## Future Directions

- Extend analysis of the bound of  $N$  in  $[\underbrace{1, \dots, 1}_m, 0, \dots, 0, N]$ , which involves the  $m$ -bonacci numbers, defined by  $[\underbrace{1, \dots, 1}_m]$ .
- Find the bound  $N$  for arbitrary coefficients  $c_2, \dots, c_{L-1}$  in  $[1, c_2, \dots, c_{L-1}, N]$ .
- Prove the conjectures made in this presentation.

# Bibliography

-  Thomas C. Martinez, Steven J. Miller, Clay Mizgerd, and Chenyang Sun. Generalizing Zeckendorf's Theorem to Homogeneous Linear Recurrences, 2020
-  Olivia Beckwith, Amanda Bower, Louis Gaudet, Rachel Insoft, Shiyu Li, Steven J. Miller, and Philip Tosteson. The Average Gap Distribution for Generalized Zeckendorf Decompositions, Dec 2012.
-  J. L. Brown. Note on complete sequences of integers. *The American Mathematical Monthly*, 68(6):557, 1961.

# Acknowledgements

- Thank you. Any questions?
- This research was conducted as part of the 2020 SMALL REU program at Williams College. This work was supported by NSF Grant DMS1947438, Williams, Yale, and Rochester.

## Legal Decompositions vs. Completeness

- Previous work on PLRS relates to *legal decompositions*, which are another way to write integers as sums of sequence terms.
- Given any PLRS, there is a legal decomposition of every positive integer. Does this mean that all PLRS are complete?
- No. For legal decompositions, sequence terms can be used more than once. This is not allowed for completeness decompositions.

### Example

The PLRS  $[1, 3]$  has terms  $1, 2, 5, 11, \dots$ . The unique *legal* decomposition for 9 is  $5 + 2(2)$ , where the term 2 is used twice. However, no *complete* decomposition for 9 exists.