

# From M&Ms to Mathematics, or, How I learned to answer questions and help my kids love math.

Steven J. Miller, Williams College

`sjml@williams.edu,`

`Steven.Miller.MC.96@aya.yale.edu`

`http://web.williams.edu/Mathematics/sjmillier/public\_html/`

**Texas State REU (zoom), June 27, 2024**



## Some Issues for the Future

- World is rapidly changing – powerful computing cheaply and readily available.
- What skills are we teaching? What skills should we be teaching?
- One of hardest skills: how to think / attack a new problem, how to see connections, what data to gather.

## Goals of the Talk: Opportunities Everywhere!

- Ask Questions! Often simple questions lead to good math.
- Gather data: observe, program and simulate.
- Use games to get to mathematics.
- Discuss implementation: `Please interrupt!`

Joint work with Cameron (age 10) and Kayla (age 8) Miller:  
Erdős number 3

Problem Editor Pi Mu Epsilon Journal, and my math riddles  
page: <http://mathriddles.williams.edu/>

## Pre-requisite: Logarithms

The logarithm of  $x$  base  $b$  is the power we raise  $b$  to get  $x$ ; it is the inverse of exponentiation.

$$x = b^y \text{ if and only if } \log_b(x) = y.$$

$$\log_{10}(10^4) = 4, \quad \log_{10}(1/100) = -2.$$

### Properties:

- $\log_{10} 1 = 0.$
- $10^{\log_{10} x} = x.$
- $\log_{10}(A \cdot B) = \log_{10} A + \log_{10} B.$

## Pre-requisites: Combinatorics Review

- $n!$ : number of ways to order  $n$  people, order matters.
- $\frac{n!}{k!(n-k)!} = nCk = \binom{n}{k}$ : number of ways to choose  $k$  from  $n$ , order doesn't matter.
- Examples:  $\binom{n}{1} = n$ ,  $\binom{4}{2} = 6$ , in general  $\binom{n}{2} = \frac{n(n-1)}{2}$ .

## Pre-requisites: Pascal's Triangle

Pascal's Triangle:  $k^{\text{th}}$  entry in row  $n$  is  $\binom{n}{k}$ .

## Pre-requisites: Pascal's Triangle


Pascal's Triangle:  $k^{\text{th}}$  entry in row  $n$  is  $\binom{n}{k}$ .

				1											
			1	1											
		1	2	1											
	1	3	3	1											
	1	4	6	4	1										
	1	5	10	10	5	1									
	1	6	15	20	15	6	1								
	1	7	21	35	35	21	7	1							
	1	8	28	56	70	56	28	8	1						
	1	9	36	84	126	126	84	36	9	1					
	1	10	45	120	210	252	210	120	45	10	1				
	1	11	55	165	330	462	462	330	165	55	11	1			
	1	12	66	220	495	792	924	792	495	220	66	12	1		
	1	13	78	286	715	1287	1716	1716	1287	715	286	78	13	1	
	1	14	91	364	1001	2002	3003	3432	3003	2002	1001	364	91	14	1
1	15	105	455	1365	3003	5005	6435	6435	5005	3003	1365	455	105	15	1

Pascal's Triangle

## Pre-requisites: Pascal's Triangle

Pascal's Triangle:  $k^{\text{th}}$  entry in row  $n$  is  $\binom{n}{k}$ .



Blocking Out the Odd Entries  
to create a Sierpinski Triangle




## Pre-requisites: Pascal's Triangle

Pascal's Triangle:  $k^{\text{th}}$  entry in row  $n$  is  $\binom{n}{k}$ .

Sierpinski's triangle: Look at Pascal's triangle modulo 2:

[https://www.youtube.com/watch?v=tt4\\_4YajqRM](https://www.youtube.com/watch?v=tt4_4YajqRM).




## The M&M Game

## Motivating Question

**Cam (4 years):** If you're born on the same day, do you die on the same day?

## M&M Game Rules

**Cam (4 years):** If you're born on the same day, do you die on the same day?



- (1) Everyone starts off with  $k$  M&Ms (we did 5).
- (2) All toss fair coins, eat an M&M if and only if head.



**Be active – ask questions!**

**What are natural questions to ask?**

# Be active – ask questions!

**What are natural questions to ask?**

**Question 1:** How likely is a tie (as a function of  $k$ )?


**Question 2:** How long until one dies?

**Question 3:** Generalize the game: More people? Biased coin?

Important to ask questions – curiosity is good and to be encouraged! Value to the journey and not knowing the answer.


Let's gather some data! Let's play!

## Probability of a tie in the M&M game (2 players)



Prob(tie)  $\approx$  33% (1 M&M), 19% (2 M&Ms), 14% (3 M&Ms), 10% (4 M&Ms).


## Probability of a tie in the M&M game (2 players)



Gave at a 110th anniversary talk....



## Probability of a tie in the M&M game (2 players)




... asked them: what will the next 110 bring us?  
Never too early to lay foundations for future classes.

## Welcome to Statistics and Inference!


- ◇ **Goal:** Gather data, see pattern, extrapolate.
- ◇ **Methods:** Simulation, analysis of special cases.
- ◇ **Presentation:** It matters **how** we show data, and **which** data we show.

# Viewing M&M Plots




Hard to predict what comes next.

## Viewing M&M Plots: Log-Log Plot



Not *just* sadistic teachers: logarithms useful!


## Viewing M&M Plots: Log-Log Plot



**Best fit line:**

$$\log(\text{Prob}(\text{tie})) = -1.42022 - 0.545568 \log(\#M\&Ms) \text{ or}$$
$$\text{Prob}(k) \approx 0.2412/k^{.5456}.$$

## Viewing M&M Plots: Log-Log Plot



**Best fit line:**

$$\log(\text{Prob}(\text{tie})) = -1.42022 - 0.545568 \log(\#M\&Ms) \text{ or} \\ \text{Prob}(k) \approx 0.2412/k^{.5456}.$$


Predicts probability of a tie when  $k = 220$  is 0.01274, but answer is 0.0137. **What gives?**

# Statistical Inference: Too Much Data Is Bad!

Small values can mislead / distort. Let's go from  $k = 50$  to 110.

## Statistical Inference: Too Much Data Is Bad!

Small values can mislead / distort. Let's go from  $k = 50$  to 110.




**Best fit line:**

$$\log(\text{Prob}(\text{tie})) = -1.58261 - 0.50553 \log(\#M\&Ms) \text{ or}$$
$$\text{Prob}(k) \approx 0.205437/k^{.50553} \text{ (had } 0.241662/k^{.5456}\text{)}.$$



## Statistical Inference: Too Much Data Is Bad!

Small values can mislead / distort. Let's go from  $k = 50$  to 110.



**Best fit line:**

$$\log(\text{Prob}(\text{tie})) = -1.58261 - 0.50553 \log(\#M\&Ms) \text{ or}$$

$$\text{Prob}(k) \approx 0.205437/k^{.50553} \text{ (had } 0.241662/k^{.5456} \text{)}.$$

Get 0.01344 for  $k = 220$  (answer 0.01347); **much better!**

From Shooting Hoops  
to the Geometric Series Formula

# Simpler Game: Hoops

Game of hoops: first basket wins, alternate shooting.



## Simpler Game: Hoops: Mathematical Formulation

**Bird** and **Magic** (I'm old!) alternate shooting; first basket wins.

- **Bird** always gets basket with probability  $p$ .
- **Magic** always gets basket with probability  $q$ .

Let  $x$  be the probability **Bird** wins – what is  $x$ ?

## Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

# Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

- **Bird** wins on 1<sup>st</sup> shot:  $p$ .

## Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

- **Bird** wins on 1<sup>st</sup> shot:  $p$ .
- **Bird** wins on 2<sup>nd</sup> shot:  $(1 - p)(1 - q) \cdot p$ .

## Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

- **Bird** wins on 1<sup>st</sup> shot:  $p$ .
- **Bird** wins on 2<sup>nd</sup> shot:  $(1 - p)(1 - q) \cdot p$ .
- **Bird** wins on 3<sup>rd</sup> shot:  $(1 - p)(1 - q) \cdot (1 - p)(1 - q) \cdot p$ .



## Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

- **Bird** wins on 1<sup>st</sup> shot:  $p$ .
- **Bird** wins on 2<sup>nd</sup> shot:  $(1 - p)(1 - q) \cdot p$ .
- **Bird** wins on 3<sup>rd</sup> shot:  $(1 - p)(1 - q) \cdot (1 - p)(1 - q) \cdot p$ .
- **Bird** wins on  $n^{\text{th}}$  shot:  
 $(1 - p)(1 - q) \cdot (1 - p)(1 - q) \cdots (1 - p)(1 - q) \cdot p$ .

## Solving the Hoop Game

Classic solution involves the geometric series.

Break into cases:

- **Bird** wins on 1<sup>st</sup> shot:  $p$ .
- **Bird** wins on 2<sup>nd</sup> shot:  $(1 - p)(1 - q) \cdot p$ .
- **Bird** wins on 3<sup>rd</sup> shot:  $(1 - p)(1 - q) \cdot (1 - p)(1 - q) \cdot p$ .
- **Bird** wins on  $n^{\text{th}}$  shot:  
 $(1 - p)(1 - q) \cdot (1 - p)(1 - q) \cdots (1 - p)(1 - q) \cdot p$ .

Let  $r = (1 - p)(1 - q)$ . Then

$$\begin{aligned}x &= \text{Prob}(\mathbf{Bird} \text{ wins}) \\ &= p + rp + r^2p + r^3p + \dots \\ &= p(1 + r + r^2 + r^3 + \dots),\end{aligned}$$

the geometric series.

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p +$$

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p + (1 - p)(1 - q)$$

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p + (1 - p)(1 - q)x$$

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p + (1 - p)(1 - q)x = p + rx.$$

## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p + (1 - p)(1 - q)x = p + rx.$$

Thus

$$(1 - r)x = p \quad \text{or} \quad x = \frac{p}{1 - r}.$$



## Solving the Hoop Game: The Power of Perspective

Showed

$$x = \text{Prob}(\text{Bird wins}) = p(1 + r + r^2 + r^3 + \dots);$$

will solve **without** the geometric series formula.

Have

$$x = \text{Prob}(\text{Bird wins}) = p + (1 - p)(1 - q)x = p + rx.$$

Thus

$$(1 - r)x = p \quad \text{or} \quad x = \frac{p}{1 - r}.$$

As  $x = p(1 + r + r^2 + r^3 + \dots)$ , find

$$1 + r + r^2 + r^3 + \dots = \frac{1}{1 - r}.$$

## Lessons from Hoop Problem

- ◇ Power of Perspective: Memoryless process.
- ◇ Can circumvent algebra with deeper understanding! (Hard)
- ◇ Depth of a problem not always what expect.
- ◇ Importance of knowing more than the minimum: [connections](#).
- ◇ Math is fun!

## The M&M Game

## Solving the M&M Game

**Overpower with algebra:** Assume  $k$  M&Ms, two people, fair coins:

$$\text{Prob}(\text{tie}) = \sum_{n=k}^{\infty} \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2} \cdot \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2},$$

where

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

is a binomial coefficient.

## Solving the M&M Game

**Overpower with algebra:** Assume  $k$  M&Ms, two people, fair coins:

$$\text{Prob}(\text{tie}) = \sum_{n=k}^{\infty} \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2} \cdot \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2},$$

where

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

is a binomial coefficient.

“Simplifies” to  $4^{-k} {}_2F_1(k, k, 1, 1/4)$ , a special value of a hypergeometric function! (Look up / write report.)

A look at your future classes, but is there a better way?

## Solving the M&M Game (cont)

Where did formula come from? Each turn one of four **equally likely** events happens:

- Both eat an M&M.
- Cam eats and M&M but Kayla does not.
- Kayla eats an M&M but Cam does not.
- Neither eat.

Probability of each event is  $1/4$  or  $25\%$ .

## Solving the M&M Game (cont)

Where did formula come from? Each turn one of four **equally likely** events happens:

- Both eat an M&M.
- Cam eats and M&M but Kayla does not.
- Kayla eats an M&M but Cam does not.
- Neither eat.

Probability of each event is  $1/4$  or  $25\%$ .

Each person has exactly  $k - 1$  heads in first  $n - 1$  tosses, then ends with a head.

$$\text{Prob}(\text{tie}) = \sum_{n=k}^{\infty} \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2} \cdot \binom{n-1}{k-1} \left(\frac{1}{2}\right)^{n-1} \frac{1}{2}$$



# Solving the M&M Game (cont)

Use the lesson from the Hoops Game: Memoryless process!



## Solving the M&M Game (cont)

Use the lesson from the Hoops Game: Memoryless process!

If neither eat, as if toss didn't happen. Now game is finite.

## Solving the M&M Game (cont)

Use the lesson from the Hoops Game: Memoryless process!

If neither eat, as if toss didn't happen. Now game is finite.

Much better perspective: each "turn" one of **three equally likely** events happens:

- Both eat an M&M.
- Cam eats and M&M but Kayla does not.
- Kayla eats an M&M but Cam does not.

Probability of each event is  $1/3$  or about **33%**

$$\sum_{n=0}^{k-1} \binom{2k-n-2}{n} \left(\frac{1}{3}\right)^n \binom{2k-2n-2}{k-n-1} \left(\frac{1}{3}\right)^{k-n-1} \left(\frac{1}{3}\right)^{k-n-1} \binom{1}{1} \frac{1}{3}.$$



## Solving the M&M Game (cont)


**Interpretation:** Let Cam have  $c$  M&Ms and Kayla have  $k$ ; write as  $(c, k)$ .

Then each of the following happens  $1/3$  of the time after a 'turn':

- $(c, k) \rightarrow (c - 1, k - 1)$ .
- $(c, k) \rightarrow (c - 1, k)$ .
- $(c, k) \rightarrow (c, k - 1)$ .




## Solving the M&M Game (cont): Assume $k = 4$




**Figure:** The M&M game when  $k = 4$ . Count the paths! Answer  $1/3$  of probability hit  $(1,1)$ .

## Solving the M&M Game (cont): Assume $k = 4$




**Figure:** The M&M game when  $k = 4$ , going down one level.

## Solving the M&M Game (cont): Assume $k = 4$




**Figure:** The M&M game when  $k = 4$ , removing probability from the second level.

## Solving the M&M Game (cont): Assume $k = 4$



**Figure:** Removing probability from two outer on third level.


## Solving the M&M Game (cont): Assume $k = 4$



**Figure:** Removing probability from the (3,2) and (2,3) vertices.




## Solving the M&M Game (cont): Assume $k = 4$




**Figure:** Removing probability from the  $(2,2)$  vertex.

## Solving the M&M Game (cont): Assume $k = 4$




**Figure:** Removing probability from the  $(4,1)$  and  $(1,4)$  vertices.

## Solving the M&M Game (cont): Assume $k = 4$



**Figure:** Removing probability from the  $(3,1)$  and  $(1,3)$  vertices.

## Solving the M&M Game (cont): Assume $k = 4$



**Figure:** Removing probability from  $(2, 1)$  and  $(1, 2)$  vertices. Answer is  $1/3$  of  $(1, 1)$  vertex, or  $245/2187$  (about 11%).

## Interpreting Proof: Connections to the Fibonacci Numbers!

**Fibonacci:**  $F_{n+2} = F_{n+1} + F_n$  with  $F_0 = 0, F_1 = 1$ .

Starts 0, 1, 1, 2, 3, 5, 8, 13, 21, . . . .

<http://www.youtube.com/watch?v=kkGeOWYOFoA>.

**Binet's Formula (can prove via 'generating functions'):**

$$F_n = \frac{1}{\sqrt{5}} \left( \frac{1 + \sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left( \frac{1 - \sqrt{5}}{2} \right)^n .$$

## Interpreting Proof: Connections to the Fibonacci Numbers!

**Fibonacci:**  $F_{n+2} = F_{n+1} + F_n$  with  $F_0 = 0, F_1 = 1$ .

Starts 0, 1, 1, 2, 3, 5, 8, 13, 21, ...

<http://www.youtube.com/watch?v=kkGeOWYOFoA>.

**Binet's Formula (can prove via 'generating functions'):**

$$F_n = \frac{1}{\sqrt{5}} \left( \frac{1 + \sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left( \frac{1 - \sqrt{5}}{2} \right)^n.$$

M&Ms: For  $c, k \geq 1$ :  $x_{c,0} = x_{0,k} = 0$ ;  $x_{0,0} = 1$ , and if  $c, k \geq 1$ :

$$x_{c,k} = \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}.$$

Reproduces the tree but a lot 'cleaner'.

## Interpreting Proof: Finding the Recurrence

What if we didn't see the 'simple' recurrence?

$$X_{c,k} = \frac{1}{3}X_{c-1,k-1} + \frac{1}{3}X_{c-1,k} + \frac{1}{3}X_{c,k-1}.$$

## Interpreting Proof: Finding the Recurrence

What if we didn't see the 'simple' recurrence?

$$x_{c,k} = \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}.$$

The following recurrence is 'natural':

$$x_{c,k} = \frac{1}{4}x_{c,k} + \frac{1}{4}x_{c-1,k-1} + \frac{1}{4}x_{c-1,k} + \frac{1}{4}x_{c,k-1}.$$



## Interpreting Proof: Finding the Recurrence

What if we didn't see the 'simple' recurrence?

$$x_{c,k} = \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}.$$

The following recurrence is 'natural':

$$x_{c,k} = \frac{1}{4}x_{c,k} + \frac{1}{4}x_{c-1,k-1} + \frac{1}{4}x_{c-1,k} + \frac{1}{4}x_{c,k-1}.$$

Obtain 'simple' recurrence by algebra: subtract  $\frac{1}{4}x_{c,k}$ :

$$\begin{aligned} \frac{3}{4}x_{c,k} &= \frac{1}{4}x_{c-1,k-1} + \frac{1}{4}x_{c-1,k} + \frac{1}{4}x_{c,k-1} \\ \text{therefore } x_{c,k} &= \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}. \end{aligned}$$

## Solving the Recurrence

$$X_{c,k} = \frac{1}{3}X_{c-1,k-1} + \frac{1}{3}X_{c-1,k} + \frac{1}{3}X_{c,k-1}.$$

## Solving the Recurrence

$$X_{c,k} = \frac{1}{3}X_{c-1,k-1} + \frac{1}{3}X_{c-1,k} + \frac{1}{3}X_{c,k-1}.$$

- $X_{0,0} = 1.$

## Solving the Recurrence

$$x_{c,k} = \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}.$$

- $x_{0,0} = 1$ .
- $x_{1,0} = x_{0,1} = 0$ .
- $x_{1,1} = \frac{1}{3}x_{0,0} + \frac{1}{3}x_{0,1} + \frac{1}{3}x_{1,0} = \frac{1}{3} \approx 33.3\%$ .

## Solving the Recurrence

$$x_{c,k} = \frac{1}{3}x_{c-1,k-1} + \frac{1}{3}x_{c-1,k} + \frac{1}{3}x_{c,k-1}.$$

- $x_{0,0} = 1.$
- $x_{1,0} = x_{0,1} = 0.$
- $x_{1,1} = \frac{1}{3}x_{0,0} + \frac{1}{3}x_{0,1} + \frac{1}{3}x_{1,0} = \frac{1}{3} \approx 33.3\%.$
- $x_{2,0} = x_{0,2} = 0.$
- $x_{2,1} = \frac{1}{3}x_{1,0} + \frac{1}{3}x_{1,1} + \frac{1}{3}x_{2,0} = \frac{1}{9} = x_{1,2}.$
- $x_{2,2} = \frac{1}{3}x_{1,1} + \frac{1}{3}x_{1,2} + \frac{1}{3}x_{2,1} = \frac{1}{9} + \frac{1}{27} + \frac{1}{27} = \frac{5}{27} \approx 18.5\%.$

## Try Simpler Cases!!!

Try and find an easier problem and build intuition.

## Try Simpler Cases!!!

Try and find an easier problem and build intuition.

Walking from  $(0,0)$  to  $(k, k)$  with allowable steps  $(1,0)$ ,  $(0,1)$  and  $(1,1)$ , hit  $(k, k)$  before hit top or right sides.

## Try Simpler Cases!!!

Try and find an easier problem and build intuition.

Walking from  $(0,0)$  to  $(k, k)$  with allowable steps  $(1,0)$ ,  $(0,1)$  and  $(1,1)$ , hit  $(k, k)$  before hit top or right sides.

Generalization of the Catalan problem. There don't have  $(1,1)$  and stay on or below the main diagonal.




## Try Simpler Cases!!!

Try and find an easier problem and build intuition.

Walking from  $(0,0)$  to  $(k,k)$  with allowable steps  $(1,0)$ ,  $(0,1)$  and  $(1,1)$ , hit  $(k,k)$  before hit top or right sides.

Generalization of the Catalan problem. There don't have  $(1,1)$  and stay on or below the main diagonal.



Interpretation: Catalan numbers are valid placings of ( and ).

## Aside: Fun Riddle Related to Catalan Numbers

Young Saul, a budding mathematician and printer, is making himself a fake ID. He needs it to say he's 21. The problem is he's not using a computer, but rather he has some symbols he's bought from the store, and that's it. He has one 1, one 5, one 6, one 7, and an unlimited supply of  $+$   $-$   $*$   $/$  (the operations addition, subtraction, multiplication and division). Using each number exactly once (but you can use any number of  $+$ , any number of  $-$ , ...) how, oh how, can he get 21 from 1,5, 6,7? Note: you can't do things like  $15+6 = 21$ . You have to use the four operations as 'binary' operations:  $((1+5)*6) + 7$ . Problem submitted by [ohadbp@infolink.net.il](mailto:ohadbp@infolink.net.il), phrasing by yours truly.

Solution involves valid sentences:  $((w + x) + y) + z, w + ((x + y) + z), \dots$

For more riddles see my riddles page:  
<http://mathriddles.williams.edu/>.

## Examining Probabilities of a Tie

When  $k = 1$ ,  $\text{Prob}(\text{tie}) = 1/3$ .

When  $k = 2$ ,  $\text{Prob}(\text{tie}) = 5/27$ .

When  $k = 3$ ,  $\text{Prob}(\text{tie}) = 11/81$ .

When  $k = 4$ ,  $\text{Prob}(\text{tie}) = 245/2187$ .

When  $k = 5$ ,  $\text{Prob}(\text{tie}) = 1921/19683$ .

When  $k = 6$ ,  $\text{Prob}(\text{tie}) = 575/6561$ .

When  $k = 7$ ,  $\text{Prob}(\text{tie}) = 42635/531441$ .

When  $k = 8$ ,  $\text{Prob}(\text{tie}) = 355975/4782969$ .

## Examining Ties: Multiply by $3^{2k-1}$ to clear denominators.

When  $k = 1$ , get 1.

When  $k = 2$ , get 5.

When  $k = 3$ , get 33.

When  $k = 4$ , get 245.

When  $k = 5$ , get 1921.

When  $k = 6$ , get 15525.

When  $k = 7$ , get 127905.

When  $k = 8$ , get 1067925.

# OEIS

Get sequence of integers: 1, 5, 33, 245, 1921, 15525, ....

# OEIS

Get sequence of integers: 1, 5, 33, 245, 1921, 15525, ....

OEIS: <http://oeis.org/>.

# OEIS

Get sequence of integers: 1, 5, 33, 245, 1921, 15525, ....

OEIS: <http://oeis.org/>.

Our sequence: <http://oeis.org/A084771>.

**The web exists!** Use it to build conjectures, suggest proofs....

## OEIS (continued)

A084771	Coefficients of $1/\sqrt{1-10*x+9*x^2}$ ; also, $a(n)$ is the central coefficient of $(1+5*x+4*x^2)^n$ .	5
	1, 5, 33, 245, 1921, 15525, 127905, 1067925, 9004545, 76499525, 653808673, 5614995765, 48416454529, 418895174885, 3634723102113, 31616937184725, 275621102802945, 2407331941640325, 21061836725455905, 184550106298084725	( <a href="#">list</a> ; <a href="#">graph</a> ; <a href="#">refs</a> ; <a href="#">listen</a> ; <a href="#">history</a> ; <a href="#">text</a> ; <a href="#">internal format</a> )
OFFSET	0,2	
COMMENTS	Also number of paths from (0,0) to (n,0) using steps $U=(1,1)$ , $H=(1,0)$ and $D=(1,-1)$ , the $U$ steps come in four colors and the $H$ steps come in five colors. - <a href="#">N.-E. Fahasi</a> , Mar 30 2008 Number of lattice paths from (0,0) to (n,n) using steps (1,0), (0,1), and three kinds of steps (1,1). [Joerg Arndt, Jul 01 2011] Sums of squares of coefficients of $(1+2*x)^n$ . [Joerg Arndt, Jul 06 2011] The Hankel transform of this sequence gives <a href="#">A103488</a> . - <a href="#">Philippe DELEHAM</a> , Dec 02 2007	
REFERENCES	Paul Barry and Aoife Hennessy, Generalized Narayana Polynomials, Riordan Arrays, and Lattice Paths, Journal of Integer Sequences, Vol. 15, 2012, #12.4.8.- From <a href="#">N. J. A. Sloane</a> , Oct 08 2012 Michael Z. Spivey and Laura L. Steil, The $k$ -Binomial Transforms and the Hankel Transform, Journal of Integer Sequences, Vol. 9 (2006), Article 06.1.1.	
LINKS	<a href="#">Table of n, a(n) for n=0..19</a> . Tony D. Noe, <a href="#">On the Divisibility of Generalized Central Trinomial Coefficients</a> , Journal of Integer Sequences, Vol. 9 (2006), Article 06.2.7.	
FORMULA	G.f.: $1/\sqrt{1-10*x+9*x^2}$ . Binomial transform of <a href="#">A059304</a> . G.f.: $\sum_{k \geq 0} \text{binomial}(2*k, k) * (2*x)^k / (1-x)^{(k+1)}$ . E.g.f.: $\exp(5*x)*\text{BesselI}(0, 4*x)$ . - <a href="#">Vladeta Jovovic</a> ( <a href="#">vladeta(AT)eunet.rs</a> ), Aug 20 2003 $a(n) = \sum_{k=0..n} \sum_{j=0..n-k} C(n,j)*C(n-j,k)*C(2*n-2*j,n-j) )$ . - <a href="#">Paul Barry</a> , May 19 2006 $a(n) = \sum_{k=0..n} 4^k * (C(n,k))^2$ [From <a href="#">heruneedollar</a> ( <a href="#">heruneedollar(AT)gmail.com</a> ), Mar 20 2010] Asymptotic: $a(n) \sim 3^{n+1} / (2*\sqrt{2*Pi*n})$ . [ <a href="#">Vaclav Kotesovec</a> , Sep 11 2012] Conjecture: $n*a(n) + 5*(-2*n+1)*a(n-1) + 9*(n-1)*a(n-2) = 0$ . - <a href="#">R. J. Mathar</a> ,	



## Future Work and Takeaways

# Future Work

## Possible Future Projects (inspired by conversations at Texas State with April Yang):

- What if each person tosses several coins simultaneously of different denominations?
- What if some of the coins are positive and some are negative, and have a starting value? Is a +5 and a -3 and a -2 different than a -5 and a +3 and a +2?

## Numerics: Code

```
fibmandmgame[start_, val1_, val2_, val3_, numdo_] := Module[{} ,
  list = {{0, start}};
  current = start;
  Print["Welcome to the Fibonacci M&M Game. You have chosen coin values of ", coin1, ", ", coin2, " and ", coin3, "."];
  Print["We toss all three coins, independently, if coin k comes up heads you get its value."];
  Print["We are doing this for ", numdo, " tosses with starting value ", start, " and will record what happens."];
  Print[
    "Notice the coin values are chosen so that on average there is no change BUT is it more likely to go
      below zero if the big value is the negative versus instead having the two smaller ones be negative?"];
  For[n = 1, n ≤ numdo, n++,
    {
      coin1 = If[Random[] ≤ .5, val1, 0];
      coin2 = If[Random[] ≤ .5, val2, 0];
      coin3 = If[Random[] ≤ .5, val3, 0];
      current = current + (coin1 + coin2 + coin3);
      If[n < 100000, list = AppendTo[list, {n, current}]];
      If[n ≥ 100000 && Mod[n, 1000] == 0, list = AppendTo[list, {n, current}]];
      If[current < 0,
        {
          Print["YOU LOSE! SURVIVED TILL n = ", n, "."];
          n = numdo + 1000;
        }]; (* end of if statement *)
    }]; (* end of n loop *)
  Print[ListLinePlot[list]];
];
```

## Numerics: Simulation

```
fibmandmgame[1000, -5, 3, 2, 200000]
```


Welcome to the Fibonacci M&M Game. You have chosen coin values of 0, 3 and 0.

We toss all three coins, independently, if coin  $k$  comes up heads you get its value.

We are doing this for 200000 tosses with starting value 1000 and will record what happens.

Notice the coin values are chosen so that on average there is no change BUT is it more likely to go below zero if the big value is the negative versus instead having the two smaller ones be negative

YOU LOSE! SURVIVED TILL  $n = 135260$ .



## Lessons

- ◇ Always ask questions.
- ◇ Many ways to solve a problem.
- ◇ Experience is useful and a great guide.
- ◇ Need to look at the data the right way.
- ◇ Often don't know where the math will take you.
- ◇ Value of continuing education: more math is better.
- ◇ Connections: My favorite quote: `If all you have is a hammer, pretty soon every problem looks like a nail.`

## Generating Functions

## Generating Function (Example: Binet's Formula)

### Binet's Formula

$$F_1 = F_2 = 1; F_n = \frac{1}{\sqrt{5}} \left[ \left( \frac{1+\sqrt{5}}{2} \right)^n - \left( \frac{-1+\sqrt{5}}{2} \right)^n \right].$$

- **Recurrence relation:**  $F_{n+1} = F_n + F_{n-1}$  (1)
- **Generating function:**  $g(x) = \sum_{n>0} F_n x^n$ .

$$(1) \Rightarrow \sum_{n \geq 2} F_{n+1} x^{n+1} = \sum_{n \geq 2} F_n x^{n+1} + \sum_{n \geq 2} F_{n-1} x^{n+1}$$

$$\Rightarrow \sum_{n \geq 3} F_n x^n = \sum_{n \geq 2} F_n x^{n+1} + \sum_{n \geq 1} F_n x^{n+2}$$

$$\Rightarrow \sum_{n \geq 3} F_n x^n = x \sum_{n \geq 2} F_n x^n + x^2 \sum_{n \geq 1} F_n x^n$$

$$\Rightarrow g(x) - F_1 x - F_2 x^2 = x(g(x) - F_1 x) + x^2 g(x)$$

$$\Rightarrow g(x) = x/(1 - x - x^2).$$

## Partial Fraction Expansion (Example: Binet's Formula)

- **Generating function:**  $g(x) = \sum_{n>0} F_n x^n = \frac{x}{1-x-x^2}$ .
- **Partial fraction expansion:**

$$\Rightarrow g(x) = \frac{x}{1-x-x^2} = \frac{1}{\sqrt{5}} \left( \frac{\frac{1+\sqrt{5}}{2}x}{1 - \frac{1+\sqrt{5}}{2}x} - \frac{\frac{-1+\sqrt{5}}{2}x}{1 - \frac{-1+\sqrt{5}}{2}x} \right).$$

**Coefficient of  $x^n$  (power series expansion):**

$$F_n = \frac{1}{\sqrt{5}} \left[ \left( \frac{1+\sqrt{5}}{2} \right)^n - \left( \frac{-1+\sqrt{5}}{2} \right)^n \right] \text{ - Binet's Formula!}$$

(using geometric series:  $\frac{1}{1-r} = 1 + r + r^2 + r^3 + \dots$ ).