Linear Recurrences from Schreier Multisets

Y. Geng[1] J. King [2] Z. L. Vasseur [3]
Mentors: H. V. Chu[4], S. J. Miller[5], G.Tresch[6]

[1]geng0114@umn.edu [2]mg168@rice.edu

[3]johnsonj93@lakeforest.edu [4]jhk12@geneseo.edu [5]zachary.l.v@tamu.edu

[6]hchu@wlu.edu [7]sjm1@williams.edu [8]treschgd@tamu.edu

Joint Mathematics Meetings AMS Contributed Paper Session on Combinatorics
Jan 04, 2026
We gratefully acknowledge support from the National Science Foundation
DMS2341670

Schreier sets

A finite set $F \subset \mathbb{N}$ is said to be Schreier if $F = \emptyset$ or $\min F \geq |F|$.

Schreier: $\{2, 3\}, \{4, 5, 10\}$

Not Schreier: $\{3, 7, 10, 13\}$

Schreier sets

A finite set $F \subset \mathbb{N}$ is said to be Schreier if $F = \emptyset$ or $\min F \geq |F|$.

Schreier: $\{2, 3\}, \{4, 5, 10\}$

Not Schreier: $\{3, 7, 10, 13\}$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$$\mathcal{A}_1 = \{\{1\}\} \Longrightarrow |\mathcal{A}_1| = 1$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$$\mathcal{A}_1 = \{\{1\}\} \Longrightarrow |\mathcal{A}_1| = 1$$

$$\mathcal{A}_2 = \{\{2\}\} \Longrightarrow |\mathcal{A}_2| = 1$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$$\mathcal{A}_1 = \{\{1\}\} \Longrightarrow |\mathcal{A}_1| = 1$$

$$\mathcal{A}_2 = \{\{2\}\} \Longrightarrow |\mathcal{A}_2| = 1$$

$$\mathcal{A}_3 = \{\{2, 3\}, \{3\}\} \Longrightarrow |\mathcal{A}_3| = 2$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$$\mathcal{A}_1 = \{\{1\}\} \Longrightarrow |\mathcal{A}_1| = 1$$

$$\mathcal{A}_2 = \{\{2\}\} \Longrightarrow |\mathcal{A}_2| = 1$$

$$\mathcal{A}_3 = \{\{2, 3\}, \{3\}\} \Longrightarrow |\mathcal{A}_3| = 2$$

$$\mathcal{A}_4 = \{\{2, 4\}, \{3, 4\}, \{4\}\} \Longrightarrow |\mathcal{A}_4| = 3$$

$\mathcal{A}_n$

$$\mathcal{A}_n = \{F \subset \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}$$

$$\mathcal{A}_1 = \{\{1\}\} \Longrightarrow |\mathcal{A}_1| = 1$$

$$\mathcal{A}_2 = \{\{2\}\} \Longrightarrow |\mathcal{A}_2| = 1$$

$$\mathcal{A}_3 = \{\{2, 3\}, \{3\}\} \Longrightarrow |\mathcal{A}_3| = 2$$

$$\mathcal{A}_4 = \{\{2, 4\}, \{3, 4\}, \{4\}\} \Longrightarrow |\mathcal{A}_4| = 3$$

$$\mathcal{A}_5 = \{\{2, 5\}, \{3, 5\}, \{4, 5\}, \{3, 4, 5\}, \{5\}\} \Longrightarrow |\mathcal{A}_5| = 5$$

Schreier sets and the Fibonacci sequence

A. Bird (2012)

For $n \in \mathbb{N}$,

$$|\{F \subseteq \{1, 2, \ldots, n\} : n \in F \text{ and } F \text{ is Schreier}\}| = F_n,$$

where $F_1 = F_2 = 1$ and $F_n = F_{n-1} + F_{n-2}$ for $n \geq 3$.

A general problem

For $(s, p, q, n) \in \mathbb{N}^4$,

$$\mathcal{A}_{p,q,n}^{(s_1,\ldots,s_n)} :=$$

$$\{F \subset \{\underbrace{1,\ldots,1}_{s_1},\ldots,\underbrace{n-1,\ldots,n-1}_{s_{n-1}},\underbrace{n,\ldots,n}_{s_n}\} : n \in F \text{ and}$$

$$q \min F \geq p|F|\}$$

A general problem

For $(s, p, q, n) \in \mathbb{N}^4$,

$$\mathcal{A}_{p,q,n}^{(s_1,\ldots,s_n)} := \{F \subset \{\underbrace{1,\ldots,1}_{s_1},\ldots,\underbrace{n-1,\ldots,n-1}_{s_{n-1}},\underbrace{n,\ldots,n}_{s_n}\} : n \in F \text{ and } q \min F \geq p|F|\}$$

What is the sequence $(|\mathcal{A}_{p,q,n}^{(s_1,\ldots,s_n)}|)_{n=1}^{\infty}$?

The special case $q = 1$, $s_1 = \cdots = s_{n-1} = s$, and $s_n = 1$

For $(s, p, n) \in \mathbb{N}^3$,

$$\mathcal{A}_{p,n}^{(s)} :=$$

$$\{F \subset \{\underbrace{1, \ldots, 1}_{s}, \ldots, \underbrace{n-1, \ldots, n-1}_{s}, n\} : n \in F \text{ and } \min F \geq p|F|\}$$

The special case $q = 1$, $s_1 = \cdots = s_{n-1} = s$, and $s_n = 1$

For $(s, p, n) \in \mathbb{N}^3$,

$$\mathcal{A}_{p,n}^{(s)} :=$$

$$\{F \subset \{\underbrace{1, \ldots, 1}_{s}, \ldots, \underbrace{n-1, \ldots, n-1}_{s}, n\} : n \in F \text{ and } \min F \geq p|F|\}$$

$$\{2, 7\}, \{3, 3, 7\} \in \mathcal{A}_{1,7}^{(3)}$$

The special case $q = 1$, $s_1 = \cdots = s_{n-1} = s$, and $s_n = 1$

For $(s, p, n) \in \mathbb{N}^3$,

$$\mathcal{A}_{p,n}^{(s)} :=$$

$$\{F \subset \{\underbrace{1, \ldots, 1}_{s}, \ldots, \underbrace{n-1, \ldots, n-1}_{s}, n\} : n \in F \text{ and } \min F \geq p|F|\}$$

$$\{2, 7\}, \{3, 3, 7\} \in \mathcal{A}_{1,7}^{(3)}$$

$$\{2, 2\} \notin \mathcal{A}_{2,5}^{(4)}$$

The special case $q = 1$, $s_1 = \cdots = s_{n-1} = s$, and $s_n = 1$

For $(s, p, n) \in \mathbb{N}^3$,

$$\mathcal{A}_{p,n}^{(s)} :=$$
$$\{F \subset \{\underbrace{1, \ldots, 1}_{s}, \ldots, \underbrace{n-1, \ldots, n-1}_{s}, n\} : n \in F \text{ and } \min F \geq p|F|\}$$

$\{2, 7\}, \{3, 3, 7\} \in \mathcal{A}_{1,7}^{(3)}$

$\{2, 2\} \notin \mathcal{A}_{2,5}^{(4)}$

$\{3, 3, 6, 7\} \notin \mathcal{A}_{1,7}^{(3)}$

Count uncolored Schreier multisets

For $(s, p) \in \mathbb{N}^2$, what is the sequence $(|\mathcal{A}_{p,n}^{(s)}|)_{n=1}^{\infty}$?

Data

$|\mathcal{A}^{(2)}_{1,n}|$: $1, 1, 2, 4, 7, 13, 24, 44, 81, 149, 274, 504, 927, \ldots$

$a_n = a_{n-1} + a_{n-2} + a_{n-3}$

$|\mathcal{A}^{(2)}_{2,n}|$: $0, 1, 1, 1, 2, 3, 5, 8, 12, 19, 30, 47, 74, 116, \ldots$

$a_n = a_{n-1} + a_{n-3} + a_{n-5}$

$|\mathcal{A}^{(3)}_{1,n}|$: $1, 1, 2, 4, 8, 15, 29, 56, 108, 208, 401, 773, 1490, \ldots$

$a_n = a_{n-1} + a_{n-2} + a_{n-3} + a_{n-4}$

$|\mathcal{A}^{(3)}_{2,n}|$: $0, 1, 1, 1, 2, 3, 5, 8, 13, 21, 33, 53, 85, \ldots$

$a_n = a_{n-1} + a_{n-3} + a_{n-5} + a_{n-7}$

Data

$|\mathcal{A}_{1,n}^{(2)}|$: $1, 1, 2, 4, 7, 13, 24, 44, 81, 149, 274, 504, 927, \ldots$

$a_n = a_{n-1} + a_{n-2} + a_{n-3}$

$|\mathcal{A}_{2,n}^{(2)}|$: $0, 1, 1, 1, 2, 3, 5, 8, 12, 19, 30, 47, 74, 116, \ldots$

$a_n = a_{n-1} + a_{n-3} + a_{n-5}$

$|\mathcal{A}_{1,n}^{(3)}|$: $1, 1, 2, 4, 8, 15, 29, 56, 108, 208, 401, 773, 1490, \ldots$

$a_n = a_{n-1} + a_{n-2} + a_{n-3} + a_{n-4}$

$|\mathcal{A}_{2,n}^{(3)}|$: $0, 1, 1, 1, 2, 3, 5, 8, 13, 21, 33, 53, 85, \ldots$

$a_n = a_{n-1} + a_{n-3} + a_{n-5} + a_{n-7}$

Number of terms: $\boxed{s+1}$; **gap btwn consecutive indices:** $\boxed{p}$

Theorem

Polymath Jr. 25 Schreier Group

For $(s, p) \in \mathbb{N}^2$ and $n \geq sp + 2$,

$$|\mathcal{A}_{p,n}^{(s)}| = |\mathcal{A}_{p,n-1}^{(s)}| + |\mathcal{A}_{p,n-1-p}^{(s)}| + |\mathcal{A}_{p,n-1-2p}^{(s)}| + \cdots + |\mathcal{A}_{p,n-1-sp}^{(s)}|.$$

Proof

$$I_s(n) := \{\underbrace{1,\ldots,1}_{s},\ldots,\underbrace{n-1,\ldots,n-1}_{s},n\}$$

$$c(F,k) := \text{number of copies of } k \text{ in } F$$

Proof

$$I_s(n) := \{\underbrace{1,\ldots,1}_{s},\ldots,\underbrace{n-1,\ldots,n-1}_{s},n\}$$

$$c(F,k) := \text{number of copies of } k \text{ in } F$$

$$\begin{aligned}
&\mathcal{A}_{p,n}^{(s)}\\
=\ &\{F\subset I_s(n): n\in F, c(F,n-1)=0, \min F\geq p|F|\}\cup\\
&\{F\subset I_s(n): n\in F, c(F,n-1)=1, \min F\geq p|F|\}\cup\\
&\{F\subset I_s(n): n\in F, c(F,n-1)=2, \min F\geq p|F|\}\cup\\
&\ldots\\
&\{F\subset I_s(n): n\in F, c(F,n-1)=s, \min F\geq p|F|\}
\end{aligned}$$

Proof

$$\mathcal{A}_{p,n}^{(s)} = \bigcup_{k=0}^{s} \{F \subset I_s(n) : n \in F, c(F, n-1) = k, \min F \geq p|F|\}$$

Proof

$$\mathcal{A}_{p,n}^{(s)} = \bigcup_{k=0}^{s} \{F \subset I_s(n) : n \in F, c(F, n-1) = k, \min F \geq p|F|\}$$

$$\mathcal{A}_{p,n-1-kp}^{(s)} \to \{F \subset I_s(n) : n \in F, c(F, n-1) = k, \min F \geq p|F|\}$$

$$F \mapsto \begin{cases} (F \backslash \{n-1\}) \cup \{n\}, & \text{if } k = 0; \\ (F + kp) \cup \{\underbrace{n-1, \ldots, n-1}_{k-1}, n\}, & \text{if } k \geq 1. \end{cases}$$

Proof

$$\mathcal{A}_{p,n}^{(s)} = \bigcup_{k=0}^{s} \{F \subset I_s(n) : n \in F, c(F, n-1) = k, \min F \geq p|F|\}$$

$$\mathcal{A}_{p,n-1-kp}^{(s)} \to \{F \subset I_s(n) : n \in F, c(F, n-1) = k, \min F \geq p|F|\}$$

$$F \mapsto \begin{cases} (F \backslash \{n-1\}) \cup \{n\}, & \text{if } k = 0; \\ (F + kp) \cup \{\underbrace{n-1, \ldots, n-1}_{k-1}, n\}, & \text{if } k \geq 1. \end{cases}$$

$$|\mathcal{A}_{p,n}^{(s)}| = |\mathcal{A}_{p,n-1}^{(s)}| + |\mathcal{A}_{p,n-1-p}^{(s)}| + |\mathcal{A}_{p,n-1-2p}^{(s)}| + \cdots + |\mathcal{A}_{p,n-1-sp}^{(s)}|$$

Colored Schreier multisets

Let k_i denote the integer k with color i.

$$B_n^{(s)} = \{1_1, 1_2, \ldots, 1_s, \ldots, (n-1)_1, \ldots, (n-1)_s, n\}$$

$$\mathcal{B}_{p,n}^{(s)} := \{F \subset B_n^{(s)} : n \in F \text{ and } \min F \geq p|F|\}.$$

Colored Schreier multisets

Let k_i denote the integer k with color i.

$$B_n^{(s)} = \{1_1, 1_2, \ldots, 1_s, \ldots, (n-1)_1, \ldots, (n-1)_s, n\}$$

$$\mathcal{B}_{p,n}^{(s)} := \{F \subset B_n^{(s)} : n \in F \text{ and } \min F \geq p|F|\}.$$

$$\{2_1, 3_2\} \neq \{2_2, 3_2\}$$

Colored Schreier multisets

Let k_i denote the integer k with color i.

$$B_n^{(s)} = \{1_1, 1_2, \ldots, 1_s, \ldots, (n-1)_1, \ldots, (n-1)_s, n\}$$

$$\mathcal{B}_{p,n}^{(s)} := \{F \subset B_n^{(s)} : n \in F \text{ and } \min F \geq p|F|\}.$$

$$\{2_1, 3_2\} \neq \{2_2, 3_2\}$$

For $(s, p) \in \mathbb{N}^2$, what is the sequence $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^{\infty}$?

Data

$\mathcal{B}^{(2)}_{1,n}$: $1, 1, 3, 6, 13, 28, 60, 129, 277, 595, 1278, 2745, 5896, \ldots$

$a_n = a_{n-1} + 2a_{n-2} + a_{n-3}$

$\mathcal{B}^{(2)}_{2,n}$: $0, 1, 1, 1, 3, 5, 8, 15, 26, 45, 80, 140, 245, 431, \ldots$

$a_n = a_{n-1} + 2a_{n-3} + a_{n-5}$

$\mathcal{B}^{(3)}_{1,n}$: $1, 1, 4, 10, 26, 69, 181, 476, 1252, 3292, 8657, 22765, 59864, \ldots$

$a_n = a_{n-1} + 3a_{n-2} + 3a_{n-3} + a_{n-4}$

$\mathcal{B}^{(3)}_{2,n}$: $0, 1, 1, 1, 4, 7, 13, 28, 53, 105, 211, 413, 819, 1624, 3206, \ldots$

$a_n = a_{n-1} + 3a_{n-3} + 3a_{n-5} + a_{n-7}$

Data

$\mathcal{B}_{1,n}^{(2)}$: $1, 1, 3, 6, 13, 28, 60, 129, 277, 595, 1278, 2745, 5896, \ldots$

$a_n = a_{n-1} + 2a_{n-2} + a_{n-3}$

$\mathcal{B}_{2,n}^{(2)}$: $0, 1, 1, 1, 3, 5, 8, 15, 26, 45, 80, 140, 245, 431, \ldots$

$a_n = a_{n-1} + 2a_{n-3} + a_{n-5}$

$\mathcal{B}_{1,n}^{(3)}$: $1, 1, 4, 10, 26, 69, 181, 476, 1252, 3292, 8657, 22765, 59864, \ldots$

$a_n = a_{n-1} + 3a_{n-2} + 3a_{n-3} + a_{n-4}$

$\mathcal{B}_{2,n}^{(3)}$: $0, 1, 1, 1, 4, 7, 13, 28, 53, 105, 211, 413, 819, 1624, 3206, \ldots$

$a_n = a_{n-1} + 3a_{n-3} + 3a_{n-5} + a_{n-7}$

Coefficients: $\boxed{\binom{s}{i}}, 0 \leq i \leq s$; **gap btwn consecutive indices:** $\boxed{p}$

Theorem

Polymath Jr. 25 Schreier Group

For $(s, p) \in \mathbb{N}^2$ and $n \geq sp + 2$,

$$|\mathcal{B}_{p,n}^{(s)}| = \binom{s}{0}|\mathcal{B}_{p,n-1}^{(s)}| + \binom{s}{1}|\mathcal{B}_{p,n-1-p}^{(s)}| + \cdots + \binom{s}{s}|\mathcal{B}_{p,n-1-sp}^{(s)}|.$$

Proof

Linear recurrences are encoded by polynomials.

Proof

Linear recurrences are encoded by polynomials.

Step 1: Find $(b^s_{p,n})_{n=1}^{\infty}$ such that $(|\mathcal{B}^s_{p,n}|)_{n=1}^{\infty}$ is an s-periodic subsequence of $(b^s_{p,n})_{n=1}^{\infty}$

$(b^s_{p,n})_{n=1}^{\infty}$ satisfies the polynomial $p(x)$

Proof

Linear recurrences are encoded by polynomials.

Step 1: Find $(b^s_{p,n})_{n=1}^{\infty}$ such that $(|\mathcal{B}^s_{p,n}|)_{n=1}^{\infty}$ is an s-periodic subsequence of $(b^s_{p,n})_{n=1}^{\infty}$

$(b^s_{p,n})_{n=1}^{\infty}$ satisfies the polynomial $p(x)$

Step 2: To prove $(|\mathcal{B}^s_{p,n}|)_{n=1}^{\infty}$ satisfies a recurrence encoded by $q(x)$, it suffices to verify $p(x)$ divides $q(x^s)$

Sample case: $\mathcal{B}_{1,n}^{(2)}$

Step 1: Find $b_{1,n}^{(2)}$

$\mathcal{B}_{1,n}^{(2)}$: 1, 1, 3, 6, 13, 28, 60, 129, 277, 595, 1278, 2745, 5896, . . .

$b_{1,n}^{(2)}$: 1, 1, 1, 2, 3, 4, 6, 9, 13, 19, 28, 41, 60, 88, 129, 189,
277, 406, 595, . . .

Sample case: $\mathcal{B}^{(2)}_{1,n}$

Step 1: Find $b^{(2)}_{1,n}$

$\mathcal{B}^{(2)}_{1,n}$: $1, 1, 3, 6, 13, 28, 60, 129, 277, 595, 1278, 2745, 5896, \ldots$

$b^{(2)}_{1,n}$: $1, 1, 1, 2, 3, 4, 6, 9, 13, 19, 28, 41, 60, 88, 129, 189,$
$277, 406, 595, \ldots$

$b^{(2)}_{1,1} = b^{(2)}_{1,2} = b^{(2)}_{1,3} = 1$ and $b^{(2)}_{1,n} = b^{(2)}_{1,n-1} + b^{(2)}_{1,n-3}, n \geq 4$

Sample case: $\mathcal{B}_{1,n}^{(2)}$

Step 1: Find $b_{1,n}^{(2)}$

$\mathcal{B}_{1,n}^{(2)}$: $1, 1, 3, 6, 13, 28, 60, 129, 277, 595, 1278, 2745, 5896, \ldots$

$b_{1,n}^{(2)}$: $1, 1, 1, 2, 3, 4, 6, 9, 13, 19, 28, 41, 60, 88, 129, 189,$
$277, 406, 595, \ldots$

$b_{1,1}^{(2)} = b_{1,2}^{(2)} = b_{1,3}^{(2)} = 1$ and $b_{1,n}^{(2)} = b_{1,n-1}^{(2)} + b_{1,n-3}^{(2)}, n \geq 4$

$(b_{1,n}^{(2)})_{n=1}^{\infty}$ satisfies $p(x) = 1 - x - x^3$.

Sample case: $\mathcal{B}^{(2)}_{1,n}$

Step 2: To prove $(|\mathcal{B}^2_{1,n}|)_{n=1}^{\infty}$ satisfies a recurrence encoded by $q(x) = 1 - x - 2x^2 - x^3$, it suffices to verify $p(x)$ divides $q(x^2)$.

Sample case: $\mathcal{B}_{1,n}^{(2)}$

Step 2: To prove $(|\mathcal{B}_{1,n}^{2}|)_{n=1}^{\infty}$ satisfies a recurrence encoded by $q(x) = 1 - x - 2x^2 - x^3$, it suffices to verify $p(x)$ divides $q(x^2)$.

$$q(x^2) \;=\; 1 - x^2 - 2x^4 - x^6 \;=\; \underbrace{(1 - x - x^3)}_{p(x)}(1 + x + x^3)$$

Schreier sets without multiples of u

For $u \geq 2$, $g_u(n)$ denotes the n^{th} positive integer not divisible by u.

Schreier sets without multiples of u

For $u \geq 2$, $g_u(n)$ denotes the n^{th} positive integer not divisible by u.

$G_{u,n} := \{g_u(1), g_u(2), ..., g_u(n)\}$and define

$$\mathcal{D}_{u,n} := \left\{ F \subset G_{u,n} : \left\lfloor \frac{un-1}{u-1} \right\rfloor \in F \text{ and } F \text{ is Schreier} \right\}$$

Schreier sets without multiples of u

Polymath Jr. 25 Schreier Group

For $1 \leq n \leq 2u-1$, we have $|\mathcal{D}_{u,n}| = F_n$. For $n \geq 2u$,

$$|\mathcal{D}_{u,n}| = \sum_{i=1}^{u} (-1)^{i-1} \binom{u}{i} |\mathcal{D}_{u,n-i}| + |\mathcal{D}_{u,n-2u+1}|.$$

Schreier sets without multiples of u

$$|\mathcal{D}_{u,n}| = \sum_{k=1}^{\left\lfloor \frac{(n+1)u-1}{2u-1} \right\rfloor} \binom{n-k+\left\lfloor \frac{k-1}{u} \right\rfloor}{k-1}$$

Schreier sets without multiples of u

Step 1: Find $(d_{u,n})_{n=1}^{\infty}$ such that $(|\mathcal{D}_{u,n}|)_{n=1}^{\infty}$ is a periodic subsequence of $(d_{u,n})_{n=1}^{\infty}$

Schreier sets without multiples of u

Step 1: Find $(d_{u,n})_{n=1}^{\infty}$ such that $(|\mathcal{D}_{u,n}|)_{n=1}^{\infty}$ is a periodic subsequence of $(d_{u,n})_{n=1}^{\infty}$

$$d_{u,1} = d_{u,2} = \cdots = d_{u,2u-1} = 1$$

$$d_{u,n} = d_{u,n-u} + d_{u,n-2u+1} \text{ for } n \geq 2u$$

Schreier sets without multiples of u

Step 1: Find $(d_{u,n})_{n=1}^{\infty}$ such that $(|\mathcal{D}_{u,n}|)_{n=1}^{\infty}$ is a periodic subsequence of $(d_{u,n})_{n=1}^{\infty}$

$$d_{u,1} = d_{u,2} = \cdots = d_{u,2u-1} = 1$$

$$d_{u,n} = d_{u,n-u} + d_{u,n-2u+1} \text{ for } n \geq 2u$$

$(d_{u,n})_{n=1}^{\infty}$ satisfies

$$p_u(x) := 1 - x^u - x^{2u-1}$$

Schreier sets without multiples of u

For each $u \geq 2$, $(|\mathcal{D}_{u,n}|)_{n=1}^{\infty}$ is a u-periodic subsequence of $(d_{u,n})_{n=1}^{\infty}$:

$$|\mathcal{D}_{u,n}| \;=\; d_{u,un-(u-1)}, \text{ for all } n \in \mathbb{N}.$$

Schreier sets without multiples of u

Step 2:

To prove that $(|\mathcal{D}_{u,n}|)_{n=1}^{\infty}$ satisfies a linear reccurence encoded by $q_u(x) = (1-x)^u - x^{2u-1}$, it suffices to verify that $p_u(x) := 1 - x^u - x^{2u-1}$ divides $q_u(x^u)$.

Future investigations

Allow different elements to have different number of copies

Future investigations

Allow different elements to have different number of copies

Study the more general condition $q \min F \geq p|F|$

Future investigations

Allow different elements to have different number of copies

Study the more general condition $q \min F \geq p|F|$

Study nonlinear Schreier-type conditions for multisets:
$(\min F)^q \geq |F|^p$

Paper

Link to the paper:

https://arxiv.org/abs/2509.05158