

PRIMES IN TUPLES II

D. A. GOLDSTON, J. PINTZ AND C. Y. YILDIRIM

ABSTRACT. We prove that

$$\liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\sqrt{\log p_n} (\log \log p_n)^2} < \infty,$$

where p_n denotes the n^{th} prime. Since on average $p_{n+1} - p_n$ is asymptotically $\log p_n$, this shows that we can always find pairs of primes much closer together than the average. We actually prove a more general result concerning the set of values taken on by the differences $p - p'$ between primes which includes the small gap result above.

1. INTRODUCTION

In the first paper in this series [7] we proved that, letting p_n denote the n^{th} prime,

$$(1.1) \quad \Delta = \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} = 0,$$

culminating 80 years of work on this problem. Since the average spacing $p_{n+1} - p_n$ in the sequence of primes is asymptotically $\log p_n$, this result showed for the first time that the prime numbers do not eventually become isolated from each other in the sense that there will always be pairs of primes closer than any fraction of the average spacing. For the history of this problem, we refer the reader to [7] and [19].

The information about primes used to obtain (1.1) is contained in the Bombieri-Vinogradov theorem. Let

$$(1.2) \quad \theta(N; q, a) = \sum_{\substack{n \leq N \\ n \equiv a \pmod{q}}} \theta(n), \quad \text{where } \theta(n) = \begin{cases} \log n, & \text{if } n \text{ is prime,} \\ 0, & \text{otherwise.} \end{cases}$$

The Bombieri-Vinogradov theorem states that for any $A > 0$ there is a $B = B(A)$ such that, for $Q = N^{\frac{1}{2}}(\log N)^{-B}$,

$$(1.3) \quad \sum_{q \leq Q} \max_{\substack{a \\ (a, q) = 1}} \left| \theta(N; q, a) - \frac{N}{\phi(q)} \right| \ll \frac{N}{(\log N)^A}.$$

Thus the primes tend to be equally distributed among the arithmetic progressions modulo q that allow primes, and this holds for the primes up to N and at least for almost all the progressions with modulus q up to nearly $N^{\frac{1}{2}}$. The principle can be quantified by saying that the primes have an *admissible level of distribution* ϑ (or *satisfy a level of distribution* ϑ) if (1.3) holds for any $A > 0$ and any $\epsilon > 0$ with

$$(1.4) \quad Q = N^{\vartheta - \epsilon}.$$

Elliott and Halberstam [3] conjectured that the primes have the maximal admissible level of distribution 1, while by the Bombieri-Vinogradov theorem we have immediately that $1/2$ is an admissible level of distribution for the primes. In [7]

Date: October 15, 2007.

The first author was supported in part by an NSF Grant, the second author by OTKA grants No. K 67676, T 43623, T 49693 and the Balaton program, the third author by TÜBITAK.

we proved that *if* the primes satisfy a level of distribution $\vartheta > \frac{1}{2}$ then there is an absolute constant $M(\vartheta)$ for which

$$(1.5) \quad p_{n+1} - p_n \leq M(\vartheta), \quad \text{for infinitely many } n.$$

In particular assuming the Elliott-Halberstam conjecture (or just $\vartheta \geq 0.98$) then

$$(1.6) \quad p_{n+1} - p_n \leq 16, \quad \text{for infinitely many } n.$$

These are surprising results because they show that going beyond an admissible level of distribution $1/2$ implies there are infinitely often bounded gaps between primes, and therefore questions as hard as the twin prime conjecture can nearly be dealt with using this type of information.

Since we obtained our results in 2005 there has been no further progress toward (1.5), and it appears now that an extension of the Bombieri-Vinogradov theorem of sufficient strength to obtain bounded gaps between primes will require some basic new ideas. One can also pursue improving the approximations we used or improving the method used to detect primes, and again this now appears to require some essentially new idea. Our goal in this paper is to extend the current method as much as possible in order to obtain strong quantitative results. In particular we obtain the following quantitative version of (1.1).

Theorem 1. *The differences of consecutive primes satisfy*

$$(1.7) \quad \liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\sqrt{\log p_n (\log \log p_n)^2}} < \infty.$$

This result is remarkable in that it shows that there exist pairs of primes nearly within the square root of the average spacing. By comparison, the best result for large gaps between primes [18] is that

$$(1.8) \quad \limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{(\log p_n)(\log \log p_n)(\log \log \log p_n)^{-2}(\log \log \log \log p_n)} \geq 2e^\gamma,$$

where γ is Euler's constant. Thus the best large gap result produces gaps larger than the average by a factor a bit smaller than $\log \log p_n$, while now the small gaps are smaller than the average by a factor a bit bigger than $(\log p_n)^{-\frac{1}{2}}$. In this sense the small gap result now greatly surpasses the large gap result. (There are conflicting conjectures on how large the gap between consecutive primes can get, but all of these conjectures suggest that there can be gaps at least as large as $c(\log p_n)^2$ for some constant c .)

This paper is organized as follows. In Section 2 we present a generalization of Theorem 1 which applies to many interesting situations and which is the result we will prove in this paper. Unlike in [7], to obtain our results we need to take into account the possibility of exceptional characters associated with Landau-Siegel zeros. In Theorem 2 we assume that there are no Landau-Siegel zeros in a certain range and are able to obtain our main results including the gaps between primes in Theorem 1 in intervals $[N, 2N]$ for all sufficiently large N . Next, using the Landau-Page Theorem we can find a sequence of ranges which avoid possible Landau-Siegel zeros. Thus we obtain Theorem 3 which unconditionally gives the same results as Theorem 2 but without being able to localize them to a dyadic interval. The proof of these theorems requires substantial refinements of the methods of [7], and in Section 3 we will discuss some of these refinements and how they arise. The main technical tools needed in our proof, Theorems 4 and 5, are stated in Section 4. The proof of Theorems 4 and 5 take up Sections 5 through 13. In proving Theorem 5 in our general setting we need a modified Bombieri-Vinogradov theorem which is the topic of Section 12. Our method, as in [7], requires a result on the average of the singular series. In [7] the well-known result of Gallagher [5] was used, but

in our current setting this result is not applicable, and therefore in Section 14 we prove a new result well adapted for our needs. With Theorems 4 and 5 in hand together with the new singular series average result, the proof of Theorems 2 and 3 is completed in Section 15.

Notation. In the following c and C will denote (sufficiently) small and (sufficiently) large absolute positive constants, respectively, which have been chosen appropriately. This is also true for constants formed from c or C with subscripts or accents. We will allow these constants to be different at different occurrences. Constants implied by pure o , O , $\ll$ symbols will be absolute, unless otherwise stated. The ν times iterated logarithm will be denoted by $\log_\nu N$. $\mathcal{P}$ denotes the set of primes.

2. A GENERALIZATION OF THEOREM 1

Our method will allow us to prove a generalization of Theorem 1 where instead of seeking two neighboring primes of the form $n + i$, $n + j$ with

$$(2.1) \quad 1 \leq i < j \leq h, \quad h = h(n) = C\sqrt{\log n}(\log \log n)^2,$$

we look for two primes of the form $n + a_i$, $n + a_j$ where (2.1) is satisfied and

$$(2.2) \quad \mathcal{A} = \{a_i\}_{i=1}^h \subset [1, n]$$

is an arbitrarily given set of integers.

We remark that an extension of this type is a trivial consequence of the prime number theorem if

$$(2.3) \quad h' = h'(n) > (1 + c) \log n, \quad c > 0 \text{ fixed},$$

but that none of the earlier methods of Erdős [4], Bombieri–Davenport [1] and Maier [14] which produce small gaps between primes seem capable of proving a result of this type for any function satisfying

$$(2.4) \quad h'' = h''(n) < (1 - c) \log n, \quad c > 0 \text{ fixed}.$$

According to a conjecture of de Polignac [20] from 1849, every even number may be written as the difference of two primes. Although we know that this is true for almost all even numbers, there is no known way to specify these values. Our generalization makes a first step in this direction by proving that we can explicitly find sparse sequences $\mathcal{A}$ such that infinitely many of the elements

$$(2.5) \quad \mathcal{A} - \mathcal{A}$$

are differences of two primes, i.e.

$$(2.6) \quad |(\mathcal{P} - \mathcal{P}) \cap (\mathcal{A} - \mathcal{A})| = \infty.$$

(Here we make use of the usual notation that for sets $\mathcal{A}$ and $\mathcal{B}$, $\mathcal{A} - \mathcal{B} = \{a - b : a \in \mathcal{A}, b \in \mathcal{B}\}$.) Some sequences for which our method applies are:

$$(2.7) \quad \mathcal{A} = \{k^m\}_{m=1}^\infty, \quad k \geq 2 \text{ fixed } (k \in \mathbb{N}),$$

$$(2.8) \quad \mathcal{A} = \{k^{x^2+y^2}\}_{x,y=1}^\infty, \quad k \geq 2 \text{ fixed } (k \in \mathbb{N}),$$

$$(2.9) \quad \mathcal{A} = \{k^{f(x,y)}\}_{r=1}^\infty, \quad k \geq 2 \text{ fixed } (k \in \mathbb{N}),$$

where the value set $\mathcal{R} = \{r \in \mathbb{N}; \exists x, y : f(x, y) = r\}$ satisfies

$$(2.10) \quad \mathcal{R}(X) = |\{m; m \leq X, m \in \mathcal{R}\}| > C' \sqrt{X} \log^2 X$$

(this would happen e.g. for $f(x, y) = x^2 + y^m$ with arbitrary $m \geq 2$, and for $f(x, y) = x^3 + y^3$), or in general any set of type

$$(2.11) \quad \mathcal{A} = \{k^{r_j}\}_{j=1}^\infty, \quad k \geq 2 \text{ fixed } (k \in \mathbb{N}),$$

if $\mathcal{R} = \{r_j\}_{j=1}^\infty \subseteq \mathbb{N}$ satisfies the density condition (2.10). Among these sets the only trivial one is 2^m , that is (2.7) for $k = 2$ which has $\lfloor \log X / \log 2 \rfloor$ elements below X , thereby corresponding to the case (2.3).

Unfortunately, the possible existence of Landau-Siegel zeros (cf. Section 12) makes it impossible to formulate a localized version of our result for all $n \in \mathbb{N}$ satisfying the conditions (2.1) and (2.2). Even in the special case when $\mathcal{A}$ is an interval we cannot guarantee the existence of gaps of size $O(\sqrt{\log N} \log_2^2 N)$ between primes in any interval of type $[N, 2N]$ for any large N .

The first formulation of our main result assumes there are no exceptional characters in a certain range. (This is Hypothesis $S(Y)$ from Section 12, with $Y = Y(N) = \exp(3\sqrt{\log N})$.)

Theorem 2. *Let us suppose that an $N > N_0$ is given such that for any real primitive character $\chi \pmod{q}$, $q \leq \exp(3\sqrt{\log N})$ we have*

$$(2.12) \quad L(s, \chi) \neq 0 \quad \text{for } s \in \left(1 - \frac{1}{9\sqrt{\log N}}, 1\right].$$

Let $\mathcal{A} = \mathcal{A}_N = \{a_i\}_{i=1}^h \subseteq [1, N] \cap \mathbb{N}$ be arbitrary ($a_i \neq a_j$) with

$$(2.13) \quad h \geq C\sqrt{\log N} \log_2^2 N,$$

where C is an appropriate absolute constant. Then there exists $n \in [N, 2N]$ such that at least two numbers of the form

$$(2.14) \quad n + a_i, \quad n + a_j, \quad (1 \leq i < j \leq h),$$

are primes.

In Section 12 we show that (2.12), i.e. Conjecture $S(Y(N))$, is true for an infinite sequence $N = N_\nu \rightarrow \infty$, and thus Theorem 2 implies Theorem 1 by choosing $N = N_\nu$ and

$$(2.15) \quad \mathcal{A} = \mathcal{A}_N = \{1, 2, \dots, h\}$$

with $h = \lceil C\sqrt{\log N} \log_2^2 N \rceil$. A more general formulation of this result which covers the special cases mentioned in (2.7)–(2.9) and (2.11) as well as Theorem 1 can be stated as follows.

Theorem 3. *Let $\mathcal{A} \subseteq \mathbb{N}$ be an arbitrary sequence satisfying*

$$(2.16) \quad \mathcal{A}(N) = |\{n; n \leq N, n \in \mathcal{A}\}| > C\sqrt{\log N} \log_2^2 N \quad \text{for } N > N_0.$$

Then infinitely many elements of $\mathcal{A} - \mathcal{A}$ can be written as the difference of two primes, that is,

$$(2.17) \quad |(\mathcal{P} - \mathcal{P}) \cap (\mathcal{A} - \mathcal{A})| = \infty.$$

3. SOME INITIAL CONSIDERATIONS

The main tool of our method is an approximation for prime tuples and almost prime tuples. Consider the tuple $(n + h_1, n + h_2, \dots, n + h_K)$ as n runs over the integers. If these K values are all primes for some n then we call this a prime tuple, and we wish to examine the existence of prime tuples. A first consideration is that the set of shifts

$$(3.1) \quad \mathcal{H} = \{h_1, h_2, \dots, h_K\}, \quad \text{with } h_i \neq h_j \text{ (if } i \neq j),$$

imposes divisibility conditions on the components of the tuple which can effect the likelihood of obtaining prime tuples or even preclude the possibility of more than a single prime tuple. Specifically, let $\nu_p(\mathcal{H})$ denote the number of distinct residue classes modulo p occupied by the elements of $\mathcal{H}$, and for squarefree integers d extend

this definition to $\nu_d(\mathcal{H})$ multiplicatively. The singular series for the set $\mathcal{H}$ is defined to be

$$(3.2) \quad \mathfrak{S}(\mathcal{H}) = \prod_p \left(1 - \frac{1}{p}\right)^{-K} \left(1 - \frac{\nu_p(\mathcal{H})}{p}\right).$$

If $\mathfrak{S}(\mathcal{H}) \neq 0$ then $\mathcal{H}$ is called *admissible*. Thus $\mathcal{H}$ is admissible if and only if $\nu_p(\mathcal{H}) < p$ for all p , while if $\nu_p(\mathcal{H}) = p$ then one component of the tuple is always divisible by p and there can be at most one prime tuple of this form. Hardy and Littlewood [9] conjectured an asymptotic formula for the number of prime tuples $(n + h_1, n + h_2, \dots, n + h_K)$, with $1 \leq n \leq N$, as $N \rightarrow \infty$. Letting

$$(3.3) \quad \theta(n) = \begin{cases} \log n, & \text{if } n \text{ is prime,} \\ 0, & \text{otherwise;} \end{cases}$$

we define

$$(3.4) \quad \Lambda(n; \mathcal{H}) := \theta(n + h_1)\theta(n + h_2) \cdots \theta(n + h_K)$$

and use this function to detect prime tuples. The Hardy–Littlewood prime-tuple conjecture is the asymptotic formula

$$(3.5) \quad \sum_{n \leq N} \Lambda(n; \mathcal{H}) = N(\mathfrak{S}(\mathcal{H}) + o(1)), \quad \text{as } N \rightarrow \infty,$$

which is trivial if $\mathcal{H}$ is not admissible, but is otherwise only known to be true in the case $K = 1$ which is the prime number theorem.

The starting point for our method in [7] is to find approximations of $\Lambda(n; \mathcal{H})$ for which we can obtain asymptotic formulas similar to (3.5). A further essential idea is that rather than approximating just prime tuples, we should approximate almost-prime K -tuples with a total of $\leq K + \ell$ prime factors in all the components, which if $0 \leq \ell \leq K - 2$ guarantees at least two of the components are prime. The almost prime tuple approximation used in [7] and which we also use here is

$$(3.6) \quad \Lambda_R(n; \mathcal{H}, \ell) := \frac{1}{(K + \ell)!} \sum_{\substack{d | P_{\mathcal{H}}(n) \\ d \leq R}} \mu(d) \left(\log \frac{R}{d} \right)^{K + \ell},$$

where $|\mathcal{H}| = K$, and

$$(3.7) \quad \mathcal{P}_{\mathcal{H}}(n) := (n + h_1)(n + h_2) \cdots (n + h_K).$$

Our method for proving (1.1) in [7] is based on a comparison of the two sums

$$(3.8) \quad \sum_{n \leq N} \Lambda_R(n; \mathcal{H}, \ell)^2 \quad \text{and} \quad \sum_{n \leq N} \theta(n + h_0) \Lambda_R(n; \mathcal{H}, \ell)^2.$$

An asymptotic formula for the first sum can be obtained if $R \leq N^{1/2 - \epsilon}$, while for the second sum we can use an admissible level of distribution of primes ϑ to obtain an asymptotic formula when $R \leq N^{\vartheta/2 - \epsilon}$. In [7] it was assumed that K and ℓ are fixed, i.e. independent of N . Using these asymptotic formulas we can now evaluate

$$(3.9) \quad \mathcal{S}_R := \sum_{n=N+1}^{2N} \left(\sum_{1 \leq h_0 \leq h} \theta(n + h_0) - \log 3N \right) \sum_{\substack{1 \leq h_1, h_2, \dots, h_K \leq h \\ \text{distinct}}} \Lambda_R(n; \mathcal{H}, \ell)^2,$$

If $\mathcal{S}_R > 0$ then the sum over h_0 must have at least two non-zero terms and thus there must be some n and $h_i \neq h_j$ such that $n + h_i$ and $n + h_j$ are both prime. We find with $\vartheta = 1/2$ and $h = \lambda \log N$ with any fixed $\lambda > 0$ that we can choose K and

ℓ for which $\mathcal{S}_R > 0$, which proves (1.1). In order to obtain this for any arbitrarily small $\lambda > 0$, the fixed K and ℓ are chosen sufficiently large in an appropriate way.

To obtain quantitative bounds to replace (1.1), the first step is to obtain asymptotic formulas which are uniform in K and ℓ so that these can be chosen as functions of N that go to infinity with N . One also needs explicit error terms, and these error terms arise not only from lower order terms and prime number theorem type error terms, but also in (3.8) from the Bombieri-Vinogradov theorem error terms.

We will now establish the relations between our parameters that will be used throughout the paper. Recalling the set $\mathcal{A}$ from (2.2), we will always take $\mathcal{H} \subset \mathcal{A}$. Next, R and ℓ will be chosen as

$$(3.10) \quad K \leq h, \quad \ell \asymp \sqrt{K}, \quad R := (3N)^\Theta = (3N)^{1/4-\xi}, \quad \xi = o(1).$$

We will make use of two important parameters U and V defined by

$$(3.11) \quad V := \sqrt{\log N}, \quad U = e^V,$$

and will choose K later to be slightly smaller than V . We next denote the product of primes not exceeding V by

$$(3.12) \quad P := \prod_{p \leq V} p,$$

where p will always denote primes.

As just mentioned above, our present treatment requires a much more delicate analysis of the error terms than in [7], and therefore we make an initial simplification to facilitate this analysis. In [7] the irregular behavior of $\nu_p(\mathcal{H})$ for small primes greatly complicated the estimate of the function $G(s_1, s_2)$ and its partial derivatives. We can avoid these difficulties, at least for primes dividing P , by proceeding somewhat similarly to Heath-Brown in [11]. We call a residue class $a \pmod{P}$ *regular with respect to $\mathcal{H}$ and P* if

$$(3.13) \quad (P, P_{\mathcal{H}}(a)) = 1$$

and denote by $A(\mathcal{H}) = A_P(\mathcal{H})$ the set of all regular residue classes mod P . Thus

$$(3.14) \quad A(\mathcal{H}) := \{a; 1 \leq a \leq P; (P, P_{\mathcal{H}}(a)) = 1\}.$$

The number of regular residue classes mod P is clearly

$$(3.15) \quad |A(\mathcal{H})| = \prod_{p|P} (p - \nu_p(\mathcal{H}))$$

and their proportion of all the residue classes mod P is

$$(3.16) \quad \frac{|A(\mathcal{H})|}{P} = \prod_{p|P} \left(1 - \frac{\nu_p(\mathcal{H})}{p}\right),$$

which is positive if $\mathcal{H}$ is admissible. Thus in particular for a given $\mathcal{H}$ and all P there exists at least one regular residue class mod P , if and only if $\mathcal{H}$ is admissible.

With this notation, we now consider the sums

$$(3.17) \quad \sum_{\substack{n=N+1 \\ n \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2)}}^{2N} \Lambda_R(n; \mathcal{H}_1, \ell) \Lambda_R(n; \mathcal{H}_2, \ell)$$

and

$$(3.18) \quad \sum_{\substack{n=N+1 \\ n \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2)}}^{2N} \Lambda_R(n; \mathcal{H}_1, \ell) \Lambda_R(n; \mathcal{H}_2, \ell) \theta(n + h_0)$$

with $h_0 \in [1, h]$, which are asymptotically evaluated in Theorems 4 and 5, respectively. A new feature in the proof of these theorems which does not occur in [7] is that (3.17) and (3.18) are first evaluated for each residue class $a(\bmod P)$ with

$$(3.19) \quad a \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2) = A(\mathcal{H}_1 \cup \mathcal{H}_2)$$

separately, and then the results are added over all regular residue classes modulo P . It turns out that the asymptotic main term (and even secondary terms) are independent of the particular choice of the regular residue class a , so this summing presents no difficulty. However, the restriction of the values of n to a single residue class $a(\bmod P)$ in (3.18) requires a stronger form of the Bombieri–Vinogradov theorem (cf. Section 12).

To detect primes, in place of (3.9) we consider

$$(3.20) \quad S'_R(N, K, \ell, P) := \frac{1}{Nh^{2K+1}} \sum_{n=N+1}^{2N} \left(\sum_{\substack{p \\ p-n \in \mathcal{A}}} \log p - \log 3N \right) (\Psi'_R(K, \ell, n, h))^2,$$

where

$$(3.21) \quad \Psi'_R(K, \ell, n, h) := \sum_{\substack{\mathcal{H}, |\mathcal{H}|=K \\ n \in A(\mathcal{H})}} \Lambda_R(n; \mathcal{H}, \ell).$$

On applying Theorems 4 and 5 we can asymptotically evaluate S'_R , which we carry out in Section 15. One condition that arises from the main terms is that in order to prove the existence of prime pairs in intervals of length h we need

$$(3.22) \quad h > \frac{C \log N}{K}.$$

Since $K \leq h$ this immediately implies that

$$(3.23) \quad h > C \sqrt{\log N}.$$

Our goal is to take h as small as possible, and therefore we can not obtain anything better than (3.23) when using the approximation in (3.6) together with (3.20). Apart from powers of $\log_2 N$, we are able to prove our results for h of this size.

Our actual choices for K and h are

$$(3.24) \quad K \leq c_1 \frac{\sqrt{\log N}}{\log_2^2 N}, \quad h = \frac{25 \log N}{K} \geq \frac{25}{c_1} \sqrt{\log N} \log_2^2 N,$$

with a sufficiently small explicitly calculable absolute constant c_1 (to be chosen later). We will need the error terms in Theorems 4 and 5 to be uniform in K with a relative error of size η_1 satisfying

$$(3.25) \quad \eta_1 < \frac{c}{\sqrt{K}}.$$

However, we do not achieve this for all admissible pairs $\mathcal{H}_1$ and $\mathcal{H}_2$ of size K . Instead, for all admissible pairs $\mathcal{H}_1, \mathcal{H}_2$ we obtain a weaker error term, but if

$$(3.26) \quad K - |\mathcal{H}_1 \cap \mathcal{H}_2| \ll \sqrt{K}.$$

then we do obtain the error estimate in (3.25). This turns out to be sufficient for our proof, since such pairs $\mathcal{H}_1, \mathcal{H}_2$ will be dominant in (3.20).

4. TWO BASIC THEOREMS

In the following let N be a sufficiently large integer, c_1 a sufficiently small positive constant,

$$(4.1) \quad K \leq c_1 \frac{\sqrt{\log N}}{(\log_2 N)^2},$$

$$(4.2) \quad K \ll k_1, k_2 \leq K, \quad \sqrt{K} \ll \ell_1, \ell_2 \ll \sqrt{K}.$$

We will consider sets $\mathcal{H} := \mathcal{H}_1 \cup \mathcal{H}_2$, $\mathcal{H}_1, \mathcal{H}_2 \subseteq [1, N]$ of sizes

$$(4.3) \quad |\mathcal{H}_i| = k_i, \quad |\mathcal{H}_1 \cap \mathcal{H}_2| = r.$$

Let

$$(4.4) \quad \bar{m} := K - m \text{ for } m \in [0, K], \quad n^* := \max(\sqrt{K}, n), \quad \bar{n}^* := (\bar{n})^*.$$

Our first main result is the following theorem.

Theorem 4. *We have for $N^c < R \leq N^{1/2} \exp(-c\sqrt{\log N})$, as $N \rightarrow \infty$*

$$(4.5) \quad \sum_{\substack{n \leq N \\ n \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2)}} \Lambda_R(n; \mathcal{H}_1, \ell_1) \Lambda_R(n; \mathcal{H}_2, \ell_2) = \\ = N \binom{\ell_1 + \ell_2}{\ell_1} \frac{(\log R)^{r+\ell_1+\ell_2}}{(r+\ell_1+\ell_2)!} \frac{\mathfrak{S}(\mathcal{H})P}{|A(\mathcal{H})|} \left(1 + O\left(\frac{K\bar{r}^* \log_2 N}{\log R} \right) \right) + O\left(N e^{-c\sqrt{\log N}} \right).$$

For the next theorem we suppose that the following form of the Bombieri–Vinogradov theorem holds (see Section 12). For a given, sufficiently large N , and recalling the parameter P defined in (3.12), we have

$$(4.6) \quad \sum_{\substack{q \leq Q^* \\ (q, P)=1}} \max_{\substack{(a, q)=1 \\ (a, q)=1}} \left| \sum_{\substack{N < p \leq 2N \\ p \equiv a \pmod{Pq}}} \log p - \frac{N}{\varphi(Pq)} \right| \ll \frac{N}{P} \exp(-c\sqrt{\log N}),$$

where

$$(4.7) \quad Q^* = N^{1/2} P^{-3} \exp(-c^* \sqrt{\log N}),$$

with an arbitrary positive constant c^* .

Letting $\mathcal{H}^0 = \mathcal{H} \cup \{h_0\}$, our second main result is as follows.

Theorem 5. *Suppose (4.6)–(4.7) hold and let $N^c \leq R \leq \sqrt{Q^*}$. Then*

$$(4.8) \quad \sum_{\substack{N < n \leq 2N \\ n \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2)}} \Lambda_R(n; \mathcal{H}_1, \ell_1) \Lambda_R(n; \mathcal{H}_2, \ell_2) \theta(n + h_0) \\ = N \frac{C_R(\ell_1, \ell_2, \mathcal{H}_1, \mathcal{H}_2, h_0)}{(r + \ell_1 + \ell_2)!} \binom{\ell_1 + \ell_2}{\ell_1} \mathfrak{S}(\mathcal{H}^0) (\log R)^{r+\ell_1+\ell_2} \left(1 + O\left(\frac{CK\bar{r}^* \log_2 N}{\log R} \right) \right) \\ + O\left(N e^{-c\sqrt{\log N}} \right),$$

where

$$(4.9) \quad C_R(\ell_1, \ell_2, \mathcal{H}_1, \mathcal{H}_2, h_0) = \begin{cases} 1, & \text{if } h_0 \notin \mathcal{H}; \\ \frac{(\ell_1 + \ell_2 + 1) \log R}{(\ell_1 + 1)(r + \ell_1 + \ell_2 + 1)}, & \text{if } h_0 \in \mathcal{H}_1 \text{ and } h_0 \notin \mathcal{H}_2; \\ \frac{(\ell_1 + \ell_2 + 2)(\ell_1 + \ell_2 + 1) \log R}{(\ell_1 + 1)(\ell_2 + 1)(r + \ell_1 + \ell_2 + 1)}, & \text{if } h_0 \in \mathcal{H}_1 \cap \mathcal{H}_2. \end{cases}$$

For the applications to Theorems 1–3 the simpler case $\ell_1 = \ell_2 = \ell$ will be sufficient.

5. LEMMAS

We will use standard properties of the Riemann zeta function $\zeta(s)$. Proceeding slightly differently from [7] we use the zero-free region, with $s = \sigma + it$,

$$(5.1) \quad \zeta(1+s) \neq 0 \quad \text{for } s \in \mathcal{R}_N := \left\{ s; \sigma \geq -\frac{1}{\log_2 N + 6 \log(|t|+3)} \right\}.$$

Further we have for $s \in \mathcal{R}_N$ by Titchmarsh [23, Ch. 3]

$$(5.2) \quad \max \left(\left| \zeta(1+s) - \frac{1}{s} \right|, \left| \frac{1}{\zeta(1+s)} \right|, \left| \frac{\zeta'}{\zeta}(1+s) + \frac{1}{s} \right| \right) \ll \log(|t|+3).$$

In the course of the proof the following contours which lie in the zero-free region $\mathcal{R}_N$ will be used (with U and V given in (3.11))

$$(5.3) \quad \begin{aligned} \mathcal{L}_1 &:= \left\{ \sigma = \frac{1}{28V}, |t| \leq U \right\}, & \mathcal{L}_2 &:= \left\{ \sigma = \frac{1}{14V}, |t| \leq 2U \right\}, \\ \mathcal{L}_3 &:= \left\{ \sigma = \frac{-1}{28V}, |t| \leq U \right\}, & \mathcal{L}_4 &:= \left\{ \sigma = \frac{-1}{14V}, |t| \leq 2U \right\}, \\ \mathcal{L}_5 &:= \left\{ -\frac{1}{28V} \leq \sigma \leq \frac{1}{28V}, |t| = U \right\}, & \mathcal{L}_6 &:= \left\{ -\frac{1}{14V} \leq \sigma \leq \frac{1}{14V}, |t| = 2U \right\}, \end{aligned}$$

$$(5.4) \quad \begin{aligned} \mathcal{L}' &= \mathcal{L}'_0 \cup \mathcal{L}'_1; \quad \mathcal{L}'_0 = \left\{ s; s = \delta_0 e^{i\varphi}, \frac{\pi}{2} \leq \varphi \leq \frac{3\pi}{2} \right\}, \\ \mathcal{L}'_1 &= \{s = it; \delta_0 \leq |t| \leq U\}, \quad \delta_0 = \left(\sqrt{K} \log_2 N \right)^{-1}. \end{aligned}$$

Similarly to Lemma 1 of [7], we have

Lemma 1. *Let $k(\log_2 N)^2 \leq \sqrt{\log R}$, $N^c \leq R \leq N$, $N \geq C$, $k \geq 2$, $B \leq Ck$. Then*

$$(5.5) \quad \int_{\mathcal{L}_i} (\log(|t|+3))^B \left| \frac{R^s ds}{s^k} \right| \ll e^{-c\sqrt{\log N}}, \quad (3 \leq i \leq 6),$$

where the constant implied by the $\ll$ symbol depends only on C .

Proof. The integral I in (5.5) satisfies for all i

$$(5.6) \quad \begin{aligned} I &\ll \int_0^{2U} R^{-1/(28V)} \frac{(\log(|t|+3))^B}{\max(|t|, \frac{1}{28V})^k} dt + \int_{|\sigma| \leq 1/14V} R^\sigma \cdot \frac{d\sigma}{U^{3/2}} \\ &\ll e^{-c\sqrt{\log N}} \left(\int_0^C (28V)^k dt + \int_C^\infty \frac{dt}{t^{3/2}} \right) + e^{\sqrt{\log N}(1/14-1/2)} \ll e^{-c\sqrt{\log N}}. \end{aligned}$$

We will prove a generalization of the combinatorial identity (8.16) of [7] in order to evaluate the terms $I_{1,1}$ of Section 8. Let us define for triplets of integers d, u, y with $d \geq 0$, $u \geq 0$, $y+u \geq 0$ (to be called suitable triplets) the quantity

$$(5.7) \quad Z(d, u, y) := \frac{1}{u!} \sum_{\substack{m=0 \\ m \geq -y}}^u \binom{u}{m} (-1)^m \frac{d(d+1) \cdots (d+m-1)}{(y+m)!}.$$

Lemma 2. *We have for any suitable triplet d, u, y the relation*

$$(5.8) \quad Z(d, u, y) = \frac{(y-d+1) \cdots (y-d+u)}{u!(y+u)!}.$$

Proof. We will prove this by induction on u . For $u = 0$ we have trivially for any non-negative d and y , $Z(d, 0, y) = (y!)^{-1}$ (the empty product in the numerator of Z is 1 by definition). We can suppose $u \geq 1$ and that our statement is true for all suitable triplets $d, u-1, y$. Making the convention that we define for $n < 0$

$$(5.9) \quad \frac{x}{n!} = 0$$

for any real number x (in other words, we just neglect in a sum all terms with an $n!$ in the denominator with $n < 0$), we obtain by $\binom{u}{i} = \binom{u-1}{i} + \binom{u-1}{i-1}$ (where we define $\binom{u-1}{-1} = \binom{u-1}{-1} = 0$), with the notation $[S] = 1$ if the statement S is true and $[S] = 0$ if S is false,

$$\begin{aligned} Z(d, u, y) &= \frac{1}{u!} \left\{ \sum_{\substack{i=0 \\ i \geq -y}}^{u-1} \binom{u-1}{i} (-1)^i \frac{d(d+1) \dots (d+i-1)}{(y+i)!} \right. \\ &\quad \left. - \sum_{\substack{j=0 \\ j \geq -y-1}}^{u-1} \binom{u-1}{j} (-1)^j \frac{d(d+1) \dots (d+j-1)}{(y+j+1)!} \right\} \\ &= \frac{1}{u!} \left\{ \sum_{\substack{i=0 \\ i \geq -y}}^{u-1} \binom{u-1}{i} (-1)^i \frac{d(d+1) \dots (d+i-1)}{(y+i)!} \left(1 - \frac{d+i}{y+i+1} \right) \right. \\ &\quad \left. - [-y-1 \geq 0] \binom{u-1}{-y-1} (-1)^{-y-1} d(d+1) \dots (d-y-2)(d-y-1) \right\} \\ &= \frac{1}{u} \cdot \frac{1}{(u-1)!} \sum_{\substack{i=0 \\ i \geq -y-1}}^{u-1} \binom{u-1}{i} (-1)^i \frac{d(d+1) \dots (d+i-1)(y+1-d)}{(y+i+1)!} \\ &= \frac{y+1-d}{u} Z(d, u-1, y+1) = \frac{(y+1-d)(y+2-d) \dots (y+u-d)}{u!(y+u)!}. \end{aligned}$$

Finally we mention a simple lemma for the mean value of the generalized divisor function

$$(5.10) \quad d_m(q) := m^{\omega(q)},$$

where $\omega(q)$ denotes the number of prime-factors of q for a squarefree q .

Lemma 3. *If $m > 0$, $\nu \geq \max(c' \log(K+1), 1)$ then there exists a constant C' depending on c' such that, for $K \geq 1$ and $x \geq 1$ we have*

$$(5.11) \quad \sum_{q \leq x}^b d_m(q) \leq x(1 + \log x)^{\lceil m \rceil}$$

and

$$(5.12) \quad \sum_{q \leq x}^b \frac{(d_{3K}(q))^{1+1/\nu}}{q} \leq (1 + \log x)^{C'K}.$$

Proof. Equation (5.11) follows from

$$(5.13) \quad \sum_{q \leq x}^b d_m(q) \leq x \left(\sum_{q \leq x}^b \frac{d_{\lceil m \rceil}(q)}{q} \right) \leq x \left(\sum_{j \leq x} \frac{1}{j} \right)^{\lceil m \rceil} \leq x(1 + \log x)^{\lceil m \rceil}.$$

Further, by (5.10) we have

$$(5.14) \quad (d_{3K}(q))^{1+1/\nu} = d_j(q)$$

with

$$(5.15) \quad j = (3K)^{1+1/\nu} \leq 9e^{1/c'} K,$$

and Lemma 3 follows with $C' = 9e^{1/c'} + 1$.

6. PREPARATION FOR THE PROOF OF THEOREM 4

Since the preparation for the proof of Theorem 4 is nearly the same as in Sections 6 and 7 of [7] for the analogous Proposition 1 (or 3 or 4), we will briefly summarize it and the reader is referred for the details to [7]. Let

$$(6.1) \quad \mathcal{H}(p) = \left\{ h'_1, \dots, h'_{\nu_p(\mathcal{H})} : h'_j \equiv h_i \pmod{p}, h_i \in \mathcal{H} \text{ for some } i, 1 \leq h'_j \leq p \right\},$$

$$(6.2) \quad \bar{\nu}_p(\mathcal{H}_1 \bar{\cap} \mathcal{H}_2) := \nu_p(\mathcal{H}_1(p) \cap \mathcal{H}_2(p)) = \nu_p(\mathcal{H}_1) + \nu_p(\mathcal{H}_2) - \nu_p(\mathcal{H}).$$

For any $a \in A(\mathcal{H}_1) \cap A(\mathcal{H}_2)$ (cf. (3.14)), we have similarly to Section 7 of [7]

$$(6.3) \quad \begin{aligned} S_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, a) &:= \sum_{\substack{n=N+1 \\ n \equiv a \pmod{P}}}^{2N} \Lambda_R(n; \mathcal{H}_1, \ell_1) \Lambda_R(n; \mathcal{H}_2, \ell_2) \\ &= \frac{N}{P} \mathcal{T}_R(\ell_1, \ell_2; \mathcal{H}_1, \mathcal{H}_2) + O(R^2 (3 \log R)^{7K}), \end{aligned}$$

where

$$(6.4) \quad \mathcal{T}_R(\ell_1, \ell_2; \mathcal{H}_1, \mathcal{H}_2) := \frac{1}{(2\pi i)^2} \int \int_{(1)(1)} F(s_1, s_2) \frac{R^{s_1}}{s_1^{K+\ell_1+1}} \frac{R^{s_2}}{s_2^{K+\ell_2+1}} ds_1 ds_2,$$

$$(6.5) \quad F(s_1, s_2) := \prod_{p>V} \left(1 - \frac{\nu_p(\mathcal{H}_1)}{p^{1+s_1}} - \frac{\nu_p(\mathcal{H}_2)}{p^{1+s_2}} + \frac{\bar{\nu}_p(\mathcal{H}_1 \bar{\cap} \mathcal{H}_2)}{p^{1+s_1+s_2}} \right),$$

where now, differently from [7], primes not exceeding V do not appear in $F(s_1, s_2)$ since by the regularity of a , $(P_{\mathcal{H}_1}(n), P) = (P_{\mathcal{H}_2}(n), P) = 1$.

Let

$$(6.6) \quad \Delta := \left| \prod_{1 \leq i < j \leq K} (h_i - h_j) \right| \leq N^{K(K-1)/2}.$$

Then if $p \nmid \Delta$ (consequently, for all sufficiently large primes p),

$$(6.7) \quad \nu_p(\mathcal{H}_1) = |\mathcal{H}_1| = K, \quad \nu_p(\mathcal{H}_2) = |\mathcal{H}_2| = K, \quad \bar{\nu}_p(\mathcal{H}_1 \bar{\cap} \mathcal{H}_2) = |\mathcal{H}_1 \cap \mathcal{H}_2| = r.$$

We therefore factor out the dominant zeta-factors and write

$$(6.8) \quad F(s_1, s_2) = G_{\mathcal{H}_1, \mathcal{H}_2}(s_1, s_2) \frac{\zeta(1+s_1+s_2)^d}{\zeta(1+s_1)^a \zeta(1+s_2)^b}$$

with a function $G(s_1, s_2)$, regular for $\sigma_i > -1/5$, say, which we write slightly more generally for future application in Theorem 5 as

$$(6.9) \quad G_{\mathcal{H}_1, \mathcal{H}_2}(s_1, s_2) = G(s_1, s_2) = G = G_1 G_2 G_3 = G_1 G_4,$$

where now $a = b = K$, $d = r$, $\nu_1(p) = \nu_p(\mathcal{H}_1)$, $\nu_2(p) = \nu_p(\mathcal{H}_2)$, $\nu_3(p) = \bar{\nu}_p(\mathcal{H}_1 \bar{\cap} \mathcal{H}_2)$,

$$(6.10) \quad G_1(s_1, s_2) = \prod_{p \leq V} \left(1 - \frac{1}{p^{1+s_1}} \right)^{-a} \prod_{p \leq V} \left(1 - \frac{1}{p^{1+s_2}} \right)^{-b} \prod_{p \leq V} \left(1 - \frac{1}{p^{1+s_1+s_2}} \right)^d,$$

$$(6.11) \quad G_4(s_1, s_2) = \prod_{p>V} \left(\frac{\left(1 - \frac{\nu_1(p)}{p^{1+s_1}} - \frac{\nu_2(p)}{p^{1+s_2}} + \frac{\nu_3(p)}{p^{1+s_1+s_2}} \right) \left(1 - \frac{1}{p^{1+s_1+s_2}} \right)^d}{\left(1 - \frac{1}{p^{1+s_1}} \right)^a \left(1 - \frac{1}{p^{1+s_2}} \right)^b} \right)$$

$$= \prod_{p|\Delta, p>V} \cdot \prod_{p \nmid \Delta, p>V} =: G_2(s_1, s_2) G_3(s_1, s_2).$$

Let us use the notation

$$(6.12) \quad \delta_i := \max(0, -\sigma_i), \quad \delta := \delta_1 + \delta_2, \quad s_3 := s_1 + s_2,$$

and

$$(6.13) \quad \mathcal{R}'_N := \left\{ s; \sigma \geq -\frac{1/2}{\log_2 N + 6 \log(|t| + 3)} \right\}.$$

We will estimate the order of $G(s_1, s_2)$ in the region $s_1, s_2 \in \mathcal{R}'_N$ under the more general conditions

$$(6.14) \quad a, b, d \leq K, \quad \nu_i(p) \leq K,$$

$$(6.15) \quad \nu_1(p) = a, \quad \nu_2(p) = b, \quad \nu_3(p) = d \text{ for } p \nmid \Delta.$$

(Later we will examine more delicate properties of $G(s_1, s_2)$ with further conditions on $a, b, d, \nu_i(p)$.) We have

$$(6.16) \quad |G_1(s_1, s_2)| \leq \exp \left(C \sum_{p \leq V} \frac{K}{p^{1-\delta}} \right) \leq \exp(CK \log_3 N),$$

$$(6.17) \quad |G_2(s_1, s_2)| \leq \exp \left(C \sum_{p|\Delta} \frac{K}{p^{1-\delta}} \right) \leq \exp \left(CK \sum_{p \leq \log \Delta(1+o(1))} \frac{1}{p^{1-\delta}} \right) \\ \leq \exp(CK \log_3 N),$$

and

$$(6.18) \quad |G_3(s_1, s_2)| \leq \exp \left(C \sum_{p>V} \frac{K^2}{p^{2-2\delta}} \right) \leq \exp \left(\frac{CK^2}{V} \right) \leq \exp(CK),$$

where in (6.16)–(6.18) we made use of the estimates

$$(6.19) \quad \max(V^\delta, (\log \Delta)^\delta) \leq (\log^2 N)^{\frac{1/2}{\log_2 N}} = e;$$

further in (6.17) the sum which was originally over $p \mid \Delta$ has been majorized by using the set of the smallest possible primes which could divide Δ .

Summarizing (6.16)–(6.18) we obtain

$$(6.20) \quad |G(s_1, s_2)| \leq e^{CK \log_3 N} \quad \text{for } s_1, s_2 \in \mathcal{R}'_n,$$

and further

$$(6.21) \quad |F(s_1, s_2)| \leq e^{CK \log_3 N} ((\log(|t_1| + 3)) \log(|t_2| + 3))^{2K} \text{ for } s_1, s_2, s_3 \in \mathcal{R}'_n.$$

The above estimate shows that the integrand in (6.4) vanishes as either $|t_1| \rightarrow \infty$ or $|t_2| \rightarrow \infty$, $s_1, s_2, s_3 \in \mathcal{R}'_N$. We will examine the integral (analogously to (7.15) in [7])

$$(6.22) \quad I := \mathcal{T}_R^*(d, a, b, u, v, \mathcal{H}_1, \mathcal{H}_2) := \frac{1}{(2\pi i)^2} \int \int_{(1)(1)} \frac{D(s_1, s_2) R^{s_1+s_2} ds_1 ds_2}{s_1^{u+1} s_2^{v+1} (s_1 + s_2)^d},$$

where we introduce the function $D(s_1, s_2)$, regular for $s_1, s_2, s_3 \in \mathcal{R}'_N$,

$$(6.23) \quad D(s_1, s_2) := D_0(s_1, s_2) G(s_1, s_2), \quad D_0(s_1, s_2) := \frac{W^d(s_1 + s_2)}{W^a(s_1) W^b(s_2)}, \quad W(s) := s\zeta(1+s),$$

$$(6.24) \quad 0 \leq d \leq a, \quad b \leq K, \quad \min(a, b) \geq cK, \quad \sqrt{K}/8 \leq u, \quad v \leq \sqrt{K}$$

and by symmetry we can assume $u \leq v$. (In the applications we will have $|a-b| \leq 1$, $|u-v| \leq 1$.)

First step. Move the contour (1) for the integral over s_1 to $\mathcal{L}_1$, over s_2 to $\mathcal{L}_2$. The vertical parts $|t| \geq U$, and $|t| \geq 2U$, resp. can be neglected similarly to Lemma 1. After this move the integral in s_1 from $\mathcal{L}_1$ to $\mathcal{L}_3 \cup \mathcal{L}_5$. The horizontal segments $\mathcal{L}_5$ can be again neglected. We pass a pole of order $u+1$ at $s_1 = 0$, and obtain

$$(6.25) \quad I = I_1 + \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_2} \int_{\mathcal{L}_3} \frac{D(s_1, s_2) R^{s_1+s_2} ds_1 ds_2}{s_1^{u+1} s_2^{v+1} (s_1 + s_2)^d} = I_1 + I_2 + O(e^{-c\sqrt{\log N}}),$$

where

$$(6.26) \quad \begin{aligned} I_1 &:= \frac{1}{2\pi i} \int_{\mathcal{L}_2} \operatorname{Res}_{s_1=0} \left(\frac{D(s_1, s_2) R^{s_1+s_2}}{s_1^{u+1} s_2^{v+1} (s_1 + s_2)^d} \right) ds_2 \\ &= \frac{1}{2\pi i} \int_{\mathcal{L}_2} \frac{1}{u!} \left\{ \sum_{i=0}^u \binom{u}{i} (\log R)^{u-i} \frac{\partial^i}{\partial s_1^i} \left(\frac{D(s_1, s_2)}{(s_1 + s_2)^d} \right) \right\} \Big|_{s_1=0} \frac{R^{s_2}}{s_2^{v+1}} ds_2. \end{aligned}$$

We denote the complete integrand above by $Z(s_2)$ and express

$$(6.27) \quad \begin{aligned} \frac{\partial^i}{\partial s_1^i} \left(\frac{D(s_1, s_2)}{(s_1 + s_2)^d} \right) \Big|_{s_1=0} &= (-1)^i \frac{D(0, s_2) d(d+1) \dots (d+i-1)}{s_2^{d+i}} \\ &+ \sum_{j=1}^i \binom{i}{j} \frac{\partial^j}{\partial s_1^j} D(s_1, s_2) \Big|_{s_1=0} \cdot (-1)^{i-j} \frac{d(d+1) \dots (d+i-j-1)}{s_2^{d+i-j}} \end{aligned}$$

where in case of $i = j$ (including also the case when $i = j = 0$ and $d \geq 0$ arbitrary) the empty product in the numerator is 1.

Second step. Let us denote the contribution of the first term in (6.27) to (6.26) by $I_1(i, 0)$ and the others by $I_1(i, j)$ ($1 \leq j \leq i$). $I_1(i, 0)$ will belong to the main term, all $I_1(i, j)$ with $j \geq 1$ will just contribute to the secondary terms. Let us move now the contour $\mathcal{L}_2$ for the integral over s_2 to $\mathcal{L}_4 \cup \mathcal{L}_6$ in (6.26). The horizontal segments $\mathcal{L}_6$ can be neglected again. We pass a pole of order $v+1+d+i-j$ in case of $I_1(i, j)$ and we obtain in this way

$$(6.28) \quad \begin{aligned} I_1 &= \frac{1}{u!} \sum_{i=0}^u \binom{u}{i} (\log R)^{u-i} \sum_{j=0}^i (-1)^{i-j} \binom{i}{j} \frac{d(d+1) \dots (d+i-j-1)}{(v+d+i-j)!} \times \\ &\times \sum_{\nu=0}^{v+d+i-j} \binom{v+d+i-j}{\nu} (\log R)^{v+d+i-j-\nu} \cdot \frac{\partial^\nu}{\partial s_2^\nu} \frac{\partial^j}{\partial s_1^j} D(s_1, s_2) \Big|_{s_1=s_2=0} \\ &+ \frac{1}{2\pi i} \int_{\mathcal{L}_4} Z(s_2) ds_2 + O(e^{-c\sqrt{\log N}}) =: I_{1,1} + I_{1,2} + O(e^{-c\sqrt{\log N}}). \end{aligned}$$

7. ESTIMATES OF THE PARTIAL DERIVATIVES OF $D(s_1, s_2)$

In this section we will estimate partial derivatives $\frac{\partial^i}{\partial s_1^i} \frac{\partial^j}{\partial s_2^j} D(s_1, s_2)$ of $D(s_1, s_2)$ for $i+j \leq CK$ with $s_i = s_i^*$ in $\mathcal{R}'_N$ for $1 \leq i \leq 3$. We will often use Cauchy's estimate for functions regular in $|z - z_0| \leq \eta$:

$$(7.1) \quad \frac{1}{j!} |f^{(j)}(z_0)| \leq \eta^{-j} \max_{|z-z_0|=\eta} |f(z)|.$$

Applying this for $D(s_1, s_2)$ we obtain

$$(7.2) \quad \frac{1}{i!j!} \left| \frac{\partial^i}{\partial s_1^i} \frac{\partial^j}{\partial s_2^j} D(s_1^*, s_2^*) \right| \ll \eta^{-(i+j)} \max_{|s'_1 - s_1^*| \leq \eta, |s'_2 - s_2^*| \leq \eta} |D(s'_1, s'_2)|.$$

In order to substitute the above maximum for $D(s_1^*, s_2^*)$, we have to estimate

$$(7.3) \quad L(s_1, s_2) := \max \left(\left| \frac{\partial}{\partial s_1} \log D(s_1, s_2) \right|, \left| \frac{\partial}{\partial s_2} \log D(s_1, s_2) \right| \right),$$

for $s_1, s_2, s_3 \in \mathcal{R}'_N$; since by the regularity of $\log D(s_1, s_2)$ for $s_i \in \mathcal{R}'_N$ ($1 \leq i \leq 3$) (cf. (6.10), (6.11), (6.23)) we have, for $\eta \leq (\log_2 N + \log(|t_1| + 3) + \log(|t_2| + 3))^{-1}/100$,

$$(7.4) \quad \left| \frac{D(s'_1, s'_2)}{D(s_1^*, s_2^*)} \right| \leq \exp(2\eta \cdot \max_{|s_1 - s_1^*| \leq \eta, |s_2 - s_2^*| \leq \eta} L(s_1, s_2)).$$

By symmetry it is enough to deal with

$$(7.5) \quad L_1(s_1, s_2) := \left| \frac{\partial}{\partial s_1} \log D(s_1, s_2) \right|.$$

Since the logarithm is an additive function, using the representation (6.9)–(6.11) and (6.23) of $D(s_1, s_2)$ it is sufficient to examine the factors D_0, G_1, G_2, G_3 separately.

We will choose a positive η

$$(7.6) \quad \eta \leq \frac{1/100}{\log_2 N + \log T}, \quad T = T_1 + T_2, \quad T_i = |t_i| + 3, \quad t_3 = t_1 + t_2 \quad (1 \leq i \leq 3)$$

where by $\delta = \delta_1 + \delta_2 \leq 2/\log_2 N$, $V = (\log N)^{1/2}$, $\log \Delta \leq K^2 \log N \leq \log^2 N$ we have

$$(7.7) \quad \max(V^\delta, V^\eta, (\log \Delta)^\delta, (\log \Delta)^\eta) \ll 1.$$

We have by (5.2) and (6.23)

$$(7.8) \quad \begin{aligned} \frac{\partial}{\partial s_1} (\log D_0(s_1, s_2)) &= d \cdot \left(\frac{\zeta'}{\zeta} (1 + s_1 + s_2) + \frac{1}{s_1 + s_2} \right) - a \left(\frac{\zeta'}{\zeta} (1 + s_1) + \frac{1}{s_1} \right) \\ &\ll K \log T. \end{aligned}$$

Further we have

$$(7.9) \quad \frac{\partial}{\partial s_1} (\log G_1(s_1, s_2)) = \sum_{p \leq V} \frac{\log p}{p^{1+s_1}} \left(\frac{dp^{-s_2}}{1 - p^{-(1+s_1+s_2)}} - \frac{a}{1 - p^{-(1+s_1)}} \right) \ll K \log_2 N.$$

Similarly to (6.17) we obtain by (7.7)

$$(7.10) \quad \begin{aligned} &\frac{\partial}{\partial s_1} (\log G_2(s_1, s_2)) \\ &= \sum_{\substack{p|\Delta \\ p > V}} \frac{\log p}{p^{1+s_1}} \left\{ \frac{\nu_1(p) - p^{-s_2} \nu_3(p)}{1 - \frac{\nu_1(p)}{p^{1+s_1}} - \frac{\nu_2(p)}{p^{1+s_2}} + \frac{\nu_3(p)}{p^{1+s_1+s_2}}} - \frac{a}{1 - \frac{1}{p^{1+s_1}}} + \frac{dp^{-s_2}}{1 - \frac{1}{p^{1+s_1+s_2}}} \right\} \\ &\ll K \sum_{p|\Delta} \frac{\log p}{p^{1-\delta}} \ll K \sum_{p \leq \log \Delta(1+o(1))} \frac{\log p}{p^{1-\delta}} \ll K \log_2 \Delta \ll K \log_2 N. \end{aligned}$$

Finally, analogously to (6.18) we have by (6.15)

$$\begin{aligned}
(7.11) \quad & \frac{\partial}{\partial s_1}(\log G_3(s_1, s_2)) \\
&= \sum_{\substack{p \nmid \Delta \\ p > V}} \frac{\log p}{p^{1+s_1}} \left\{ \frac{a - dp^{-s_2}}{1 - \frac{a}{p^{1+s_1}} - \frac{b}{p^{1+s_2}} + \frac{d}{p^{1+s_1+s_2}}} - \frac{a}{1 - \frac{1}{p^{1+s_1}}} + \frac{dp^{-s_2}}{1 - \frac{1}{p^{1+s_1+s_2}}} \right\} \\
&\ll \sum_{p > V} \frac{\log p}{p^{1-\delta_1}} \left(a \cdot \frac{K}{p^{1-\delta}} + dp^{\delta_2} \cdot \frac{K}{p^{1-\delta}} \right) \ll K^2 \sum_{p > V} \frac{\log p}{p^{2-2\delta}} \ll \frac{K^2}{V^{1-2\delta}} \ll \frac{K^2}{V} \ll K.
\end{aligned}$$

Summarizing (7.3)–(7.11) we have

$$(7.12) \quad \max_{|s'_1 - s_1^*| \leq \eta, |s'_2 - s_2^*| \leq \eta} |D(s'_1, s'_2)| \leq e^{C\eta K(\log_2 N + \log T)} |D(s_1^*, s_2^*)| \text{ if } s_1^*, s_2^*, s_3^* \in \mathcal{R}'_N.$$

Hence, (7.2) and (7.12) imply by the choice $\eta^{-1} = 100K(\log_2 N + \log T)$ the following estimate.

Lemma 4. *We have for $s_1, s_2, s_3 \in \mathcal{R}'_N$*

$$(7.13) \quad \frac{1}{i!j!} \left| \frac{\partial^i}{\partial s_1^i} \frac{\partial^j}{\partial s_2^j} D(s_1, s_2) \right| \ll (CK(\log_2 N + \log T))^{i+j} |D(s_1, s_2)|.$$

The above estimate is sufficient for our purposes at every point (s_1, s_2) apart from $(0, 0)$, which will appear in the main term. We will show an analogous result for the point $(0, 0)$ where η in (7.6) will be replaced by the larger value

$$(7.14) \quad \eta_0 = \frac{1}{\bar{d}^* \log_2 N}.$$

where we use the notation $\bar{d}, \bar{d}^*$ of (4.4). Next we have

$$\textbf{Lemma 5.} \quad \frac{1}{i!j!} \left| \frac{\partial^i}{\partial s_1^i} \frac{\partial^j}{\partial s_2^j} D(s_1, s_2) \right|_{s_1=s_2=0} \ll (\bar{d}^* \log_2 N)^{i+j} D(0, 0).$$

Proof. Let $d_1 = a - d$. Analogously to (7.8)–(7.9), we have, for $|s_1|, |s_2| \leq \eta_0$,

$$\begin{aligned}
(7.15) \quad & \frac{\partial}{\partial s_1} \log D_0(s_1, s_2) = d \frac{W'}{W}(s_1 + s_2) - a \frac{W'}{W}(s_1) \\
&= d \left(\frac{W'}{W}(s_1 + s_2) - \frac{W'}{W}(s_1) \right) - (a - d) \frac{W'}{W}(s_1) \\
&\ll K\eta_0 + d_1 \ll \frac{K}{\bar{d}^*} + d_1 \ll \sqrt{K} + d_1 \ll \bar{d}^*,
\end{aligned}$$

and

$$\begin{aligned}
(7.16) \quad & \frac{\partial}{\partial s_1}(\log G_1(s_1, s_2)) = \sum_{p \leq V} \frac{\log p}{p^{1+s_1}} \left(\frac{d}{p^{s_2} - p^{-(1+s_1)}} - \frac{d}{1 - p^{-(1+s_1)}} + \frac{d - a}{1 - p^{-(1+s_1)}} \right) \\
&\ll \sum_{p \leq V} \frac{\log p}{p^{1-\delta_1}} \left(\frac{K|p^{s_2} - 1|}{p^{-\delta_2}} + d_1 \right) \ll K \sum_{p \leq V} \frac{\eta_0 \log^2 p}{p^{1-\delta}} + d_1 \sum_{p \leq V} \frac{\log p}{p^{1-\delta}} \\
&\ll V^\delta \log V (K\eta_0 \log V + d_1) \ll K\eta_0 \log_2^2 N + d_1 \log_2 N \ll \bar{d}^* \log_2 N.
\end{aligned}$$

The treatment of G_2 will be similar to this and (7.10). By $|\nu_1(p) - \nu_3(p)| = |\nu_p(\mathcal{H}_1) - \nu_p(\mathcal{H}_1(p) \cap \mathcal{H}_2(p))| \leq |\mathcal{H}_1 \setminus \mathcal{H}_2| = a - d$ we have

$$\begin{aligned}
(7.17) \quad & \frac{\partial}{\partial s_1}(\log G_2(s_1, s_2)) = \\
& \sum_{\substack{p|\Delta \\ p>V}} \frac{\log p}{p^{1+s_1}} \left\{ \frac{\nu_1(p) - \nu_3(p) + \nu_3(p)(1 - p^{-s_2})}{1 - \frac{\nu_1(p)}{p^{1+s_1}} - \frac{\nu_2(p)}{p^{1+s_2}} + \frac{\nu_3(p)}{p^{1+s_1+s_2}}} - \frac{d_1}{1 - \frac{1}{p^{1+s_1}}} - d \left(\frac{1}{1 - \frac{1}{p^{1+s_1}}} - \frac{1}{p^{s_2} - \frac{1}{p^{1+s_1}}} \right) \right\} \\
& \ll \sum_{p|\Delta, p>V} \frac{\log p}{p^{1-\delta}} (d_1 + K(p^{\eta_0} - 1)) \ll d_1 \sum_{p|\Delta} \frac{\log p}{p^{1-\delta}} + K\eta_0 \sum_{\substack{p|\Delta \\ p \leq e^{2/\eta_0}}} \frac{\log^2 p}{p^{1-\delta}} + K \sum_{p|\Delta} \frac{\log p}{e^{1/\eta_0}} \\
& \ll d_1 \sum_{p \leq \log \Delta(1+o(1))} \frac{\log p}{p^{1-\delta}} + K\eta_0 \sum_{p \leq \log \Delta(1+o(1))} \frac{\log^2 p}{p^{1-\delta}} + \frac{K \log \Delta}{(\log N)^{\bar{d}^*}} \\
& \ll d_1 \log_2 \Delta + K\eta_0 \log_2^2 \Delta + \frac{1}{(\log N)^{\sqrt{K}-3}} \ll \log_2 N (d_1 + K\eta_0 \log_2 N) \ll \bar{d}^* \log_2 N.
\end{aligned}$$

Finally we have, similarly to above and (7.11), using $a - dp^{-s_2} = a - d + d(1 - p^{-s_2})$,

$$\begin{aligned}
(7.18) \quad & \frac{\partial}{\partial s_1} \log G_3(s_1, s_2) = \sum_{\substack{p|\Delta \\ p>V}} \frac{\log p}{p^{1+s_1}} \left\{ a \left(\frac{1}{1 - \frac{1}{p^{1+s_1+s_2}}} - \frac{1}{1 - \frac{1}{p^{1+s_1}}} \right) \right. \\
& \quad \left. + (a - dp^{-s_2}) \left(\frac{1}{1 - \frac{a}{p^{1+s_1}} - \frac{b}{p^{1+s_2}} + \frac{d}{p^{1+s_1+s_2}}} - \frac{1}{1 - \frac{1}{p^{1+s_1+s_2}}} \right) \right\} \\
& \ll \sum_{p>V} \frac{\log p}{p^{1-\delta_1}} \left\{ \frac{K}{p^{1-\delta}} + \frac{K}{p^{1-\delta}} (d_1 + K(p^{\eta_0} - 1)) \right\} \\
& \ll Kd_1 \sum_{p>V} \frac{\log p}{p^{2-2\delta}} + K^2\eta_0 \sum_{V < p \leq e^{1/\eta_0}} \frac{\log^2 p}{p^{2-2\delta}} + K^2 \sum_{p > e^{1/\eta_0}} \frac{\log p}{p^{2-2\delta-\eta_0}} \\
& \ll \frac{Kd_1}{V^{1-2\delta}} + \frac{K^2\eta_0 \log V}{V^{1-2\delta}} + \frac{K^2}{(\log N)^{(1-2\delta-\eta_0)\bar{d}^*}} \\
& \ll \frac{K}{V} (d_1 + K\eta_0 \log_2 N) + o(1) \ll d_1 + K\eta_0 \log_2 N \ll \bar{d}^*.
\end{aligned}$$

Now, (7.15)–(7.18) imply, by symmetry for $i = 1, 2$, that

$$(7.19) \quad \left| \frac{\partial}{\partial s_i} \log D(s_1, s_2) \right| \ll \eta_0^{-1}, \quad \text{for } |s_1|, |s_2| \leq \eta_0,$$

and therefore, similarly to (7.12) we have

$$(7.20) \quad \max_{|s'_1| \leq \eta_0, |s'_2| \leq \eta_0} |D(s'_1, s'_2)| \ll D(0, 0),$$

which by (7.2) proves Lemma 5.

8. CONTRIBUTION OF THE RESIDUE AT $s_1 = s_2 = 0$

This section will be devoted to the examination of $I_{1,1}$, the sum of the residues in (6.28).

The rather complicated formula (6.28) yields the main term and all secondary terms of the form $(\log R)^m$ exclusively for $m \in [d, d+u+v-1]$ and will additionally contribute to other secondary terms for $m \in [0, d-1]$. However, from the terms $I_{1,1}(i, j, \nu)$ belonging to the triplet (i, j, ν) in the triple summation, only those with

$\nu = 0, j = 0$ contribute to the main term of order $(\log R)^{d+u+v}$, since in all other terms the exponent of $\log R$ is $d + u + v - j - \nu$.

We have to work now more carefully than in [7]. For example, by the aid of Lemma 2 (a generalization of (8.16) in [7]) we will exactly evaluate the coefficients $A_{j,\nu}$ of $\frac{1}{j!\nu!} \frac{\partial^\nu}{\partial s_2^\nu} \frac{\partial^j}{\partial s_1^j} D(s_1, s_2)(\log R)^{v+d+u-j-\nu}$ in (6.28) as follows. Let $j, \nu \geq 0$,

$$(8.1) \quad m := i - j \geq 0, \quad y := v + d - \nu,$$

where we can assume by (6.28)

$$(8.2) \quad \nu \leq v + d + m \iff m \geq \nu - v - d = -y.$$

Then we have from (6.28), by notation (5.7), (8.1) and Lemma 2

$$(8.3) \quad \begin{aligned} A_{j,\nu} &= \frac{j!\nu!}{u!} \sum_{\substack{m=0 \\ m \geq -y}}^{u-j} \binom{u}{m+j} (-1)^m \binom{m+j}{j} \frac{d(d+1) \dots (d+m-1)}{(v+d+m-\nu)! \nu!} \\ &= \sum_{\substack{m=0 \\ m \geq -y}}^{u-j} \frac{(-1)^m}{(u-j-m)!m!} \cdot \frac{d(d+1) \dots (d+m-1)}{(v+d+m-\nu)!} \\ &= Z(d, u-j, v+d-\nu) = \frac{(v-\nu+1) \dots (v-\nu+u-j)}{(u-j)!(d+v-\nu+u-j)!}. \end{aligned}$$

We have to compare $A_{j,\nu}$ with $A_{0,0}$. This will be furnished by the following

Lemma 6. $|A'_{j,\nu}| := \left| \frac{A_{j,\nu}}{A_{0,0}} \right| \leq (CK)^{j+\nu}.$

Proof. $|A'_{j,\nu}| = \frac{(d+v+u)!}{(d+v+u-\nu-j)!} \cdot \frac{(u-j+1) \dots u}{(v+u-j+1) \dots (v+u)} \cdot |A''_{j,\nu}| \leq (CK)^{j+\nu} |A''_{j,\nu}|$, where

$$(8.4) \quad |A''_{j,\nu}| = \frac{|(v-\nu+1) \dots (v-\nu+u-j)|}{(v+1) \dots (v+u-j)}.$$

If $\nu \leq 2(v+1)$, then clearly $A''_{j,\nu} \leq 1$, so we may suppose

$$(8.5) \quad \nu = B(v+1), \quad B > 2.$$

In this case we have by $u-j \leq u \leq v < v+1$:

$$(8.6) \quad A''_{j,\nu} \leq \left(\frac{\nu}{v+1} \right)^{u-j} \leq B^{v+1} = B^{\nu/B} < 2^\nu,$$

since the maximum of $x^{1/x}$ in $[1, \infty)$ is attained at $x = e$ and $e^{1/e} < 2$.

Now we are ready to evaluate the crucial term $I_{1,1}$ by the aid of Lemmas 2, 5 and 6. Namely by (4.1), (4.4), $R \gg N^c$, (6.28), (8.3) and (6.24) we have

$$(8.7) \quad \begin{aligned} I_{1,1} &= A_{0,0}(\log R)^{d+u+v} \left\{ D(0,0) + \sum_{j=0}^u \sum_{\substack{\nu=0 \\ j+\nu \geq 1}}^{v+d+u-j} \frac{A'_{j,\nu}}{(\log R)^{j+\nu}} \cdot \frac{\frac{\partial^j}{\partial s_1^j} \frac{\partial^\nu}{\partial s_2^\nu} D(0,0)}{j!\nu!} \right\} \\ &= Z(d, u, v+d)(\log R)^{d+u+v} D(0,0) \left(1 + O \left(\sum_{j=0}^{\infty} \sum_{\substack{\nu=0 \\ j+\nu \geq 1}}^{\infty} \left(\frac{CK \bar{d}^* \log_2 N}{\log R} \right)^{j+\nu} \right) \right) \\ &= \frac{\binom{v+u}{u} (\log R)^{d+v+u} D(0,0)}{(d+v+u)!} \left(1 + O \left(\frac{CK \bar{d}^* \log_2 N}{\log R} \right) \right). \end{aligned}$$

The integral $I_{1,2}$ in (6.28) does not contribute to the main term and can be estimated relatively easily due to the presence of the term R^{s_2} ($s_2 \in \mathcal{L}_4$). In fact,

choosing $\eta^{-1} = 100(\log_2 N + \log T)$ as earlier, we obtain by Lemma 4, (5.2), (6.20), (6.26)–(6.27) for any $s_2 \in \mathcal{L}_4$,

$$(8.8) \quad Z(s_2) \ll \sum_{i=0}^u \sum_{j=0}^i \frac{(\log R)^{u-i} (CK)^{i-j}}{(u-i)!(i-j)!} (CK(\log_2 N + \log T))^j \frac{|D(0, s_2)| R^{\sigma_2}}{|s_2|^{d+i-j+v+1}} \\ \ll e^{C\sqrt{K} \log_2 N} \frac{(\log(|t_2| + 3))^{3K+O(\sqrt{K})} R^{\sigma_2}}{|s_2|^{b+O(\sqrt{K})}}.$$

Now Lemma 1 yields immediately by (6.24) and $R \gg N^c$

$$(8.9) \quad I_{1,2} = \frac{1}{2\pi i} \int_{\mathcal{L}_4} Z(s_2) ds_2 \ll e^{C\sqrt{K} \log_2 N - c\sqrt{\log N}} \\ \ll e^{-c\sqrt{\log N}}.$$

We may summarize (8.7) and (8.9) by $D(0, 0) = D_0(0, 0)G(0, 0) = G(0, 0) \neq 0$ (which is true by the admissibility condition) by (6.9)–(6.11) and (6.23) as

Lemma 7. *The integral I_1 in (6.26) satisfies the asymptotic*

$$(8.10) \quad I_1 = \frac{\binom{v+u}{u} (\log R)^{d+v+u} G(0, 0)}{(d+v+u)!} \left(1 + O\left(\frac{K \bar{d}^* \log_2 N}{\log R} \right) \right) + O(e^{-c\sqrt{\log N}}).$$

9. ESTIMATE OF THE INTEGRAL I_2

For I_2 in (6.25), after interchange of the two integrations we move the contour $\mathcal{L}_2$ for the inner integral over s_2 to the left to $\mathcal{L}_4$ passing a pole of order d at $s_2 = -s_1$ if $|t_2| \leq U$ and a pole of order $v+1$ at $s_2 = 0$ and obtain

$$(9.1) \quad I_2 = \frac{1}{2\pi i} \int_{\mathcal{L}_3} \text{Res}_{s_2=-s_1} \left(\frac{D(s_1, s_2) R^{s_1+s_2}}{s_1^{u+1} s_2^{v+1} (s_1+s_2)^d} \right) ds_1 + \frac{1}{2\pi i} \int_{\mathcal{L}_3} \text{Res}_{s_2=0} \left(\frac{D(s_1, s_2) R^{s_1+s_2}}{s_1^{u+1} s_2^{v+1} (s_1+s_2)^d} \right) ds_1 \\ + \frac{1}{(2\pi i)^2} \int_{\mathcal{L}_4} \int_{\mathcal{L}_3} F(s_1, s_2) \frac{R^{s_1}}{s_1^{a+u+1}} \frac{R^{s_2}}{s_2^{b+v+1}} ds_1 ds_2 + O(e^{-c\sqrt{\log N}}) \\ := I_{2,1} + I_{2,2} + I_{2,3} + O(e^{-c\sqrt{\log N}}).$$

By the argument of Lemma 1 and (6.21), the third integral $I_{2,3}$ is $\ll e^{-c\sqrt{\log N}}$. The second integral $I_{2,2}$ is completely analogous to $I_{1,2}$ in (6.28), which was estimated by $e^{-c\sqrt{\log N}}$ in (8.9), the only change being that the role of s_1 and s_2 is interchanged.

The residue in $I_{2,1}$ is zero if $d = 0$, while for $d \geq 1$ we have

$$(9.2) \quad \text{Res}_{s_2=-s_1} \left(\frac{D(s_1, s_2) R^{s_1+s_2}}{s_1^{u+1} s_2^{v+1} (s_1+s_2)^d} \right) = \lim_{s_2 \rightarrow -s_1} \frac{1}{(d-1)!} \frac{\partial^{d-1}}{\partial s_2^{d-1}} \left(\frac{D(s_1, s_2) R^{s_1+s_2}}{s_1^{u+1} s_2^{v+1}} \right) \\ = \frac{1}{(d-1)!} \sum_{j=0}^{d-1} \mathcal{B}_j(s_1, \mathcal{H}_1, \mathcal{H}_2) (\log R)^{d-1-j},$$

where

$$(9.3) \quad \mathcal{B}_j(s_1, \mathcal{H}_1, \mathcal{H}_2) = \binom{d-1}{j} \sum_{\nu=0}^j \binom{j}{\nu} \frac{\partial^{j-\nu}}{\partial s_2^{j-\nu}} D(s_1, s_2) \Big|_{s_2=-s_1} \cdot \frac{(-1)^\nu (v+1) \dots (v+\nu)}{(-1)^{\nu+v+1} s_1^{u+v+\nu+2}}.$$

We thus obtain

$$(9.4) \quad I_2 = \frac{1}{(d-1)!} \sum_{j=0}^{d-1} \mathcal{C}_j(\mathcal{H}_1, \mathcal{H}_2) (\log R)^{d-1-j} + O(e^{-c\sqrt{\log N}}),$$

where

$$(9.5) \quad \mathcal{C}_j(\mathcal{H}_1, \mathcal{H}_2) = \frac{1}{2\pi i} \int_{\mathcal{L}_3} \mathcal{B}_j(s_1, \mathcal{H}_1, \mathcal{H}_2) ds_1 \quad (j = 0, 1, 2, \dots, d-1).$$

It remains to estimate these quantities, which are independent of R .

We are allowed to transform the contour $\mathcal{L}_3$ in (9.5) to the contour $\mathcal{L}'$, defined in (5.4). Our task is now the estimation of the integral $\mathcal{C}_j$ on the new contour $\mathcal{L}' = \mathcal{L}'_0 \cup \mathcal{L}'_1$ since the integral on the horizontal segments $|t| = U$ is $O(e^{-c\sqrt{\log N}})$.

10. COMPARISON OF $D(s, -s)$ AND $D(0, 0)$

We have seen in Section 7 that by Lemma 4 we can estimate $\frac{\partial^i}{\partial s_1^i} \frac{\partial^j}{\partial s_2^j} D(s_1, s_2)$ with the aid of $D(s_1, s_2)$. We will show now how to estimate $|D(s, -s)|/D(0, 0)$ from above when s is on the contour $\mathcal{L}'$. This, together with Lemma 8 will play a crucial role in the estimation of $I_{2,1}$ which is the main part of I_2 .

First we note that if $s \in \mathcal{L}'$ is on the semicircle $\mathcal{L}'_0$, then by (7.12) we obtain

$$(10.1) \quad |D(s, -s)| \leq e^{C\sqrt{K}} D(0, 0), \quad (s \in \mathcal{L}'_0).$$

Thus, in the following we may suppose

$$(10.2) \quad s = it, \quad t > 0,$$

since $|D(-it, it)| = |D(it, -it)|$.

First we will examine the behavior of the functions $D_0(s, -s)$ and $G_1(s, -s)$ on the imaginary axis, which requires a lemma concerning $W(s)$ from (6.23).

Lemma 8. *There exist positive absolute constants t_0 and $t_1 > 1$ such that*

$$(10.3) \quad |W(it)| \geq e^{t^2/6} \geq 1 = W(0) \quad \text{for } |t| \leq t_0,$$

$$(10.4) \quad |W(it)| \geq t^{2/3} \quad \text{for } |t| \geq t_1.$$

Proof. We will use that in a neighborhood of $s = 0$ we have for the entire function $W(s)$ the representation

$$(10.5) \quad W(s) = 1 + \gamma_0 s + \sum_{\nu=1}^{\infty} \gamma_{\nu} s^{\nu+1}$$

where $\gamma_0 = \gamma$ is Euler's constant and (see [13], Notes on p. 49)

$$(10.6) \quad \gamma_0 = \gamma = 0.5772157\dots, \quad \gamma_1 = 0.07281\dots$$

This implies

$$(10.7)$$

$$\begin{aligned} |W(it)|^2 &= W(it)W(-it) = (1 + i\gamma t - \gamma_1 t^2 + O(t^3))(1 - i\gamma t - \gamma_1 t^2 + O(t^3)) \\ &= 1 + t^2(\gamma^2 - 2\gamma_1) + O(t^3) \quad \text{if } t \rightarrow 0. \end{aligned}$$

Now (10.6)–(10.7) prove (10.3) for $|t| \leq t_0$, while (10.4) clearly holds by (5.2).

Remark. If (10.3) is true for any t (which could be checked by computers, since t_0, t_1 are explicitly calculable), then the following simple lemma is not necessary.

Lemma 9. *Given any positive constants B_0, B_1, ε we have for any $t \in [B_0, B_1]$ and any $X > C(B_0, B_1, \varepsilon)$*

$$(10.8) \quad J(t, X) := \prod_{p \leq X} \frac{|1 - p^{-1-it}|}{1 - p^{-1}} \geq c(B_0, B_1)(\log X)^{1/2-\varepsilon}.$$

Proof. Let us fix t . Since every factor is at least 1, we can neglect those with $\cos(t \log p) > 0$. On the other hand, if $\cos(t \log p) \leq 0$, then we have

$$(10.9) \quad \log \left(\frac{|1 - p^{-1-it}|}{1 - p^{-1}} \right) > \log \frac{1}{1 - p^{-1}} > \frac{1}{p}.$$

The primes satisfying $\cos(t \log p) \leq 0$ are in intervals of type

$$(10.10) \quad I_j = \left[\exp \left(\frac{\pi (2j + \frac{1}{2})}{t} \right), \exp \left(\frac{\pi (2j + \frac{3}{2})}{t} \right) \right] =: [e^{m_j}, e^{m_j + \pi/t}]$$

and $I_j \subset [1, X]$ if $2\pi(j + 3/4)/t \leq \log X$, that is, if

$$(10.11) \quad j \leq \frac{t \log X}{2\pi} - \frac{3}{4} =: j^*.$$

Using the prime number theorem we obtain by partial summation

$$(10.12) \quad \sum_{p \in I_j} \frac{1}{p} \sim \int_{e^{m_j}}^{e^{m_j + \pi/t}} \frac{dx}{x \log x} = \log \frac{2j + \frac{3}{2}}{2j + \frac{1}{2}} = \frac{1}{2j} + O\left(\frac{1}{j^2}\right).$$

Hence, by (10.9) we have

$$(10.13) \quad \log J(t, X) > \sum_{1 \leq j \leq j^*} \frac{1 - \varepsilon}{2j} + O(1) > \frac{1 - \varepsilon}{2} \log_2 X - c'(B_0, B_1).$$

Remark. Working more carefully we could prove Lemma 9 with $(\log X)^{\frac{1}{2} - \varepsilon}$ replaced by $\log X$. But actually any lower bound larger than $C(t_0, t_1) = \max_{t_0 \leq t \leq t_1} |W(it)|^{-1}$ would suffice for us.

Taking into account the trivial relation

$$(10.14) \quad |1 - p^{-1-it}|^{-1} \leq |1 - p^{-1}|^{-1},$$

we obtain from Lemmas 8, 9 the following

Lemma 10. *We have, with a sufficiently small constant $t_0 < 1$ and suitable positive constants c' and c'' ,*

$$(10.15) \quad E_0(t) := \left| \frac{D_0(it, -it)G_1(it, -it)}{D_0(0, 0)G_1(0, 0)} \right| \leq e^{-c'(a+b)t^2}, \quad \text{if } |t| \leq t_0,$$

and for any $t > t_0$ and $N > N_0$,

$$(10.16) \quad E_0(t) \leq e^{-c''(a+b)} \max(1, |t|)^{-(a+b)/2}.$$

Proof. By (10.14) and the definition of D_0 in (6.23) we clearly have

$$(10.17) \quad |G_1(it, -it)| \leq G_1(0, 0),$$

$$(10.18) \quad |D_0(it, -it)| = |W(it)|^{-(a+b)}, \quad D_0(0, 0) = W(0) = 1,$$

which immediately imply

$$(10.19) \quad E_0(t) \leq |W(it)|^{-(a+b)}.$$

Hence, by Lemma 8 we have (10.15) and (10.16) for $|t| \geq t_1$. Finally, for $t_0 \leq |t| \leq t_1$ we have by Lemma 9

$$(10.20) \quad \begin{aligned} E_0(t) &= (J(|t|, V)|W(it)|)^{-(a+b)} \leq (c(t_0, t_1)(\log V)^{1/3} \cdot C^{-1}(t_0, t_1))^{-(a+b)} \\ &\leq (c \log_2 N)^{-(a+b)/3} \leq (e^{-c''} t_1)^{-(a+b)}, \end{aligned}$$

which proves (10.16).

We will continue our study of $D(it, -it)$ with that of

$$(10.21) \quad L_4(t) := \log \frac{|G_4(it, -it)|}{G_4(0, 0)} = \operatorname{Re} \log \frac{G_4(it, -it)}{G_4(0, 0)}.$$

We first divide each term by $\left(1 + \frac{\nu_3(p)}{p}\right) \left(1 - \frac{1}{p}\right)^d$ in the product representation of both $G_4(it, -it)$ and $G_4(0, 0)$. After this we take the logarithm of each term and use the formula

$$(10.22) \quad \log(1 - z) = -\left(z + \sum_{m=2}^{\infty} \frac{z^m}{m}\right), \quad \text{if } |z| < 1.$$

Now we separate the effect of the linear terms and those of order $m \geq 2$ and write accordingly

$$(10.23) \quad L_4(t) = L_{4,1}(t) + L_{4,2}(t).$$

We have by the trivial relations $\nu_1(p) \leq a$, $\nu_2(p) \leq b$,

$$(10.24) \quad L_{4,1}(t) = \sum_{p>V} \left(\frac{a+b}{p} - \frac{\nu_1(p) + \nu_2(p)}{p(1 + \nu_3(p)/p)} \right) (\cos(t \log p) - 1) \leq 0.$$

(We remark that the sum is convergent, since $\nu_1(p) = a$, $\nu_2(b) = p$ for $p \nmid \Delta$.)

The logarithms of the higher order terms of G_4 which do not involve the functions $\nu_i(p)$ can be estimated from above in modulus for any t by

$$(10.25) \quad (a+b) \sum_{p>V} \sum_{m=2}^{\infty} \frac{2}{mp^m} \leq C(a+b) \frac{1}{V \log V} \leq \frac{C}{(\log_2 N)^3}.$$

Similarly we have for the contribution of the numerator to $L_{4,2}(t)$ the upper estimate (valid for any t)

$$(10.26) \quad \sum_{p>V} \sum_{m=2}^{\infty} \frac{2(a+b)^m}{mp^m} \leq \frac{C(a+b)^2}{V \log V} \leq \frac{C(a+b)}{(\log_2 N)^3}.$$

We see from (10.16) that this estimation is sufficient for $|t| \geq t_0$ but not for small values of t . Then, working more carefully we have for the contribution of the terms of G_4 involving the functions $\nu_i(p)$ to $L_{4,2}(t)$ the upper estimate

$$(10.27) \quad \begin{aligned} & \sum_{p>V} \sum_{m=2}^{\infty} \sum_{j=0}^m \binom{m}{j} \frac{(\nu_1(p))^j (\nu_2(p))^{m-j}}{m(p + \nu_3(p))^m} (1 - \cos((m-2j)t \log p)) \\ & \leq C \sum_{p>V} \sum_{m=2}^{\infty} \frac{(a+b)^m}{mp^m} t^2 m^2 \log^2 p \leq C(a+b)^2 t^2 \sum_{p>V} \frac{\log^2 p}{p^2} \\ & \leq \frac{C(a+b)^2 t^2 \log V}{V} \leq \frac{C(a+b)t^2}{\log_2 N}. \end{aligned}$$

It is easier to see that the contribution of the terms of G_4 which do not involve the functions $\nu_i(p)$ to $L_{4,2}$ are majorized by

$$(10.28) \quad C(a+b)t^2 \sum_{p>V} \frac{\log^2 p}{p^2} \leq \frac{Ct^2}{\log_2 N}.$$

Summarizing (10.21)–(10.28) we have proved

Lemma 11. $\frac{|G_4(it, -it)|}{G_4(0, 0)} \leq \exp \left(C \frac{(a+b)}{\log_2 N} \min(1, t^2) \right).$

Comparing the above with (10.15)–(10.16) we see that (10.15)–(10.16) remain valid if we multiply them by $G_4(it, -it)/G_4(0, 0)$. This proves the final result of this section:

Lemma 12. $|D(it, -it)| \leq \max(1, |t|)^{-(a+b)/2} D(0, 0)$ for any real t .

Together with (10.1) this implies

Lemma 13. $|D(s, -s)| \leq e^{C\sqrt{K}} \max(1, |t|)^{-(a+b)/2} D(0, 0)$ for $s \in \mathcal{L}'$.

11. ESTIMATE OF I_2 . EVALUATION OF I

In this section we will estimate the integral $I_{2,1}$ based on formulas (9.3)–(9.5), using Lemmas 4 and 13.

First we obtain from the above lemmas for $s \in \mathcal{L}'$ by $j \leq d \leq \min(a, b) \leq K$, $v \leq \sqrt{K}$

(11.1)

$$\begin{aligned} \mathcal{B}_j(s, \mathcal{H}_1, \mathcal{H}_2) &\ll \frac{d^j}{|s|^{u+v+2}} \sum_{\nu=0}^j (CK(\log_2 N + \log T))^{j-\nu} \prod_{i=1}^{\nu} \frac{\left(\frac{v}{i} + 1\right)}{|s|} |D(s, -s)| \\ &\ll \frac{e^{C\sqrt{K}} D(0, 0) d^j (\log(|t| + 3))^j \delta_0^{-(u+v)}}{\max(1, \sqrt{|t|})^{a+b}} \frac{1}{|s|^2} \sum_{\nu=0}^j (CK \log_2 N)^{j-\nu} (K \log_2 N)^{\nu} \\ &\ll e^{C\sqrt{K}} (CK^2 \log_2 N)^j \cdot \frac{\delta_0^{-(u+v)}}{|s|^2} D(0, 0). \end{aligned}$$

Integrating the above upper bound along $\mathcal{L}'$ we obtain

$$(11.2) \quad \mathcal{C}_j(\mathcal{H}_1, \mathcal{H}_2) \ll e^{C\sqrt{K}} (CK^2 \log_2 N)^j \delta_0^{-(u+v+1)} D(0, 0).$$

Finally, summation over $j \leq d-1$ yields in (9.4) by $R \gg N^c$

$$\begin{aligned} (11.3) \quad I_{2,1} &\ll \frac{e^{C\sqrt{K}} D(0, 0) \delta_0^{-(u+v+1)} (\log R)^{d-1}}{(d-1)!} \cdot \sum_{j=0}^{d-1} \left(\frac{CK^2 \log_2 N}{\log R} \right)^j \\ &\ll \frac{e^{C(u+v)} D(0, 0) (\log R)^{d-1} (\sqrt{K} \log_2 N)^{u+v+1}}{(d-1)!} \\ &\ll \frac{D(0, 0) (\log R)^{d+u+v}}{(d+u+v)!} \cdot \left(\frac{CK^{3/2} \log_2 N}{\log R} \right)^{u+v+1} \\ &\ll \frac{D(0, 0) (\log R)^{d+u+v}}{(d+u+v)!} (\log N)^{-\sqrt{K}/50}. \end{aligned}$$

This implies by (9.1) and (9.4)

$$(11.4) \quad I_2 \ll \frac{D(0, 0) (\log R)^{d+u+v}}{(d+u+v)!} (\log N)^{-\sqrt{K}/50} + e^{-c\sqrt{\log N}}.$$

This yields by Lemma 7 the final asymptotic evaluation of I in (6.22) by $D(0, 0) = G(0, 0)$ as

$$(11.5) \quad I = \frac{\binom{v+u}{u} (\log R)^{d+v+u} G(0, 0)}{(d+v+u)!} \left(1 + O\left(\frac{K \bar{d}^* \log_2 N}{\log R} \right) \right) + O(e^{-c\sqrt{\log N}}),$$

where

$$(11.6) \quad G(0, 0) = \prod_{p \nmid P} \left(1 - \frac{\nu_p(\mathcal{H})}{p} \right) \prod_p \left(1 - \frac{1}{p} \right)^{-|\mathcal{H}|} = \frac{\mathfrak{S}(\mathcal{H})P}{|A(\mathcal{H})|},$$

thereby proving Theorem 4.

12. A BOMBIERI–VINOGRADOV TYPE THEOREM

In the present section we will prove a modified Bombieri–Vinogradov theorem, where the examined moduli are all multiples of a single modulus M . It would facilitate our task if we were entitled to use the following hypothesis.

Hypothesis $S(Y)$. *If $L(1 - \delta, \chi) = 0$ for a $\delta > 0$ and a real primitive character $\chi \pmod{q}$, $q \leq Y$, then*

$$(12.1) \quad \delta \geq \frac{1}{3 \log Y}$$

for $Y > C_0$, an explicitly calculable absolute constant.

We note that we have the effective unconditional estimate ([6], [16]), valid for $q > q_0$:

$$(12.2) \quad \delta \geq \frac{1}{\sqrt{q}}.$$

A further observation (similar to that of Maier [14]) is that by the Landau–Page theorem (cf. Davenport [2, §14]), with some constant c in place of $1/3$, or Pintz [17] with $(1/2 + o(1))$ for any given Y there is at most one real primitive character χ_1 which does not fulfill (12.1). This makes it possible to turn Hypothesis $S(Y)$ into a theorem, valid for a sequence $Y = Y_n \rightarrow \infty$ (for $n > n_0$, an explicitly calculable absolute constant) with

$$(12.3) \quad Y_n \leq \exp\left(\sqrt{Y_{n-1}}\right).$$

In order to show this, suppose that (12.1) is false for a sufficiently large Y' , i.e. by (12.2) there exists a $\chi_1 \pmod{q_1 \leq Y'}$ such that $L(1 - \delta_1, \chi_1) = 0$ with

$$(12.4) \quad \frac{1}{\sqrt{Y'}} \leq \min\left(\frac{1}{\sqrt{q_1}}, c_0\right) \leq \delta_1 < \frac{1}{3 \log Y'}.$$

Let us choose $\tilde{Y} > Y'$ in such a way, that

$$(12.5) \quad \tilde{Y} = \exp\left(\frac{1}{3\delta_1}\right) \Leftrightarrow \delta_1 = \frac{1}{3 \log \tilde{Y}}.$$

Then for any other zero $1 - \delta_2$ belonging to a real primitive $\chi_2 \pmod{q_2}$, $q_2 \leq \tilde{Y}$, we have by the Landau–Page theorem in the version of Pintz [17]

$$(12.6) \quad \max(\delta_1, \delta_2) > \frac{1}{3 \log \tilde{Y}} \Leftrightarrow \delta_2 > \frac{1}{3 \log \tilde{Y}}.$$

Now, (12.4)–(12.6) show that (12.1) is true for a value $Y = \tilde{Y}$ satisfying

$$(12.7) \quad Y' < \tilde{Y} < \exp(\sqrt{Y'}/3).$$

We can formulate this as

Lemma 14. *Hypothesis $S(Y)$ holds for a sequence $Y_n \rightarrow \infty$ with*

$$(12.8) \quad Y_n \leq \exp\left(\sqrt{Y_{n-1}}\right)$$

where Y_0 can be chosen with $Y_0 < C_0$, an explicitly calculable absolute constant.

An alternative to this Lemma and this approach would be to use Heath-Brown's theorem [10] (but only in case of Theorem 1) according to which either

(i) $S(Y)$ holds for every $Y > C$, with some absolute constant C ,

or

(ii) there are infinitely many twin primes.

The significance of the real zeros in Hypothesis $S(Y)$ is that a similar inequality holds with $\operatorname{Re} \varrho$ in place of $1 - \delta$ and with a constant c_0 in place of $1/3$ if $\operatorname{Im} \varrho$ is not too large; this is the standard zero-free region of L -functions (cf. Davenport [2, §14]).

Lemma 15. *There exists an explicitly calculable absolute constant $c_0 < 1/3$ such that $L(s, \chi) \neq 0$ in the region*

$$(12.9) \quad \sigma > 1 - \frac{c_0}{\log(q(|t| + 3))},$$

apart from possible real exceptional zeros of real L -functions.

Now we are in a position to formulate and prove the following theorem (which is similar to but stronger than Lemma 6 of Maier [14]).

Theorem 6. *Let c^* be an arbitrary, fixed constant. Let $Y = Y(X)$ be a strictly monotonically increasing function of X with*

$$(12.10) \quad \exp(2\sqrt{\log X}) \leq Y(X) \leq X.$$

Then there exists a sequence $X_n \rightarrow \infty$ satisfying $X_1 < C'_0$, an explicitly calculable constant, with the following property. Let $X = X_n$, $\mathcal{L} = \log X$, M be a natural number $\leq \min(\sqrt{Y(X)}/4, X^{1/8})$,

$$(12.11) \quad Q^* = X^{1/2} M^{-3} \exp(-c^* \sqrt{\log X}),$$

$$(12.12) \quad E^*(X, q) := \max_{x \leq X} \max_{(a, q)=1} |E(X, q, a)| := \max_{x \leq X} \max_{(a, q)=1} \left| \sum_{\substack{p \leq x \\ p \equiv a \pmod{q}}} \log p - \frac{x}{\varphi(q)} \right|,$$

Then

$$(12.13) \quad \sum_{\substack{q \leq Q^* \\ (q, M)=1}} E^*(X, Mq) \ll \frac{X}{M} \mathcal{L}^{15} \exp\left(-c_2 \frac{\log X}{\log Y(X)}\right),$$

where $c_2 = \min(c^/6, c_0/4)$.*

Remark. The above theorem holds with any X , for which $S(Y(X)) = S(Y)$ is true, i.e.

$$(12.14) \quad L(s, \chi) \neq 0 \quad \text{for } s \in \left(1 - \frac{1}{3 \log Y}, 1\right]$$

holds without exception for all real primitive characters $\chi \pmod{q}$, where

$$(12.15) \quad q \leq Y = Y(X).$$

Proof. We will choose our sequence $X_n = Y^{-1}(Y_n)$, where Y_n is the sequence supplied by Lemma 14 (for which $S(Y)$ is true) and Y^{-1} is the inverse function of $Y(X)$. Alternatively, if (12.14)–(12.15) hold, then we can choose X as an arbitrary sufficiently large number. In both cases $S(Y)$, i.e. (12.14)–(12.15) hold. Using the explicit formula for primes in arithmetic progressions with $T^* = \sqrt{X} \log^2 X$ ($\varrho = \beta + i\gamma = 1 - \delta + i\gamma$ denotes a generic zero of an L -function) we obtain (cf. Davenport [2, §19] for any a with $(a, q) = 1$, $q \leq Q^*$, $y \leq X$ the relation

$$(12.16) \quad E(y, q, a) = -\frac{1}{\varphi(q)} \sum_{\chi(q)} \bar{\chi}(a) \sum_{\substack{\varrho = \varrho_\chi \\ \beta \geq 1/2, |\gamma| \leq T^*}} \frac{y^\varrho}{\varrho} + O(\mathcal{L}^2 \sqrt{y}).$$

The effect of the last error term is clearly suitable, $O(Q^* \mathcal{L}^2 \sqrt{X})$ in total. We can classify zeros of all primitive L -functions mod $q\widetilde{M} \leq Q^* M, (\widetilde{M} \mid M)$, up to height T^* into $O(\mathcal{L}^4)$ classes $B(\kappa, \lambda, \mu, \nu)$ by Lemma 15, as

$$(12.17) \quad \widetilde{M} \in [M_\lambda/2, M_\lambda), \quad q \in [Q_\nu/2, Q_\nu), \quad \gamma \in [T_\mu/2, T_\mu), \quad \delta \in \left[\frac{\kappa c_0}{\mathcal{L}}, \frac{(\kappa+1)c_0}{\mathcal{L}} \right),$$

where

$$(12.18) \quad M_\lambda = 2^\lambda \leq 2M, \quad Q_\nu = 2^\nu \leq 2Q^*, \quad T_\mu = 2^\mu \leq 2T^*, \quad \frac{\kappa c_0}{\mathcal{L}} \leq \frac{1}{2},$$

with the additional class of index 0: $\gamma \in [0, 1) = [0, T_0)$. The set of quadruples $\kappa, \lambda, \mu, \nu$ satisfying (12.18) with $\nu \geq 1, \mu \geq 0, \lambda \geq 1, \kappa \geq 0$ will be denoted by $\mathcal{B}$.

In this case we have clearly by (12.16), similarly to Davenport [2, §28],

$$(12.19) \quad \sum_{\substack{q \leq Q^* \\ (q, \widetilde{M})=1}} E^*(X, qM) \ll \frac{X}{M} \mathcal{L}^6 \max_{\kappa, \lambda, \mu, \nu \in \mathcal{B}} \frac{N^*(1 - \frac{(\kappa+1)c_0}{\mathcal{L}}, M_\lambda Q_\nu, T_\mu)}{Q_\nu T_\mu} X^{-c_0 \kappa / \mathcal{L}},$$

where

$$(12.20) \quad N^*(\sigma, Q, T) = \sum_{Q/2 < q \leq Q} \sum_{\substack{\chi(q) \\ \chi \text{ primitive}}} \sum_{\substack{\varrho = \varrho_\chi \\ \beta \geq \sigma, |\gamma| \leq T}} 1.$$

We will see that, in order to prove our theorem, it will be enough to prove for any quadruple $\delta, M'Q, T$ with the property (cf. (12.9), (12.14)–(12.15))

$$(12.21) \quad \begin{aligned} \frac{c_0}{\log(QM'T)} &\leq \delta \leq \frac{1}{2}, \quad 2 \leq M' \leq M, \quad 2 \leq Q \leq Q^*, \quad 1 \leq T \leq T^* \quad \text{or} \\ \frac{c_0}{\log Y} &\leq \delta \leq \frac{1}{2}, \quad 2 \leq M' \leq M, \quad Q \leq \sqrt{Y}, \quad T = T_0 = 1, \\ 0 &\leq \delta \leq \frac{1}{2}, \quad 2 \leq M' \leq M, \quad Q > \sqrt{Y}, \quad T = T_0 = 1 \end{aligned}$$

the crucial inequality

$$(12.22) \quad N^*(1 - \delta, M'Q, T) \ll \mathcal{L}^9 QT X^\delta \exp(-c \log X / \log Y)$$

with some positive absolute constant c . The first line in (12.21) is meant to cover all non-real zeros, the second and third lines are meant to cover the real zeros.

We will use Theorem 12.2 of Montgomery [15]

$$(12.23) \quad N^*(1 - \delta, Q, T) \ll (Q^2 T)^{\frac{3\delta}{1+\delta}} (\log QT)^9.$$

(We do not need for the range $\delta \leq 1/5$ the stronger inequality of Theorem 12.2 of [15] with the exponent $3\delta/(1+\delta)$ replaced by the smaller $2\delta/(1-\delta)$.) Since $3\delta/(1+\delta) \leq 1$, (12.22) will follow if we can show

$$(12.24) \quad M^{6\delta}(Q)^{\frac{6\delta}{1+\delta}-1} \ll X^\delta e^{-c_2 \sqrt{\log X}} \text{ with } c_2 = c^*/6.$$

Since in the range $0 \leq \delta \leq 1/2$ we have $\frac{6\delta}{1+\delta} - 1 \leq 2\delta$, this is true by the definition $Q^* = X^{1/2} M^{-3} \exp(-c^* \sqrt{\log X})$, if $\delta \geq 1/12$.

In case of $\delta \leq 1/12$ we have by (12.23)

$$(12.25) \quad N^*(1 - \delta, M'Q, T) \ll (QT)^{1/2} M^{6\delta}.$$

If we have here $QT \geq \exp(\sqrt{\log X})$, then (12.25) directly implies (12.22), since

$$(12.26) \quad \frac{N^*(1 - \delta, M'Q, T)}{QT} \ll (QT)^{-1/2} M^{6\delta} \ll X^\delta \exp(-\sqrt{\log X}/2).$$

If $\delta \leq 1/12$ and $QT \leq e^{\sqrt{\log X}} \leq \sqrt{Y}$, then $\delta \geq c_0/\log(MQT)$ or $\delta \geq \frac{1}{3 \log Y} > \frac{c_0}{\log Y}$ by (12.14)–(12.15), since the modulus of the corresponding primitive character is $q\widetilde{M} \leq 4QM \leq Y$. Hence,

$$(12.27) \quad \left(\frac{M^6}{X}\right)^\delta \leq X^{-\delta/4} \leq \exp\left(-\frac{c_0}{8} \min\left(\frac{\log X}{\log QT}, \frac{\log X}{\log M}, \frac{\log X}{\log \sqrt{Y}}\right)\right) \\ = \exp\left(-\frac{c_0}{8} \frac{\log X}{\log \sqrt{Y}}\right).$$

Remark. The condition for Q^* could be weakened to $Q^* < X^{1/2}M^{-1} \exp(-cf(X, Y))$ but this has no significance in our application.

13. PROOF OF THEOREM 5

The method of proof of Theorem 5 is quite similar to that of Theorem 4. The basic difference is that instead of the trivial problem of the distribution of integers in arithmetic progressions we have to use properties of the distribution of primes in arithmetic progressions. Since we have to consider the (weighted) sum of the error terms in the formula for the number of primes in arithmetic progressions, the Bombieri–Vinogradov theorem can help us. However, due to the relatively weak estimate of the original Bombieri–Vinogradov theorem, it does not lead to better results than $\liminf_{n \rightarrow \infty} (p_{n+1} - p_n)/\log p_n = 0$. That is partly why we need to use Theorem 6 instead. Our situation is even more complicated here, since we need the moduli of the progressions to be the multiples of a number V . Fortunately our present Theorem 6 solves this problem in a completely satisfactory way, even without loss if $P = M \leq \exp((1 + o(1))\sqrt{\log N})$ which is now the case by $V = \sqrt{\log N}$.

We will suppose that $N = X_n/3$, n is sufficiently large, and $M = P$ in Theorem 6. (If we use Heath-Brown’s theorem [10] we may assume Hypothesis $S(Y)$ for any N and then N can be an arbitrary, sufficiently large integer.)

In the course of proof we will follow closely the analogous proofs of Propositions 4 and 5 in Sections 7–9 of [7], so we will sometimes omit details. Let

$$(13.1) \quad \Theta(x; q, a) := \sum_{\substack{p \leq x \\ p \equiv a \pmod{q}}} \log p = [(a, q) = 1] \frac{x}{\phi(q)} + E(x; q, a),$$

where $[S]$ is 1 if the statement S is true and 0 if S is false. We have for a regular residue class $\tilde{a}$ with respect to $\mathcal{H}$ and P

$$(13.2) \quad \tilde{S}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, \tilde{a}, h_0) := \sum_{n=N+1}^{2N} \Lambda_R(n; \mathcal{H}_1, \ell_1) \Lambda_R(n; \mathcal{H}_2, \ell_2) \theta(n + h_0) \\ = \frac{1}{(K + \ell_1)!(K + \ell_2)!} \sum_{d, e \leq R} \mu(d) \mu(e) \left(\log \frac{R}{d}\right)^{K + \ell_1} \left(\log \frac{R}{e}\right)^{K + \ell_2} \sum_{\substack{1 \leq n \leq N, n \equiv \tilde{a} \pmod{P} \\ d | P_{\mathcal{H}_1}(n), e | P_{\mathcal{H}_2}(n)}} \theta(n + h_0).$$

For the inner sum, we let $d = a_1 a_{12}$, $e = a_2 a_{12}$ where $(d, e) = a_{12}$, and thus a_1, a_2 , and a_{12} are pairwise relatively prime. We may suppose in the following $(d, P) = (e, P) = 1$, otherwise the last sum would be zero, since by the regularity of $\tilde{a}$ we have $(P_{\mathcal{H}_1}(n), P) = (P_{\mathcal{H}_2}(n), P) = 1$. The n for which $d | P_{\mathcal{H}_1}(n)$ and $e | P_{\mathcal{H}_2}(n)$ cover certain residue classes modulo $[d, e]$. If $n \equiv b' \pmod{a_1 a_2 a_{12}}$ is such a residue class, then letting $m = n + h_0 \equiv b' + h_0 \pmod{a_1 a_2 a_{12}}$, $b \equiv b' \pmod{a_1 a_2 a_{12}}$,

$b \equiv \tilde{a} \pmod{P}$ we see this residue class contributes to the inner sum

$$(13.3) \quad \sum_{\substack{N+1+h_0 \leq m \leq 2N+h_0 \\ m \equiv b+h_0 \pmod{a_1 a_2 a_{12} P}}} \theta(m) = \theta(2N+h_0; a_1 a_2 a_{12} P, b+h_0) - \theta(N+h_0; a_1 a_2 a_{12} P, b+h_0) \\ = [(b+h_0, a_1 a_2 a_{12} P) = 1] \frac{N}{\phi(a_1 a_2 a_{12} P)} + O(E^*(3N, a_1 a_2 a_{12} P)).$$

We need to determine the number of these residue classes where $(b+h_0, a_1 a_2 a_{12} P) = 1$ so that the main term is non-zero. The condition $(\tilde{a}+h_0, P) = (b+h_0, P) = 1$ is equivalent to $\tilde{a}$ being regular with respect to $\mathcal{H}^0$, since $\tilde{a}$ is regular with respect to $\mathcal{H}$. Thus we will assume from now on that $\tilde{a}$ is regular with respect to $\mathcal{H}^0$. If $p|a_1$ then $b \equiv -h_j \pmod{p}$ for some $h_j \in \mathcal{H}_1$, and therefore $b+h_0 \equiv h_0 - h_j \pmod{p}$. Thus, if h_0 is distinct modulo p from all the $h_j \in \mathcal{H}_1$ then all $\nu_p(\mathcal{H}_1)$ residue classes satisfy the relatively prime condition, while otherwise $h_0 \equiv h_j \pmod{p}$ for some $h_j \in \mathcal{H}_1$ leaving $\nu_p(\mathcal{H}_1) - 1$ residue classes with a non-zero main term. We introduce the notation $\nu_p^*(\mathcal{H}_1)$ for this number in either case, where we define for a set $\mathcal{G}$

$$(13.4) \quad \nu_p^*(\mathcal{G}) = \nu_p(\mathcal{G}^0) - 1.$$

and

$$(13.5) \quad \mathcal{G}^0 = \mathcal{G} \cup \{h_0\}.$$

We extend this definition to $\nu_d^*(\mathcal{H}_1)$ for squarefree numbers d by multiplicativity. (The function ν_d^* is familiar in sieve theory, see [8].) The same applies for $\nu_d^*(\mathcal{H}_2)$ and $\overline{\nu}_d^*((\mathcal{H}_1 \overline{\cap} \mathcal{H}_2))$, as in (6.2).

Since $E(n; q, a) \ll (\log N)$ if $(a, q) > 1$ and $q \leq N$ we conclude

$$(13.6) \quad \sum_{\substack{N+1 \leq n \leq 2N, n \equiv \tilde{a} \pmod{P} \\ d|P_{\mathcal{H}_1}(n), e|P_{\mathcal{H}_2}(n)}} \theta(n+h_0) = \nu_{a_1}^*(\mathcal{H}_1) \nu_{a_2}^*(\mathcal{H}_2) \overline{\nu}_{a_{12}}^*((\mathcal{H}_1 \overline{\cap} \mathcal{H}_2)) \frac{N}{\phi(a_1 a_2 a_{12} P)} \\ + O(d_K(a_1 a_2 a_{12}) (|E^*(3N; a_1 a_2 a_{12} P)|)).$$

Let $\sum^{(P)}$ denote that the summation variables are relatively prime to P and to each other. Substituting this into (13.2) we conclude by $\ell_i \leq K$

$$(13.7) \quad \tilde{\mathcal{S}}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, \tilde{a}, h_0) \\ = \frac{N}{\varphi(P)(K+\ell_1)!(K+\ell_2)!} \sum_{\substack{a_1 a_{12} \leq R \\ a_2 a_{12} \leq R}}^{(P)} \frac{\mu(a_1) \mu(a_2) \mu(a_{12})^2 \nu_{a_1}^*(\mathcal{H}_1) \nu_{a_2}^*(\mathcal{H}_2) \overline{\nu}_{a_{12}}^*((\mathcal{H}_1 \overline{\cap} \mathcal{H}_2))}{\phi(a_1 a_2 a_{12})} \\ \times \left(\log \frac{R}{a_1 a_{12}} \right)^{K+\ell_1} \left(\log \frac{R}{a_2 a_{12}} \right)^{K+\ell_2} \\ + O \left((\log R)^{4K} \sum_{\substack{a_1 a_{12} \leq R \\ a_2 a_{12} \leq R}}^{(P)} d_K(a_1 a_2 a_{12}) E^*(3N; a_1 a_2 a_{12} P) \right) \\ = \frac{N}{\varphi(P)} \tilde{\mathcal{T}}_R(\mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, h_0) + O((\log R)^{4K} \mathcal{E}_K(N)).$$

Using the notation $R^2 = Q^*$, we obtain from Theorem 6 by the trivial estimate $|E(X, Pq, a)| \leq 2q^{-1}P^{-1}X \log X$ (for $Pq \leq X$), Lemma 3 and by Hölder's inequality with parameters $\alpha = \nu + 1$, $\beta = (\nu + 1)/\nu$ where $\nu \in \mathbb{Z}^+$, $c' \log(K + 1) \leq \nu \leq c'' \log(K + 1)$, uniformly for $K \leq (\log N)/(2C)$, ($\sum^{b*}$ means summation over squarefree integers which are relatively prime to P)

(13.8)

$$\begin{aligned}
|\mathcal{E}_K(N)| &\leq \sum_{q \leq Q^*}^{b*} d_K(q) E^*(3N, Pq) \sum_{q=a_1 a_2 a_{12}} 1 \\
&\leq \sum_{q \leq Q^*}^{b*} d_K(q) d_3(q) E^*(3N, Pq) = \sum_{q \leq Q^*}^{b*} \frac{d_{3K}(q)}{q^{1/\beta}} \cdot q^{1/\beta} E^*(3N, Pq) \\
&\leq \left(\sum_{q \leq Q^*}^{b*} \frac{(d_{3K}(q))^\beta}{q} \right)^{1/\beta} \left(\sum_{q \leq Q^*}^{b*} q^{\alpha/\beta} (E^*(3N, Pq))^\alpha \right)^{1/\alpha} \\
&\leq \left(1 + \frac{1}{2} \log N \right)^{CK} (6NP^{-1} \log 3N)^{\nu/(\nu+1)} \left(\sum_{q \leq Q^*}^{b*} E^*(3N, Pq) \right)^{\frac{1}{\nu+1}} \\
&\ll (\log N)^{CK+1} NP^{-1} \exp \left(-\frac{c_2 \sqrt{\log N}}{\nu+1} \right) \\
&\leq NP^{-1} \exp \left((CK+1) \log_2 N - c_2(\nu+1)^{-1} \sqrt{\log N} \right) \\
&\leq NP^{-1} \exp \left(-c \frac{\sqrt{\log N}}{\log(K+1)} \right).
\end{aligned}$$

Since, by (3.24), K satisfies the inequality

$$(13.9) \quad K \log_2 N < c \sqrt{\log N} / \log K.$$

From (13.9) we have, finally

$$(13.10) \quad (\log R)^{4K} |\mathcal{E}_K(N)| \leq P^{-1} N \exp \left(-c \frac{\sqrt{\log N}}{\log(K+1)} \right).$$

So, our task is reduced to the evaluation of $\tilde{T}_R$ which is very similar to $\mathcal{T}_R^*$ in (6.22). Due to the more general treatment of $\mathcal{T}_R^*$ in Section 5 than needed, the crucial part, the error analysis will remain the same. The difference will be only the fact that we have now $\varphi(a_1 a_2 a_{12})$ in the denominator in (13.7) in place of $a_1 a_2 a_{12}$. Therefore $\frac{\nu_i(p)}{p^{1+s_i}}$ has to be replaced by $\frac{\nu_i(p)}{(p-1)p^{s_i}}$ in the definition of $F(s_1, s_2)$ and $G(s_1, s_2)$ in (6.5) and (6.11) (where $s_i = s_1, s_2$ or $s_3 = s_1 + s_2$). However, factors of type $(1 - p^{-(1+s_i)})$ remain unchanged, since they arise from the zeta-factors.

Summarizing our results above we have

(13.11)

$$\tilde{S}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, \tilde{a}, h_0) = \frac{N}{\varphi(P)} \mathcal{T}_R(\ell_1, \ell_2; \mathcal{H}_1, \mathcal{H}_2) + O \left(\frac{N}{P} \exp \left(-\frac{c \sqrt{\log N}}{\log_2 N} \right) \right)$$

where

$$(13.12) \quad \mathcal{T}_R(\ell_1, \ell_2, \mathcal{H}_1, \mathcal{H}_2) := \frac{1}{(2\pi i)^2} \int \int_{(1)(1)} F(s_1, s_2) \frac{R^{s_1}}{s_1^{K+\ell_1+1}} \frac{R^{s_2}}{s_2^{K+\ell_2+1}} ds_1 ds_2,$$

$$(13.13) \quad F(s_1, s_2) := \prod_{p>V} \left(1 - \frac{\nu_1(p)}{(p-1)p^{s_1}} - \frac{\nu_2(p)}{(p-1)p^{s_2}} + \frac{\nu_3(p)}{(p-1)p^{s_3}} \right)$$

and for this paragraph we have with notation (6.2) ($i = 1, 2$)

$$(13.14) \quad \nu_i(p) = \nu_p^*(\mathcal{H}_i) = \nu_p(\mathcal{H}_i^0) - 1, \quad \nu_3(p) = \bar{\nu}_p((\mathcal{H}_1 \bar{\cap} \mathcal{H}_2)^0) - 1.$$

To factor out the dominant zeta-factors we write now, in place of (6.8)

$$(13.15) \quad F(s_1, s_2) = G_{\mathcal{H}_1, \mathcal{H}_2}(s_1, s_2) \frac{\zeta(1 + s_1 + s_2)^{|\mathcal{H}_1 \cap \mathcal{H}_2|^0 - 1}}{\zeta(1 + s_1)^{|\mathcal{H}_1^0| - 1} \zeta(1 + s_2)^{|\mathcal{H}_2^0| - 1}}$$

and define accordingly G_i ($1 \leq i \leq 4$) as in (6.9)–(6.11) with

$$(13.16) \quad a = |\mathcal{H}_1^0| - 1, \quad b = |\mathcal{H}_2^0| - 1, \quad d = |(\mathcal{H}_1 \cap \mathcal{H}_2)^0| - 1,$$

and with $\nu_i(p)p^{-s_i}/(p-1)$ in place of $\nu_i(p)p^{-1-s_i}$.

Similarly to Section 9 of [7] by symmetry we have to consider three cases:

Case 1. $h_0 \notin \mathcal{H} \iff a = K, b = K, d = r$.

Case 2. $h_0 \in \mathcal{H}_1 \setminus \mathcal{H}_2 \iff a = K - 1, b = K, d = r$.

Case 3. $h_0 \in \mathcal{H}_1 \cap \mathcal{H}_2 \iff a = K - 1, b = K - 1, d = r - 1$.

(Cases 1 and 3 are basically the same.)

Since the results of the previous section are more general, they apply to the error analysis here and we only have to evaluate $G(0, 0)$ in Cases 1–3. Similarly to Section 9 of [7] we have by (13.14)

$$(13.17) \quad \nu_1(p) + \nu_2(p) - \nu_3(p) = \nu_p(\mathcal{H}_1^0) + \nu_p(\mathcal{H}_2^0) - \bar{\nu}_p(\mathcal{H}_1^0 \bar{\cap} \mathcal{H}_2^0) - 1 = \nu_p(\mathcal{H}^0) - 1,$$

$$(13.18) \quad a + b - d = |\mathcal{H}^0| - 1.$$

Hence, from the analogies of (6.9)–(6.11) we have now

$$(13.19) \quad G_1(0, 0) = \prod_{p \leq V} \left(1 - \frac{1}{p}\right)^{-(|\mathcal{H}^0| - 1)} = \left(\frac{P}{\varphi(P)}\right)^{|\mathcal{H}^0| - 1},$$

$$(13.20) \quad \begin{aligned} G_4(0, 0) &= \prod_{p > V} \left(1 - \frac{\nu_p(\mathcal{H}^0) - 1}{p - 1}\right) \left(\frac{p}{p - 1}\right)^{|\mathcal{H}^0| - 1} = \\ &= \prod_{p > V} \left(\frac{p - \nu_p(\mathcal{H}^0)}{p}\right) \cdot \left(1 - \frac{1}{p}\right)^{-|\mathcal{H}^0|} := \tilde{\mathfrak{S}}_V(\mathcal{H}^0). \end{aligned}$$

Taking into account the term $\varphi(P)$ in the denominator in (13.11) we obtain

$$(13.21) \quad \frac{G(0, 0)}{\varphi(P)} = \frac{1}{P} \prod_{p \leq V} \left(1 - \frac{1}{p}\right)^{-|\mathcal{H}^0|} \tilde{\mathfrak{S}}_V(\mathcal{H}^0).$$

Further we have by the comparison of (13.12), (13.15) and (6.22)

$$(13.22) \quad u = K + \ell_1 - a = K + 1 - |\mathcal{H}_1^0| + \ell_1, \quad v = K + 1 - |\mathcal{H}_2^0| + \ell_2, \quad d = |\mathcal{H}_1^0 \cap \mathcal{H}_2^0| - 1.$$

The evaluation (11.5) of the crucial integral I defined in (6.22) yields in our case (13.11)–(13.15) the relation

$$(13.23) \quad \begin{aligned} \tilde{S}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, \tilde{a}, h_0) &= \\ &= N \frac{G(0, 0)}{\varphi(P)} \frac{\binom{v+u}{u} (\log R)^{d+v+u}}{(d+v+u)!} \left(1 + O\left(\frac{K \bar{d}^* \log_2 N}{\log R}\right)\right) \\ &\quad + O\left(\frac{N}{\varphi(P)} e^{-c\sqrt{\log N}}\right). \end{aligned}$$

Let us observe that on the right-hand side the residue class $\tilde{a}$ does not appear at all. Therefore we can add this together for all $|A(\mathcal{H}^0)|$ regular residue classes $\tilde{a} \pmod{P}$ with respect to $\mathcal{H}^0$ and P , since the contribution of those with $(\tilde{a} +$

$h_0, P) > 1$ is zero, as mentioned after (13.3). Taking into account the trivial relations (3.15)–(3.16) for $\mathcal{H}^0$ in place of $\mathcal{H}$ we obtain from (13.21)

$$(13.24) \quad \sum_{\tilde{a} \in A(\mathcal{H}^0)} \frac{G(0,0)}{\varphi(P)} = \frac{|A(\mathcal{H}^0)|}{P} \prod_{p \leq V} \left(1 - \frac{1}{p}\right)^{-|\mathcal{H}^0|} \tilde{\mathfrak{S}}_V(\mathcal{H}^0) \\ = \prod_{p \leq V} \left(1 - \frac{\nu_p(\mathcal{H}^0)}{p}\right) \left(1 - \frac{1}{p}\right)^{-|\mathcal{H}^0|} \cdot \tilde{\mathfrak{S}}_V(\mathcal{H}^0) = \mathfrak{S}(\mathcal{H}^0).$$

Inserting this into (13.23) we obtain by (13.11)

$$(13.25) \quad \tilde{\tilde{S}}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, h_0) := \sum_{\tilde{a} \in A(\mathcal{H})} \tilde{S}_R(N; \mathcal{H}_1, \mathcal{H}_2, \ell_1, \ell_2, P, \tilde{a}, h_0) \\ = N \frac{\binom{v+u}{u} (\log R)^{d+v+u} \mathfrak{S}(\mathcal{H}^0)}{(d+v+u)!} \left(1 + O\left(\frac{K \bar{d}^* \log_2 N}{\log R}\right)\right) \\ + O\left(N \exp\left(-c \min\left(\sqrt{\log R}, \frac{\sqrt{\log N}}{\log_2 N}\right)\right)\right).$$

Now, from a brief examination of the values of the parameters a, b, d in Cases 1, 2, 3 (after (13.16)) and (13.22), we see that

$$(13.26) \quad \frac{\binom{v+u}{u} (\log R)^{d+v+u}}{(d+v+u)!} = C_R(\ell_1, \ell_2, \mathcal{H}_1, \mathcal{H}_2, h_0) \cdot \frac{\binom{\ell_1+\ell_2}{\ell_1} (\log R)^{r+\ell_1+\ell_2}}{(r+\ell_1+\ell_2)!}.$$

The relations (13.25)–(13.26) prove Theorem 5.

14. THE SUM OF THE SINGULAR SERIES $\mathfrak{S}(\mathcal{H})$

Let

$$(14.1) \quad B_{\mathcal{A}}(k) = B(k) = \sum_{|\mathcal{H}|=k, \mathcal{H} \subseteq \mathcal{A}} \mathfrak{S}(\mathcal{H}),$$

where all sets $\mathcal{H} = \{h_1, h_2, \dots, h_k\} \subseteq \mathcal{A} \subseteq [1, N]$ are counted with $k!$ multiplicity according to all possible permutations of h_i , and $|\mathcal{A}| = h$.

By Gallagher's theorem [5] we have for fixed k and $\mathcal{A} = [1, h]$ as $h \rightarrow \infty$

$$(14.2) \quad B_{\mathcal{A}}(k) = h^k (1 + O_{k,\varepsilon}(h^{-\frac{1}{2}+\varepsilon})).$$

This is not uniform in k but up to some level $k \leq f(h)$ one could still show $B_{\mathcal{A}}(k) \sim h^k$. However, we will use here a completely different approach. We do not prove (14.2), just (see Lemma 16) the weaker relation that $B_{\mathcal{A}}(k)/h^k$ is, apart from a factor $1+o(1)$, non-decreasing as a function of k , at least as long as $k = o(h/\log h)$. This result is fortunately completely sufficient for our purposes.

Further, our method is much more general and works for any set $\mathcal{A}$ with $\mathcal{A} \subseteq [1, N]$, $|\mathcal{A}| = h$.

We remark that the asymptotic $B_{\mathcal{A}}(k) \sim h^k$ is probably not true if $\mathcal{A}$ is arbitrary and even for $\mathcal{A} = [1, h]$ it might fail if k is as large as $h/(\log h)^C$.

Let c be an arbitrary small constant, h, z, N and Z sufficiently large,

$$(14.3) \quad k \leq \log N, \quad h^2 \leq z = \log^5 N, \\ Z = P(z) = \prod_{p \leq z} p, \quad Y = Y_z = \prod_{p \leq z} \left(1 - \frac{1}{p}\right)^{-1} \sim e^\gamma \log z,$$

$$(14.4) \quad Q := Q_z := \{n; (n, P(z)) = 1\}, \quad M := \sum_{1 \leq n \leq Z, n \in Q} 1 = \frac{Z}{Y}.$$

Then we have for a fixed set $\mathcal{H}$ consisting of k distinct elements $h_i \in [1, N]$, similarly to Section 6, the density of z -quasi-prime tuples of pattern $\mathcal{H}$, using (6.6):

$$\begin{aligned}
 (14.5) \quad R(\mathcal{H}) &:= \frac{1}{Z} \sum_{\substack{i=1 \\ P_{\mathcal{H}(i)} \in Q}}^Z 1 = \prod_{p \leq z} \left(1 - \frac{\nu_p(\mathcal{H})}{p}\right) = Y^{-k} \prod_{p \leq z} \left(1 - \frac{\nu_p(\mathcal{H})}{p}\right) \left(1 - \frac{1}{p}\right)^{-k} \\
 &= Y^{-k} \mathfrak{S}(\mathcal{H}) \exp \left(O \left(k \sum_{\substack{p > z \\ p | \Delta}} \frac{1}{p} + k^2 \sum_{\substack{p > z \\ p \nmid \Delta}} \frac{1}{p^2} \right) \right) \\
 &= Y^{-k} \mathfrak{S}(\mathcal{H}) \exp \left(O \left(k \sum_{p | \Delta} \frac{\log p}{z \log z} + \frac{k^2}{z \log z} \right) \right) \\
 &= Y^{-k} \mathfrak{S}(\mathcal{H}) \exp \left(O \left(\frac{k^3 \log N}{z \log z} + \frac{k^2}{z \log z} \right) \right) = Y^{-k} \mathfrak{S}(\mathcal{H}) \left(1 + O \left(\frac{1}{\log N} \right) \right),
 \end{aligned}$$

uniformly in k, h, z, N satisfying (14.3), if c is fixed. Let further

$$(14.6) \quad S_{\mathcal{A}}^*(k) := \frac{1}{h^k} \sum_{|\mathcal{H}|=k, \mathcal{H} \subset \mathcal{A}} \mathfrak{S}(\mathcal{H}) = \frac{B_{\mathcal{A}}(k)}{h^k}.$$

Lemma 16. *If $k < \varepsilon(h)h / \log_2 N$, then*

$$(14.7) \quad S_{\mathcal{A}}^*(k+1) \geq S_{\mathcal{A}}^*(k) \left(1 + O(\varepsilon(h)) + O \left(\frac{1}{\log N} \right) \right).$$

Proof. For $i \in [1, Z]$ let

$$(14.8) \quad f_i = \sum_{\substack{j \\ i+a_j \in Q}} 1, \quad b_i = b_i(k) = f_i(f_i - 1) \dots (f_i - k + 1).$$

Then $b_i(k)$ is the number of all k -tuples of z -quasiprimes of type $i + a_{j_\nu}$, $a_{j_\nu} \in \mathcal{A}$ ($\nu = 1, \dots, k$, $1 \leq j_\nu \leq h$, j_ν distinct), calculated with all $k!$ permutations, while f_i is the number of z -quasiprimes of the form $i + a_j$. We have obviously for every pair $i, j \in [1, h]$

$$(14.9) \quad f_i \geq f_j \Leftrightarrow b_i \geq b_j,$$

therefore

$$(14.10) \quad \frac{1}{Z} \sum_{i=1}^Z b_i f_i \geq \frac{\sum_{i=1}^Z f_i}{Z} \frac{\sum_{i=1}^Z b_i}{Z}.$$

The above formula follows from

$$(14.11) \quad 2 \left(Z \sum_{i=1}^Z b_i f_i - \sum_{i=1}^Z f_i \sum_{i=1}^Z b_i \right) = \sum_{i=1}^Z \sum_{j=1}^Z (f_i - f_j)(b_i - b_j) \geq 0.$$

We have further $b_i(k+1) = b_i(k)(f_i - k) = b_i f_i - k b_i$ and by calculating in two different ways how many times all pairs $i, \mathcal{H}$ ($|\mathcal{H}| = k$) satisfy the relation $P_{\mathcal{H}}(i) \in Q$ we obtain

$$(14.12) \quad Z^{-1} \sum_{i=1}^Z b_i(k) = Z^{-1} \sum_{i=1}^Z \sum_{\substack{|\mathcal{H}|=k \\ P_{\mathcal{H}}(i) \in Q}} 1 = Z^{-1} \sum_{|\mathcal{H}|=k} \sum_{\substack{i=1 \\ P_{\mathcal{H}}(i) \in Q}}^Z 1 = \sum_{|\mathcal{H}|=k} R(\mathcal{H}),$$

while

$$(14.13) \quad \frac{1}{Z} \sum_{i=1}^Z f_i = \frac{hM}{Z} = \frac{h}{Y}.$$

Thus (14.10) and (14.13) imply by $b_i f_i = b_i(k+1) + k b_i$ that

$$(14.14) \quad \frac{1}{Z} \sum_{i=1}^Z b_i(k+1) + k \cdot \frac{1}{Z} \sum_{i=1}^Z b_i(k) \geq \frac{h}{Y} \cdot \frac{1}{Z} \sum_{i=1}^Z b_i(k).$$

Hence, using (14.12), we obtain

$$(14.15) \quad \sum_{|\mathcal{H}|=k+1} R(\mathcal{H}) \geq \left(\frac{h}{Y} - k \right) \sum_{|\mathcal{H}|=k} R(\mathcal{H}).$$

Multiplying by Y^{k+1} on both sides, we obtain by (14.5)

$$(14.16) \quad \sum_{|\mathcal{H}|=k+1} \mathfrak{S}(\mathcal{H}) \geq h \left(1 + O\left(\frac{kY}{h}\right) + O\left(\frac{1}{\log N}\right) \right) \sum_{|\mathcal{H}|=k} \mathfrak{S}(\mathcal{H}).$$

Now dividing by h^{k+1} on both sides we obtain (14.7) by $Y \ll \log_2 N$.

15. PROOF OF THEOREM 1

Theorems 4 and 5 allow us to express the quantity $S'_R(N, K, \ell, P)$ in (3.20) in terms of

$$(15.1) \quad S_{\mathcal{A}}^*(k) = S^*(k) := \frac{B_{\mathcal{A}}(k)}{h^k} := \frac{1}{h^k} \sum_{|\mathcal{H}|=k, \mathcal{H} \subset \mathcal{A}} \mathfrak{S}(\mathcal{H}),$$

where we consider two sets $\mathcal{H}$ and $\mathcal{H}'$ different if they contain the same elements in different permutations. The value of the parameter k will be between K and $2K+1$ since in the application the sum (15.1) will refer to sums of type $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2$, $|\mathcal{H}_i| = K$, or to $\mathcal{H}^0$.

The derivation of the proof of Theorem 1 from our present Theorems 4 and 5 will be nearly the same as that of the main result (Theorem 3) of [7] from Propositions 1 and 2 in [7], which appears in Section 10 of [7], so we will be brief. Although the restrictions for K and h will be quite different here, nearly everything will be valid without any change in the present case. Our analysis refers now for the case $\nu = 1$ of Section 10 in [7].

Let us choose, somewhat differently from [7],

$$(15.2) \quad R = (3N)^\Theta = (3N)^{\frac{1}{4}-\xi}, \quad \xi = c/\sqrt{\log N}, \quad V = \sqrt{\log N}$$

$$(15.3) \quad K = 16(\ell+1)^2 = 16\varphi^{-2} \iff \ell+1 = \varphi^{-1} = \sqrt{K}/4$$

$$(15.4) \quad x = \frac{K}{100} = \frac{\log R}{h} \iff h = \frac{100 \log R}{K} \left(\sim \frac{25 \log N}{K} \right),$$

$$(15.5) \quad r_0 = (1-2\varphi)K, \quad r_1 = (1-\varphi)K$$

$$(15.6) \quad f(r) = \binom{K}{r}^2 \frac{x^r}{(r+1) \dots (r+2\ell)}, \quad \bar{r}^* = \max(\sqrt{K}, K-r), \quad t(r) = \frac{\bar{r}^*}{\varphi K}$$

and suppose that our crucial parameter K satisfies

$$(15.7) \quad K \leq c_0 \frac{\sqrt{\log N}}{\log_2^2 N},$$

with a sufficiently small (explicitly calculable) absolute constant c_0 .

In the course of the proof of our present Theorem 1 (similar to Section 10 of [7]) a very important role is played by the fact that although the sums evaluated in Theorems 4 and 5 depend on the actual choice of $\mathcal{H}_1$ and $\mathcal{H}_2$, the asymptotic formulas for them depends just on the set $\mathcal{H} = \mathcal{H}_1 \cup \mathcal{H}_2$ and on the size of $\mathcal{H}_1 \cap \mathcal{H}_2$. On the other hand the size of the error terms may depend on the actual choice of $\mathcal{H}_1, \mathcal{H}_2$ and $\mathcal{H}$. This dependence is made explicit in our present refined version, at least in the sense that we show an asymptotic which is more precise if $r = |\mathcal{H}_1 \cap \mathcal{H}_2|$ is near $K = |\mathcal{H}_i|$.

We have seen in [7] that taking any given set $\mathcal{H}$ of given size $k = 2K - r \in [K, 2K]$, we can write it in

$$(15.8) \quad (2K - r)! \binom{K}{r}^2 r!$$

ways as the union of two sets $\mathcal{H}_1$ and $\mathcal{H}_2$ of size K , $|\mathcal{H}_1 \cap \mathcal{H}_2| = r$ if we consider sets $\mathcal{H}_i$ and $\mathcal{H}'_i$ different when the permutation of the same elements is different (cf. (10.4) of [7]). Now we can apply Theorems 4 and 5 in order to obtain similarly to Section 10 of [7]

$$(15.9) \quad S'_R(N, K, \ell, P) = \binom{2\ell}{\ell} (\log R)^{2\ell} P_{K,\ell}^*(x)$$

with

$$(15.10) \quad P_{K,\ell}^*(x) \geq \sum_{r=0}^K f(r) S^*(2K - r) \left(1 + O(\eta_2) + x \left(\frac{4K(1 - \frac{\varphi}{2})}{r + 2\ell + 1} - \frac{1}{\Theta} + O(\eta_1) \right) \right)$$

where the error terms arising from Theorems 4, 5 and Lemma 16 are now

$$(15.11) \quad \eta_1 = \frac{Kr^* \log_2 N}{\log N} = \frac{4K^{3/2} t(r) \log_2 N}{\log N}, \quad \eta_2 = \frac{1}{\log_2^3 N},$$

and by our choice of Θ in (15.2) we have

$$(15.12) \quad \frac{1}{\Theta} = 4 + O(\eta_3), \quad \eta_3 = \frac{1}{\sqrt{\log N}}.$$

We will examine the quantity in parenthesis after x in (15.10) which is clearly monotonic in r (apart from the error terms). We have now by (15.3)–(15.6) for $r \leq r_1 = K - \varphi K = K - 4\sqrt{K} \Leftrightarrow t(r) \geq 1$:

$$(15.13) \quad r + 2\ell + 1 < K - t(r)\varphi K + \frac{\sqrt{K}}{2} = K \left(1 - t(r)\varphi + \frac{\varphi}{8} \right)$$

and therefore

$$(15.14) \quad \begin{aligned} & \frac{4(1 - \frac{\varphi}{2})K}{r + 2\ell + 1} - \frac{1}{\Theta} + O(\eta_1) > 4 \left(t(r) - \frac{5}{8} \right) \varphi + O(\eta_1 + \eta_3) \\ & > \frac{16}{\sqrt{K}} \cdot \frac{3}{8} t(r) - \frac{Ct(r)K^{3/2} \log_2 N}{\log N} - \frac{C}{\sqrt{\log N}} > 0 \end{aligned}$$

by $t(r) \geq 1$ and (15.7).

On the other hand, as in (10.24)–(10.25) of [7], the contribution of all terms $r_2 > r_1$ to $P_{K,\ell}^*(x)$ is bounded by

$$(15.15) \quad e^{-\sqrt{K}} f(r_0) \max_{r > r_1} S^*(2K - r),$$

because $f(r)$ quickly decreases for $r > r_1$. (These are the terms where the quantity in parenthesis after x in (15.10) may be negative.) We have, for any $r > r_2$,

$$(15.16) \quad \frac{f(r_2)}{f(r_0)} = \prod_{r_0 < r \leq r_2} \left(\frac{K-r+1}{r} \cdot \frac{\sqrt{K}}{10} \right)^2 \leq \left(\frac{2\varphi K}{K-2\varphi K} \cdot \frac{\sqrt{K}}{10} \right)^{2\varphi K} \\ \leq (0.81)^{8\sqrt{K}} = e^{-1.6\sqrt{K}}.$$

However, all terms $r \leq r_1$ have a positive contribution and that of $r = r_0$ is at least

$$(15.17) \quad f(r_0)S^*(2K-r_0) \left(1 + O\left(\frac{1}{\log_2^3 N} \right) \right).$$

Now the quasi-monotonic property, Lemma 16, implies

$$(15.18) \quad \frac{S^*(2K-r_0)}{\max_{r>r_1} S^*(2K-r)} > e^{-(K-r_0)C/\log_2^3 N} = e^{-\frac{8C\sqrt{K}}{\log_2^3 N}}.$$

Consequently the positive term belonging to r_0 dominates all possibly negative terms belonging to $r > r_1$ and therefore we have

$$(15.19) \quad P_{K,\ell}^*(x) > 0 \iff S'_R(N, K, \ell, P) > 0.$$

This, by (3.20), proves the existence of some $n \in [N+1, 2N]$ with

$$(15.20) \quad \sum_{p=n+a_\nu, a_\nu \in \mathcal{A}} \log p > \log(3N),$$

and thereby the existence of two primes $p', p'' \in [N+1, 3N]$ with

$$(15.21) \quad 0 \neq p'' - p' \in \mathcal{A} - \mathcal{A}.$$

This proves Theorem 1 if we choose K maximal, satisfying the restriction (15.7).

REFERENCES

- [1] E. Bombieri and H. Davenport, Small differences between prime numbers, *Proc. Roy. Soc. Ser. A* **293** (1966), 1–18.
- [2] H. Davenport, *Multiplicative Number Theory*, Second Edition, Revised by Hugh L. Montgomery, Springer, Berlin, Heidelberg, New York, 1980.
- [3] P. D. T. A. Elliott and H. Halberstam, A conjecture in prime number theory, *Symposia Mathematica* **4** (INDAM, Rome, 1968/69) 59–72, Academic Press, London.
- [4] P. Erdős, The difference of consecutive primes, *Duke Math. J.* **6** (1940), 438–441.
- [5] P. X. Gallagher, On the distribution of primes in short intervals, *Mathematika* **23** (1976), 4–9.
- [6] D. M. Goldfeld and A. Schinzel, On Siegel’s zero, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **2** (1975), 571–583.
- [7] D. A. Goldston, J. Pintz and C. Y. Yıldırım, Primes in Tuples I, AIM Preprint Series N° 2005-19, <http://aimath.org/preprints.html> (accepted, Annals of Mathematics)
- [8] H. Halberstam and H.-E. Richert, *Sieve methods*, Academic Press, London, New York, 1975.
- [9] G. H. Hardy and J. E. Littlewood, Some problems of ‘Partitio Numerorum’; III: On the expression of a number as a sum of primes, *Acta Math.* **44** (1922), 1–70.
- [10] D. R. Heath-Brown, Prime twins and Siegel zeros, *Proc. London Math. Soc.* (3) **47** (1983), no. 2, 193–224.
- [11] D. R. Heath-Brown, Almost-prime k -tuples, *Mathematika* **44** (1997), 245–266.
- [12] M. N. Huxley, Small differences between consecutive primes II, *Mathematika* **24** (1977), 142–152.
- [13] A. Ivíč, *The Riemann Zeta-Function*, John Wiley & Sons, New York, 1985.
- [14] H. Maier, Small differences between prime numbers, *Michigan Math. J.* **35** (1988), 323–344.
- [15] H. L. Montgomery, *Topics in Multiplicative Number Theory*, Lecture Notes in Mathematics, Springer, Berlin, Heidelberg, New York, 1971.
- [16] J. Pintz, Elementary methods in the theory of $\mathcal{L}$ -functions II, On the greatest real zero of a real $\mathcal{L}$ -function, *Acta Arith.* **31** (1976), 273–289.
- [17] J. Pintz, Elementary methods in the theory of $\mathcal{L}$ -functions VIII, Real zeros of real $\mathcal{L}$ -functions, *Acta Arith.* **33** (1977), 89–98.

- [18] J. Pintz, Very large gaps between consecutive primes, *J. Number Th.* **63** (1997), 286–301.
- [19] J. Pintz, Approximations to the Goldbach and Twin Prime Problem and Gaps Between Consecutive Primes, *Advanced Studies in Pure Mathematics* **43** (2006), International Conference in Probability and Number Theory, Kanazawa, 2005, 1–40.
- [20] Polignac, A. de, Six propositions arithmologiques sur les nombres premiers, *Nouv. Ann. Math.* **7** (1849), 423–429.
- [21] R. A. Rankin, The difference between consecutive prime numbers. II, *Proc. Cambridge Philos. Soc.* **36** (1940), 255–266.
- [22] G. Ricci, Sull’andamento della differenza di numeri primi consecutivi, *Riv. Mat. Univ. Parma* **5** (1954), 3–54.
- [23] E. C. Titchmarsh, *The theory of the Riemann zeta-function*, Second edition, Edited and with a preface by D. R. Heath-Brown, The Clarendon Press, Oxford University Press, New York, 1986.
- [24] K. Soundararajan, Small gaps between prime numbers: the work of Goldston-Pintz-Yıldırım, *Bull. Amer. Math. Soc.* **44**, Number 1, January 2007, 1–18.

D. A. Goldston
 Department of Mathematics
 San Jose State University
 San Jose, CA 95192, USA
 e-mail: goldston@math.sjsu.edu

J. Pintz
 Rényi Mathematical Institute of the Hungarian Academy of Sciences
 Budapest, Reáltanoda u. 13-15
 H-1053
 Hungary
 e-mail: pintz@renyi.hu

C. Y. Yıldırım
 Department of Mathematics
 Boğaziçi University
 Bebek
 Istanbul 34342
 Turkey
 & Feza Gürsey Enstitüsü
 Çengelköy
 Istanbul, P.K. 6,
 81220 Turkey
 e-mail: yalciny@boun.edu.tr