

Name: \_\_\_\_\_

Section: \_\_\_\_\_

**Instructions:** You have **3 hours** to complete this exam. You should work alone, without access to the textbook or class notes. You may not use a calculator. Do not discuss this exam with anyone except your instructor.

This exam consists of 9 questions. Except for the first problem (multiple choice), you must show your work to receive full credit. Be sure to **indicate your final answer clearly** for each question.

**Note:** If you use a major theorem (such as Green's, Stokes', or Gauss' Divergence theorem), you must indicate it. Points will be deducted for failure to indicate the use of a major theorem. **You must clearly indicate each time you use such a theorem.**

Before turning in the exam, be sure to:

- Staple your exam with this cover sheet on top,
- Pledge your exam,
- Write your name and section number above.

The exam is due by **Wednesday, May 10, 4 p.m.** Good luck!

**Pledge:**

Problem	Value	Score
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
Total	90	

1. In the following problem, let  $S$  be a unit disk in the plane  $z = 5$ , centered at  $(0, 0, 5)$ , and oriented upward. Let  $C$  be a straight path from  $(2, 2, 2)$  to  $(0, 0, 0)$ . Also, let

$$f(x, y, z) = xy^2z^3,$$

$$\mathbf{F}(x, y, z) = -y\mathbf{i} + x\mathbf{j},$$

$$\mathbf{G}(x, y, z) = 3\mathbf{i} + \mathbf{j} + 2\mathbf{k}.$$

Now, for each of the given quantities, decide if it is **positive**, **negative**, or **zero**. (You do not need to justify your answers. No partial credit will be given.)

(a)  $\iint_S \nabla \times \mathbf{F} \cdot d\mathbf{S}.$

(b)  $\int_C \nabla f \cdot ds.$

(c)  $\iint_S \mathbf{F} \cdot d\mathbf{S}.$

(d)  $\iint_S \mathbf{G} \cdot d\mathbf{S}.$

(e)  $\int_{\partial S} \nabla f \cdot ds.$

2. (a) Find all critical points of the function  $g(x, y) = 2y^3 - 6y + x^2$  in  $\mathbb{R}^2$  and classify them.  
 (b) Let  $B$  be the unit ball in  $\mathbb{R}^3$ , i.e. the set of points  $(x, y, z)$  satisfying  $x^2 + y^2 + z^2 \leq 1$ . Let  $f(x, y, z) = 2x + 4y + 6z$ . Find the minimum and maximum values of  $f$  restricted to  $B$ .

3. (a) Let  $f(x, y)$  be a  $C^1$  function whose domain is the unit disk in the  $xy$ -plane, such that  $f(x, y) \geq 0$  everywhere. Suppose that the level set  $f(x, y) = 0$  is exactly the unit circle. Let  $S$  be the graph of  $f(x, y)$ , oriented upward, and let

$$\mathbf{F}(x, y, z) = \text{curl}(x^2 - z, e^z + 2x, \pi).$$

Determine  $\iint_S \mathbf{F} \cdot d\mathbf{S}.$

- (b) Let  $g(x, y)$  be a  $C^1$  function whose domain is the unit disk in the  $xy$ -plane, such that  $g(x, y) \leq 0$  everywhere. Suppose that the level set  $g(x, y) = 0$  is exactly the unit circle. Let  $T$  be the graph of  $f(x, y)$ , oriented downward.

Determine  $\iint_T \mathbf{F} \cdot d\mathbf{S}.$

4. Evaluate  $\int_0^{\sqrt[7]{3^3}} \int_{\sqrt[3]{x}}^{\sqrt{3}} x(\sqrt{1+y^7}) dy dx.$  Hint: Consider the region of integration.

5. Let  $W$  be the region in space under the graph of

$$f(x, y) = (\cos y) \exp(1 - \cos 2x) + xy$$

over the region in the  $xy$ -plane bounded by the line  $y = 2x$ , the  $x$  axis, and the line  $x = \pi/4$ .

(a) Find the volume of  $W$ .

(b) Let  $\mathbf{F} = 5x\mathbf{i} + 5y\mathbf{j} + 5z\mathbf{k}$  be the velocity field of a fluid in space. Calculate the flux of  $\mathbf{F}$  through the boundary  $\partial W$  of  $W$ , where  $W$  is the region from (a).

6. Let  $S$  be a surface in  $\mathbb{R}^3$  given as follows.  $S$  is the portion of the cylinder  $x^2 + y^2 = 9$  lying above  $z = 0$ , below the graph of  $z = \sqrt{x^2 + (y - 3)^2}$ , and with  $y \leq 0$ .

(a) Set up, **but do not evaluate**, an integral giving the surface area of  $S$ .

(b) Suppose  $S$  is made of a thin metal whose mass density at any point is given by the function  $f(x, y, z) = \sqrt{1 - \frac{y}{3}}$ . Find the total mass of  $S$ .

7. Let  $g(x, y) = xe^y$ .

(a) Compute the second-order Taylor formula for  $g$  around  $(3, 0)$ .

(b) Approximate  $2.9e^{0.1}$ .

(c) Compute the directional derivative of  $g$  based at  $(3, 0)$  in the direction of fastest increase.

8. Let  $\mathbf{F}(x, y) = 2xy\mathbf{i} + x^2\mathbf{j}$ . Show that the line integral of  $\mathbf{F}$  around the triangle  $T$  (oriented counterclockwise) with vertices  $(0, 0)$ ,  $(0, 1)$  and  $(1, 1)$  is zero in the following three ways:

(a) parameterizing  $T$  and evaluating the integral directly,

(b) showing  $\mathbf{F}$  is a gradient vector field and explaining why the integral is zero, and

(c) using Green's Theorem.

9. Compute the flux of the vector field

$$\mathbf{G}(x, y, z) = (xy^2, yz^2 + y, zx^2 + 1)$$

through the unit sphere, oriented outward.