

1. Let $f(x, y, z)$ be a scalar function, and let $\mathbf{F}(x, y, z)$ be a vector field. (Assume both f and \mathbf{F} have continuous partial derivatives of all orders.) Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be vectors in \mathbb{R}^3 .

(a) $\text{curl grad } f = \mathbf{0}$.

Solution: **True**.

(b) $\text{div grad } f = 0$.

Solution: **False**. For example if $f(x, y, z) = x^2$, then $\text{grad } f = (2x, 0, 0)$ and $\text{div grad } f = 2$.

(c) $\text{div curl } \mathbf{F} = 0$.

Solution: **True**.

(d) Let C be an oriented curve. The path integral of f along C does not change when the orientation of C is reversed.

Solution: **True**.

(e) Let C be an oriented curve. The line integral of \mathbf{F} along C does not change when the orientation of C is reversed.

Solution: **False**. The integral changes by a minus-sign.

(f) The expression $\mathbf{u} \cdot \mathbf{v}$ is a vector.

Solution: **False**.

(g) The expression $\mathbf{u} \times \mathbf{v}$ is a vector.

Solution: **True**.

(h) The expression $(\mathbf{v} \cdot \mathbf{w})\mathbf{u}$ is a vector.

Solution: **True**. First note that $(\mathbf{v} \cdot \mathbf{w})$ is a scalar, so $(\mathbf{v} \cdot \mathbf{w})\mathbf{u}$ is the scalar product of the vector \mathbf{u} with the scalar $(\mathbf{v} \cdot \mathbf{w})$.

(i) Let S be an oriented surface. The quantity $\iint_S \mathbf{F} \cdot d\mathbf{S}$ is a vector.

Solution: **False**.

(j) Let S be an oriented surface. The quantity $\iint_S f dS$ is a vector.

Solution: **False**.

2. Let $\mathbf{F}(x, y, z) = \left(\frac{-y}{x^2 + y^2}, \frac{x}{x^2 + y^2}, 0 \right)$.

(a) Show $\text{curl } \mathbf{F} = (0, 0, 0)$.

(b) Let C be the unit circle in the xy -plane, oriented **clockwise**. Evaluate $\int_C \mathbf{F} \cdot ds$.

(c) Using your answer from (b), explain why \mathbf{F} is not a gradient field, even though $\text{curl } \mathbf{F} = (0, 0, 0)$.

Solution:

(a) This is a straight-forward computation:

$$\begin{aligned} \text{curl } \mathbf{F} &= \det \begin{pmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{-y}{x^2+y^2} & \frac{x}{x^2+y^2} & 0 \end{pmatrix} \\ &= -\mathbf{i} \frac{\partial}{\partial z} \frac{x}{x^2+y^2} + \mathbf{j} \frac{\partial}{\partial z} \frac{-y}{x^2+y^2} + \mathbf{k} \left(\frac{\partial}{\partial x} \frac{x}{x^2+y^2} - \frac{\partial}{\partial y} \frac{-y}{x^2+y^2} \right) \\ &= \mathbf{k} \left(\frac{x^2+y^2-2x^2}{(x^2+y^2)^2} + \frac{x^2+y^2-2y^2}{(x^2+y^2)^2} \right) \\ &= (0, 0, 0). \end{aligned}$$

(b) We first have to parametrize C . We take

$$\mathbf{c}(t) = (\cos(-t), \sin(-t)), t \in [0, 2\pi].$$

(note that we need the $-t$ to ensure that we go clockwise). We then compute

$$\begin{aligned} \int_C \mathbf{F} \cdot ds &= \int_{t=0}^{t=2\pi} \mathbf{F}(\mathbf{c}(t)) \cdot \mathbf{c}'(t) dt \\ &= \int_{t=0}^{t=2\pi} (-\sin(-t), \cos(-t), 0) \cdot (\sin(-t), -\cos(-t), 0) dt \\ &= \int_{t=0}^{t=2\pi} -1 dt = -2\pi. \end{aligned}$$

(Note that one could also choose the usual clockwise parametrization $\mathbf{c}(t) = (\cos(t), \sin(t))$, $t \in [0, 2\pi]$, and then changed the integral by a minus-sign.)

(c) If \mathbf{F} was a gradient field, then any line integral over a closed curve would be zero. But in (b) we saw that this is not the case.

3. The surface S is parameterized by $\Phi(u, v) = (e^u - 2, 2v + 3, 5 + u^2 + v^2)$ with $u, v \in \mathbb{R}$.

(a) Determine the equation of the tangent plane to $(-1, 5, 6) \in S$.

(b) Find all points on S for which the tangent plane is parallel to the xy -plane.

Solution:

(a) We compute

$$\begin{aligned}\mathbf{T}_u &= (e^u, 0, 2u) \\ \mathbf{T}_v &= (0, 2, 2v) \\ \mathbf{T}_u \times \mathbf{T}_v &= (-4u, -2ve^u, 2e^u).\end{aligned}$$

Now note that $\Phi(u, v) = (-1, 5, 6)$ for $u = 0, v = 1$ (which can be seen by looking at the first two coordinates).

For $u = 0, v = 1$ we have

$$\mathbf{T}_u \times \mathbf{T}_v = (0, -2, 2).$$

This is the normal vector to the tangent plane at $(-1, 5, 6)$. The equation of the tangent plane is therefore given by

$$(x - (-1), y - 5, z - 6) \cdot (0, -2, 2) = 0$$

which simplifies to

$$-2y + 2z - 2 = 0.$$

(b) First note that the tangent plane is parallel to the xy -plane if the z -components of \mathbf{T}_u and \mathbf{T}_v are zero. But this means that $2u = 0$ and $2v = 0$. So the only possibility is $u = 0, v = 0$. The corresponding point on S is $\Phi(0, 0) = (-1, 3, 5)$.

4. Let $f(x, y) = \frac{1}{3}x^3 + y\sqrt{2} + 3$, and let D be the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 1)$. Let S be the surface given by the graph of $f(x, y)$ over D .

(a) Find a parametrization of S .

(b) Compute $\iint_S 4x^2 dS$.

Solution:

(a) The parametrization is given by

$$\Phi(x, y) = (x, y, \frac{1}{3}x^3 + y\sqrt{2} + 3)$$

where the domain for x, y is given by D , i.e. by the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 1)$.

(b) Let's first compute $\|\mathbf{T}_x \times \mathbf{T}_y\|$:

$$\begin{aligned}\mathbf{T}_x &= (1, 0, x^2) \\ \mathbf{T}_y &= (0, 1, \sqrt{2}) \\ \mathbf{T}_x \times \mathbf{T}_y &= (-x^2, -\sqrt{2}, 1) \\ \|\mathbf{T}_x \times \mathbf{T}_y\| &= \sqrt{3 + x^4}.\end{aligned}$$

We compute

$$\iint_S 4x^2 dS = \int_{x=0}^{x=1} \int_{y=0}^{y=x} 4x^2 \sqrt{3 + x^4} dy dx = \int_{x=0}^{x=1} 4x^3 \sqrt{3 + x^4} dx.$$

Now let $u = 3 + x^4$ and we get

$$\int_{x=0}^{x=1} 4x^3 \sqrt{3+x^4} dx = \int_{u=3}^{u=4} \sqrt{u} du = \left[\frac{2}{3} u^{\frac{3}{2}} \right]_{u=3}^{u=4} = \frac{2}{3} (4^{\frac{3}{2}} - 3^{\frac{3}{2}}).$$

5. Consider the solid hemisphere formed by taking the portion of the unit ball with $y \geq 0$. Let S be the **surface** of this region (so that S is a hemisphere, together with a flat ‘base’ in the xz -plane). Find the flux of the vector field $\mathbf{V}(x, y, z) = -z\mathbf{i} + \mathbf{j} + x\mathbf{k}$ out of the surface S .

You may find the following identity useful: $\sin^2 \alpha = \frac{1}{2}(1 - \cos 2\alpha)$.

Solution: We have to break this problem into two parts. We first integrate over the hemisphere, and then we integrate over the flat base.

For the hemisphere we take the parametrization

$$\Phi(\theta, \phi) = (\cos(\theta) \sin(\phi), \sin(\theta) \sin(\phi), \cos(\phi))$$

with $\theta \in [0, \pi]$ and $\phi \in [0, \pi]$. We compute

$$\begin{aligned} \mathbf{T}_\theta &= (-\sin(\theta) \sin(\phi), \cos(\theta) \sin(\phi), 0) \\ \mathbf{T}_\phi &= (\cos(\theta) \cos(\phi), \sin(\theta) \cos(\phi), -\sin(\phi)) \\ \mathbf{T}_\theta \times \mathbf{T}_\phi &= (-\cos(\theta) \sin^2(\phi), -\sin(\theta) \sin^2(\phi), -\sin(\phi) \cos(\phi)) \\ &= -\sin(\phi) (\cos(\theta) \sin(\phi), \sin(\theta) \sin(\phi), \cos(\phi)). \end{aligned}$$

Note that $\mathbf{T}_\theta \times \mathbf{T}_\phi$ points inward (this can be seen by considering the last line and noticing that $-\sin(\phi)$ is negative). So it has the wrong orientation. We can fix this by putting a minus-sign into the formula. So the flux through the hemisphere is given by

$$\begin{aligned} & - \int_{\theta=0}^{\theta=\pi} \int_{\phi=0}^{\phi=\pi} \mathbf{V}(\Phi(\theta, \phi)) \cdot (\mathbf{T}_\theta \times \mathbf{T}_\phi) d\phi d\theta \\ &= - \int_{\theta=0}^{\theta=\pi} \int_{\phi=0}^{\phi=\pi} (-\cos(\phi), 1, \cos(\theta) \sin(\phi)) \cdot (-\cos(\theta) \sin^2(\phi), -\sin(\theta) \sin^2(\phi), -\sin(\phi) \cos(\phi)) d\phi d\theta \\ &= - \int_{\theta=0}^{\theta=\pi} \int_{\phi=0}^{\phi=\pi} -\sin(\theta) \sin^2(\phi) d\phi d\theta \\ &= - \int_{\theta=0}^{\theta=\pi} \int_{\phi=0}^{\phi=\pi} -\frac{1}{2} \sin(\theta) (1 - \cos(2\phi)) d\phi d\theta \\ &= - \int_{\theta=0}^{\theta=\pi} \left[-\frac{1}{2} \sin(\theta) (\phi - \frac{1}{2} \sin(2\phi)) \right]_{\phi=0}^{\phi=\pi} d\theta \\ &= - \int_{\theta=0}^{\theta=\pi} -\frac{\pi}{2} \sin(\theta) d\theta \\ &= - \left[\frac{\pi}{2} \cos(\theta) \right]_{\theta=0}^{\theta=\pi} \\ &= \pi. \end{aligned}$$

Another way to approach this part of the problem is to see that $x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ is the unit normal vector to the sphere at the point (x, y, z) . Thus $\mathbf{V} \cdot \mathbf{n} = y$. To obtain the flux,

one integrates this (as a scalar surface integral) in spherical coordinates. (It is a little bit faster this way.)

Now let's integrate over the base. The base lies in the xz -plane and the normal vector is $-\mathbf{j}$ (since outside is to the left), since the \mathbf{j} -component of \mathbf{V} is constant one, we see that the flux integral is just -1 times the area of the base, which equals $-\pi$.

To compute the flux over S we have to add up the results from the hemisphere and the base, and we get that the flux equals $\pi - \pi = 0$.

A more formal approach is to find a parametrization again. We could take

$$\Phi(r, \theta) = (r \cos(\theta), 0, r \sin(\theta))$$

where $r \in [0, 1]$ and $\theta \in [0, 2\pi]$. Then

$$\begin{aligned}\mathbf{T}_r &= (\cos(\theta), 0, \sin(\theta)) \\ \mathbf{T}_\theta &= (-r \sin(\theta), 0, r \cos(\theta)) \\ \mathbf{T}_r \times \mathbf{T}_\theta &= (0, -r, 0).\end{aligned}$$

This is the correct orientation, since $(0, -1, 0)$ points outside. So we have:

$$\begin{aligned}\iint \mathbf{V} \cdot d\mathbf{S} &= \int_{r=0}^{r=1} \int_{\theta=0}^{\theta=2\pi} \mathbf{V}(\Phi(r, \theta)) \cdot (\mathbf{T}_r \times \mathbf{T}_\theta) d\theta dr \\ &= \int_{r=0}^{r=1} \int_{\theta=0}^{\theta=2\pi} (-r \sin(\theta), 1, r \cos(\theta)) \cdot (0, -r, 0) d\theta dr \\ &= \int_{r=0}^{r=1} \int_{\theta=0}^{\theta=2\pi} -r d\theta dr = -\pi.\end{aligned}$$

6. Let $\mathbf{c}(t) = (1, -t^2, \cos t)$, $0 \leq t \leq \pi$. Evaluate

$$\int_{\mathbf{c}} \sin z dx - y^2 dy + 3xz dz.$$

Solution: We compute

$$\begin{aligned}\int_{\mathbf{c}} \sin z dx - y^2 dy + 3xz dz &= \int_{t=0}^{t=\pi} \sin(\cos(t)) \frac{d}{dt}(1) - (-t^2)^2 \frac{d}{dt}(-t^2) + 3 \cos(t) \frac{d}{dt}(\cos(t)) dt \\ &= \int_{t=0}^{t=\pi} 2t^5 - 3 \cos(t) \sin(t) dt \\ &= \left[\frac{t^6}{3} + \frac{1}{2} 3 \cos(t)^2 \right]_{t=0}^{t=\pi} \\ &= \frac{\pi^6}{3}.\end{aligned}$$